

Realization of symmetry in the ERG approach to quantum field theory

Yuji IGARASHI¹, Katsumi ITOH¹, and Hidenori SONODA²

¹ *Department of Education, Niigata University, Japan*

² *Physics Department, Kobe University, Japan*

We review the use of the exact renormalization group for realization of symmetry in renormalizable field theories. The review consists of three parts. In part I (§§2,3,4), we start with the perturbative construction of a renormalizable field theory as a solution of the exact renormalization group (ERG) differential equation. We show how to characterize renormalizability by an appropriate asymptotic behavior of the solution for a large momentum cutoff. Renormalized parameters are introduced to control the asymptotic behavior. In part II (§§5–9), we introduce two formalisms to incorporate symmetry: one by imposing the Ward-Takahashi identity, and another by imposing the generalized Ward-Takahashi identity via sources that generate symmetry transformations. We apply the two formalisms to concrete models such as QED, YM theories, and the Wess-Zumino model in four dimensions, and the O(N) non-linear sigma model in two dimensions. We end this part with calculations of the abelian axial and chiral anomalies. In part III (§§10,11), we overview the Batalin-Vilkovisky formalism adapted to the Wilson action of a bare theory with a UV cutoff. We provide a few appendices to give details and extensions that can be omitted for the understanding of the main text. The last appendix is a quick summary for the reader's convenience.

Contents

1. Introduction	3
2. Derivation of the ERG differential equations	9
2.1. Splitting fields into high and low momentum modes	9
2.2. Functional integrals with a source	13
2.3. Polchinski equation	15
2.4. Wetterich equation	17
2.5. Recapitulation	23
2.6. Diagrammatic interpretation	25
2.7. Fermions	30
3. Continuum limits	32
3.1. Perturbative renormalizability	32
3.2. Asymptotic behaviors	36
3.3. Asymptotic behaviors of the Wilson action and the effective average action	40
3.4. Alternative parameterization	41
4. Composite operators	42
4.1. Definition	42
4.2. General examples	45
4.3. Composite operators defined by their asymptotic behaviors	48
4.4. 1PI composite operators	49
5. Realization of symmetry by WT identity	51

5.1. WT identity in the continuum limit	51
5.2. WT identity for the bare action	54
6. Examples	56
6.1. QED	56
6.2. Yang-Mills theories	62
6.3. Wess-Zumino model	67
6.4. $O(N)$ non-linear sigma model	73
6.5. The axial symmetry with a massive propagator	79
7. Realization of symmetry in the antifield formalism	82
7.1. Quantum master equation	82
7.2. Outline of perturbative construction	84
8. Examples	88
8.1. QED	88
8.2. YM theories	90
9. Axial and chiral anomalies	98
9.1. Axial anomaly in QED	98
9.2. Chiral anomaly	101
10. Functional integral approach to the antifield formalism	108
10.1. The Batalin-Vilkovisky antifield formalism	109
10.2. Scale change in the functional integral approach	111
10.3. WT identity and QME	114
10.4. Antifield dependence of the Wilson action	118
10.5. QME and the modified Slavnov-Taylor identity	121
10.6. On the nilpotency without antifields	124
11. Examples: the Wilson actions for QED and Yang-Mills theories	125
11.1. QED	125
11.2. Yang-Mills theories	127
12. Concluding remarks	130
Appendix	
A. Generalized diffusion equations	131
A.1. Derivation of the generalized diffusion equations	131
A.2. Extension to fermions	137
A.3. WT operator for chiral invariance	139
B. Applications of composite operators	141
B.1. Beta functions	141
B.2. Universality	144
C. ERG differential equations for fixed points	146
D. Symmetry of the effective average action	149
E. $O(N)$ linear sigma model	152
F. Quick summary	157

§1. Introduction

This is a review of the **exact renormalization group (ERG)** with an emphasis on the use of ERG for realization of symmetry in renormalizable field theories. In writing this review, we have set two goals. The first is to popularize the ERG formalism among the practitioners of quantum field theory. We believe ERG has enough merits to become part of the shared knowledge of all those who use quantum field theory to think and calculate. The second is to convince the reader of the universal applicability of ERG: if a field theoretical model can have a continuous symmetry, we can formulate it exactly using the ERG formalism. ERG is the second best after a formalism that realizes the symmetry manifestly with no fine-tuning of parameters.

Modern development of the renormalization group started with the work of K. G. Wilson, whose main results on the subject, summarized below, are reviewed in the well known lecture notes 124):

1. He introduced the exact renormalization group as a non-perturbative framework in which we can define the continuum limit of a field theory. Given a theory near criticality, he showed how to construct a massive field theory as a double limit of criticality and infinite distances.
2. He derived an approximate recursion formula that embodies the renormalization group for the real scalar theory in D dimensions, and computed a critical exponent (the anomalous dimension of the squared mass) by solving the formula numerically.
3. He, together with M. E. Fisher, devised the epsilon expansions to compute non-trivial critical exponents in powers of ϵ , the difference of the space dimension from 4 (or the dimension in which the corresponding critical theory is gaussian).

In addition to these main results Wilson introduced the exact renormalization group equations in differential form, the principal tools for the symmetry realization in this review. Many of the results we give in §2 can already be found in §11 of 124). *)**)

It took almost ten years before Wilson's ERG differential equation was taken up by Polchinski in 94). After modifying the equation to a form suitable for perturbation theory, Polchinski applied the equation to prove the perturbative renormalizability of the ϕ^4 theory in 4 dimensions. We base all our discussions of symmetry realization on Polchinski's differential equation, since we are primarily interested in perturbation theory. For those interested in Wilson's original ERG differential equation, we describe it briefly in Appendix C for the case of a real scalar field.

Let us now recall briefly the basic idea behind Wilson's exact renormalization group.***) The starting point is a bare action S_B , a functional of field variables with

*) Wilson also predicted that his ERG differential equations would become the basis for most future work on the renormalization group.

**) Wegner and Houghton introduced an exact renormalization group differential equation independently of Wilson.¹¹⁷⁾ The one-particle-irreducible version of the Wegner-Houghton equation was subsequently derived in 88).

***) The equations in this section are given for the purpose of illustration, and they require modifications to be strictly correct.

an ultraviolet cutoff Λ_0 . The correlation of field variables is given by a functional integral with the weight e^{S_B} :

$$\langle \phi(p_1) \cdots \phi(p_n) \rangle = \int [d\phi] \phi(p_1) \cdots \phi(p_n) e^{S_B[\phi]}. \quad (1.1)$$

By integrating over the field variables with momentum between Λ_0 and $\Lambda < \Lambda_0$, we obtain an equivalent action S_Λ , called the Wilson action, that gives the same correlation functions:

$$\langle \phi(p_1) \cdots \phi(p_n) \rangle = \int [d\phi] \phi(p_1) \cdots \phi(p_n) e^{S_\Lambda[\phi]}. \quad (1.2)$$

The Wilson action S_Λ has the ultraviolet cutoff Λ . As we lower Λ , we generate a flow of equivalent Wilson actions that provide the same correlation functions.^{*)}

In defining the continuum limit of a field theory, we usually look at the correlation functions as we raise $\Lambda_0 \rightarrow \infty$.^{**)} If the theory is renormalizable, we can give appropriate Λ_0 dependence to the parameters of the bare action S_B so that the correlation functions have a limit as $\Lambda_0 \rightarrow \infty$:

$$\lim_{\Lambda_0 \rightarrow \infty} \langle \phi(p_1) \cdots \phi(p_n) \rangle. \quad (1.3)$$

Alternatively, we can look at the Wilson action S_Λ at a fixed Λ . S_B and S_Λ give the same correlation functions, and if S_Λ has a limit as $\Lambda_0 \rightarrow \infty$, the correlation functions must also have a limit. Polchinski used this idea to simplify the proof of the perturbative renormalizability of the ϕ^4 theory.⁹⁴⁾

Polchinski's work was subsequently developed further. We just mention a few as examples. An early attempt was made in 115), 116) to apply Polchinski's differential equation to construct Yang-Mills, chiral, and supersymmetric Yang-Mills theories. The beta functions of renormalized parameters were derived from the Wilson action,⁵⁷⁾ and the proof of renormalizability in 94) was simplified,⁷²⁾ extended,^{70), 71)} and given more rigor.¹⁰⁾

Turning now to the subject of symmetry realization, we call a symmetry **manifest** if it exists in the bare action S_B . An example is a continuous linear symmetry, such as $O(N)$ of N real scalar fields, for which the transformation is linear in the elementary fields. Another example is the gauge symmetry of the lattice gauge theory, for which the field transformation is highly non-linear. Not all symmetry is manifest, though: even if the symmetry cannot be realized manifestly, it may exist in the continuum limit of the theory.

Let us recall that the continuum limit is described fully by the Wilson action. If a theory has symmetry in its continuum limit, the symmetry must be present in the Wilson action. This simple reasoning suggests the possibility of formulating symmetry using the Wilson action in the continuum limit. It is the second goal of this review to explain this formulation. We introduce two identities:

^{*)} The precise definition of the Wilson action is given by (2.23), and the Λ independence of the correlation functions are given by (2.43).

^{**) Here, we describe the more traditional perturbative definition of renormalizability, not the non-perturbative definition given in 124). Hence, no rescaling of space is introduced.}

1. the **Ward-Takahashi identity (WT identity)** (to be discussed in §§5, 6) — this familiar identity is usually given for the bare action in the case of manifest symmetry. In general any continuous symmetry, whether linear or non-linear, can be formulated as the invariance of the Wilson action under an infinitesimal transformation of field variables. The infinitesimal change is highly non-linear due to integration of higher momentum modes, giving rise to a non-trivial jacobian of the transformation. Hence, the Wilson action is invariant only if the jacobian is taken into account. The WT identity for the Wilson action was discussed earlier by Becchi in his seminal work on the ERG approach to YM theories.¹⁷⁾
2. the **quantum master equation** (to be discussed in §§7, 8, and again in §§10, 11) — we introduce classical external sources that generate the symmetry transformation of fields. For gauge theory, this is quite familiar from the Zinn-Justin equation satisfied by the effective action.¹²⁵⁾ The quantum master equation results from an adaptation of the general Batalin-Vilkovisky formalism^{16)*)} to the Wilson action. It is not hard to expect the presence of a rich algebraic structure in this formulation. In fact, for YM theories we must generalize the WT identity to the quantum master equation in order to complete the perturbative proof of the theory's existence. (§§6.2 and 8.2)

In formulating either the WT identity or the quantum master equation, we must introduce renormalized parameters to parameterize the radiative corrections to the field transformation. We must tune not only the parameters of the Wilson action but also these parameters to satisfy the WT identity or the quantum master equation. The necessity of giving cutoff dependence on the symmetry transformation is reminiscent of chiral symmetry on the lattice. There, the symmetry transformation is not simply specified by the standard γ_5 matrix, but by its non-trivial extension that depends on the Dirac operator as well as lattice spacing. Ginsparg and Wilson⁴⁷⁾ found a specific form of the Ward-Takahashi (WT) identity for the presence of a lattice chiral symmetry.⁷⁹⁾ In §6.5 we derive an analogous identity in the ERG framework.⁶²⁾

As we will explain in §2.4, there is an alternative way of introducing the Wilson action S_A . Instead of S_B , we start with a bare action $S_{B,\Lambda}$ not only with an UV cutoff Λ_0 but also an IR cutoff Λ . The Wilson action can be defined as the generating functional of the connected correlation functions of $S_{B,\Lambda}$. We can then introduce the effective average action $\Gamma_{B,\Lambda}$ as the Legendre transform of S_A , or the generating functional of 1PI correlation functions. Symmetry of the continuum limit can be formulated using $\Gamma_{B,\Lambda}$ instead of S_A . For YM theories, this approach was initiated by Ellwanger.^{40),41)} The corresponding WT identity is called the **modified Slavnov-Taylor (ST) identity**, because this identity approaches the Slavnov-Taylor identity or the Jinn-Zustin equation as $\Lambda \rightarrow 0$. In the modified ST identity, the jacobian of the symmetry transformation contains the inverse of the second-order differential of $\Gamma_{B,\Lambda}$. This makes the perturbative analysis of the modified ST identity a little more complicated than that of the WT identity. For this reason we use the WT

*) We recommend 49), 51) as reviews on the BV formalism.

identity and the corresponding quantum master equation in this review; we explain the modified ST identity only briefly in §10.5 and Appendix D. For a review of the modified ST identity, we refer the reader to one of the references we will cite toward the end of this introduction.

This review consists of three main parts. Assuming that most of the readers are not familiar with the exact renormalization group or Polchinski's differential equations, we give a detailed introduction to the technique before starting its application to symmetry realization. Except in §§2, 10, we treat theories perturbatively. The review is organized as follows.

Part I consists of §§2, 3, and 4. In section 2, we define the Wilson action S_A and derive the ERG differential equation (Polchinski equation) that gives the A dependence of S_A . Primarily for notational simplicity, we take a real scalar field theory as a generic example. We then introduce the effective average action as the Legendre transform of the Wilson action. The A dependence of the effective average action is given by the Wetterich equation, widely used for non-perturbative applications of ERG. The diagrammatic interpretation of these two flow equations are given for more insights. In §3, we define perturbative renormalizability in terms of a Wilson action. A renormalizable theory is characterized by the behavior of the Wilson action for the cutoff A , large compared with the momenta carried by the field variables. In §4, we introduce “composite operators.” Following Becchi,¹⁷⁾ we give a specific meaning to the composite operators, more than an arbitrary functional of field variables. The composite operators play essential roles in our later discussions of symmetry realization: the presence of symmetry amounts to the vanishing of a certain composite operator.

Part II consists of §§5 through 9. In the ordinary treatment of a field theory, the presence of continuous symmetry is expressed as the Ward-Takahashi (WT) identity. In §5, we derive the corresponding relation satisfied by the Wilson action. This relation is expressed as the vanishing of the WT composite operator, given as the sum of the change of the Wilson action under an infinitesimal transformation and the jacobian of the transformation. We provide several examples of the formalism in §6: QED, YM theories, the Wess-Zumino model, the $O(N)$ non-linear sigma model in two dimensions, and fermionic systems with an axial symmetry. The discussions in §§5 & 6 have a serious shortcoming: the field transformations for the WT identity are applied only once, and they lack an algebraic structure. Particularly for YM theories, the BRST transformation for the Wilson action lacks nilpotency, and as a result we cannot prove the possibility of satisfying the WT identity by fine-tuning the parameters of the theory. To circumvent this difficulty, we extend the WT identity by introducing classical sources that generate the symmetry transformations. The discussion in §7 emphasizes the practical aspect of the formalism, and it is developed only to the extent necessary for the concrete application to YM theories in §8.2. In §9 we apply the formalism of §5 to the axial and chiral anomalies. The discussions are quite limited: we discuss only abelian gauge theories up to 1-loop.

Part III consists of §§10 and 11. We adapt the general formalism of Batalin and Vilkovisky to an arbitrary bare action with an UV cutoff and its Wilson action. To emphasize the generality of the framework, we adopt a matrix notation that handles

bosonic and fermionic fields equally. We show how naturally the Batalin-Vilkovisky formalism applies to the Wilson action. Contrary to §§7 & 8, we do not base our discussions on loop expansions, but we use a general functional method of §2.

Throughout this review we work on the euclidean space with D dimensions, where D may be specified in concrete models. We use the notation

$$\int_q \equiv \int \frac{d^D q}{(2\pi)^D} \quad (1.4)$$

for integrals over momenta.

Please be warned that this review is by no means a comprehensive review of the exact renormalization group approach to field theory. In the main text we cite only those references that are relevant to symmetry realization and that happen to be familiar to us. We owe an apology to those whose contributions deserve citations in a better researched review article.

Before closing the introductory section, however, we would like to give references on the aspects of ERG that may not be directly relevant to this review; this may help the reader broadening his perspective. The exact renormalization group equation of Wilson and that of Wegner & Houghton were introduced originally for non-perturbative studies, and accordingly there have been many such applications. Most non-perturbative studies use an approximation called the **local potential approximation (LPA)**, where only the local potential part of the Wilson action transforms under ERG.^{89)50)80)11)3)–5)} The analytic properties of the LPA have only recently been studied.^{21),22)} General classes of approximations to the ERG differential equation have been introduced in 48).

The ERG differential equation for the effective average action $\Gamma_{B,A}$ is now commonly called the Wetterich equation,⁸⁸⁾¹²³⁾¹²²⁾⁸¹⁾²⁴⁾ which we will derive in §2.4. The added note to 81), where the relation between the Polchinski and Wetterich equations was clarified, has a short but careful analysis of the history of the Wetterich equation. For numerical works, Wetterich's equation is known to have better convergence properties than Polchinski's.¹¹³⁾¹⁾ See Refs. 18), 19) for the state-of-the-art calculations of the critical exponents for the O(N) linear sigma model. The equivalence between the LPA for Polchinski's and that for Wetterich's has been shown in 85).

As for applications of ERG to gauge theories, there have been two other approaches besides the WT identity and its extension explained in this review:

1. Manifestly gauge invariant ERG formalism for YM theories have been introduced by Morris, Rosten, and their collaborators.^{83),84)7)8),86)87)101)99)100)}
2. The background field method has been incorporated within the ERG framework.^{96)44)43),78)33),45)}

ERG has also been applied to quantum gravity, and a possible scenario for asymptotic safety has been proposed^{74),98)}.⁷⁶⁾

Finally, we give references to the review articles that complement the present review: Morris,⁸²⁾ Litim,⁷⁷⁾ Aoki,²⁾ Bagnuls & Bervillier,⁹⁾ Berges, Tetradis, and Wetterich,²⁰⁾ Polonyi,⁹⁵⁾ and Pawłowski.⁹⁰⁾ Especially, we recommend the pedagogical review on the functional RG and gauge theories by Gies.⁴⁶⁾ For applications of

ERG to condensed matter physics, we recommend Shankar,¹⁰⁵⁾ Fisher⁴²⁾ (on RG in general), Salmhofer,¹⁰³⁾ and Delamotte.³⁹⁾




Fig. 1. Cutoff function — (a) satisfying 2, (b) satisfying 2' but not 2.

§2. Derivation of the ERG differential equations

2.1. Splitting fields into high and low momentum modes

As a generic case, we consider a real scalar theory defined by a bare action

$$S_B \equiv -\frac{1}{2} \int_p \frac{p^2 + m^2}{K(p/\Lambda_0)} \phi(p) \phi(-p) + S_{I,B}[\phi], \quad (2.1)$$

where K is a cutoff function, for which we demand the following properties:

1. $K(p/\Lambda)$ is a smooth non-increasing positive function of p^2/Λ^2 .
2. $K(p/\Lambda)$ is 1 for $p^2/\Lambda^2 < 1$.
3. $K(p/\Lambda)$ damps sufficiently fast as $p^2/\Lambda^2 \rightarrow \infty$.

(See Fig. 1(a).) Hence, $K(p/\Lambda_0)$ is small for $p^2 > \Lambda_0^2$, and the high momentum modes are suppressed by the large gaussian term. Λ_0 is the UV cutoff of the theory, but the high momentum modes of ϕ are not cut off abruptly at Λ_0 : the smoothness of $K(p/\Lambda_0)$ for $p^2/\Lambda_0^2 > 1$ gives rise to smooth damping of the modes $p^2 > \Lambda_0^2$.

The regularization in terms of a smooth cutoff function K generalizes regularization in terms of higher order derivatives which corresponds to a choice of a finite order polynomial:

$$\frac{1}{K(p/\Lambda_0)} = 1 + \sum_{n=1}^N a_n \left(\frac{p^2}{\Lambda_0^2} \right)^n. \quad (2.2)$$

With N finite, this cannot satisfy the second property

$$-p^2 \frac{d}{dp^2} K(p/\Lambda) = 0 \quad (p^2 < \Lambda^2) \quad (2.3)$$

that assures the locality of the differential equation for the exact renormalization group. We usually assume 2 for locality. (See sect. 2.3.) But it is sometimes necessary to keep K strictly less than 1 except at $p = 0$; for example in §2.4, we need that division by $1 - K(p/\Lambda)$ make sense. We can replace 2 by

- 2'. $K(0) = 1$.

(See Fig. 1(b).) For practical calculations the condition 2 is often replaced by 2'.

Let us give some concrete examples of the cutoff function. One popular choice for numerical calculations is that of Litim:⁷⁵⁾

$$K(p/\Lambda) = \begin{cases} 1 - p^2/\Lambda^2 & (p^2 < \Lambda^2), \\ 0 & (p^2 > \Lambda^2). \end{cases} \quad (2.4)$$

This satisfies 1, 2', and 3, but not 2. The gaussian function

$$K(p/\Lambda) = \exp(-p^2/\Lambda^2) \quad (2.5)$$

also satisfies only 1, 2', and 3. One example, satisfying 1, 2, and 3, is

$$K(p/\Lambda) = \begin{cases} 1 & (p^2 < \Lambda^2), \\ 1 - \exp\left[-\left(\frac{\Lambda^2}{p^2 - \Lambda^2}\right)^n\right] & (p^2 > \Lambda^2), \end{cases} \quad (2.6)$$

where n is a big enough positive number for UV convergence. (For the scalar theory in $D = 4$, we can take $n = 2$, for example.) The step function, first used in 117),

$$K(p/\Lambda) = \theta(1 - p^2/\Lambda^2) \equiv \begin{cases} 1 & (p^2 < \Lambda^2) \\ 0 & (p^2 > \Lambda^2) \end{cases} \quad (2.7)$$

is not smooth, but is often convenient for practical calculations, especially for analytic calculations.

We should remark that the choice of K is unimportant theoretically; in Appendix B on universality we will show that a different choice of K merely amounts to reparameterization and renormalization of fields.

2.1.1. Functional integrals without a source

All the momentum modes $p^2 < \Lambda_0^2$ contribute to the vacuum functional integral

$$\begin{aligned} Z_B &\equiv \int [d\phi] \exp[S_B[\phi]] \\ &= \int [d\phi] \exp\left[-\frac{1}{2} \int_p \frac{p^2 + m^2}{K(p/\Lambda_0)} \phi(p) \phi(-p) + S_{I,B}[\phi]\right], \end{aligned} \quad (2.8)$$

where the contribution of the modes $p^2 > \Lambda_0^2$ are suppressed by the cutoff function. We would like to split $\phi(p)$ into the high and low momentum modes

$$\phi(p) = \phi_h(p) + \phi_l(p), \quad (2.9)$$

so that, roughly speaking, ϕ_h contains the momentum modes $\Lambda^2 < p^2 < \Lambda_0^2$, where Λ is an arbitrarily chosen momentum scale, and ϕ_l contains the low momentum modes $p^2 < \Lambda^2$. To be more precise, in what follows we will obtain the splitting formula

$$\begin{aligned} Z_B &= \int [d\phi_l][d\phi_h] \exp\left[-\frac{1}{2} \int_p \frac{p^2 + m^2}{K(p/\Lambda)} \phi_l(p) \phi_l(-p) \right. \\ &\quad \left. - \frac{1}{2} \int_p \frac{p^2 + m^2}{K(p/\Lambda_0) - K(p/\Lambda)} \phi_h(p) \phi_h(-p) + S_{I,B}[\phi_h + \phi_l]\right]. \end{aligned} \quad (2.10)$$

The propagator of ϕ_l is the low momentum propagator

$$\frac{K(p/\Lambda)}{p^2 + m^2}, \quad (2.11)$$

while that of ϕ_h is the high momentum one

$$\frac{K(p/\Lambda_0) - K(p/\Lambda)}{p^2 + m^2}. \quad (2.12)$$

The original propagator

$$\frac{K(p/\Lambda_0)}{p^2 + m^2} \quad (2.13)$$

of $\phi = \phi_h + \phi_l$ is reproduced as their sum.

The splitting of ϕ into high and low momentum modes is an example of the following general formula:

$$\begin{aligned} & \int [d\phi] \exp \left[-\frac{1}{2} \int_p \frac{1}{A(p) + B(p)} \phi(-p) \phi(p) + S_{I,B}[\phi] \right] \\ &= \int [d\phi_1][d\phi_2] \exp \left[-\frac{1}{2} \int_p \frac{1}{A(p)} \phi_1(-p) \phi_1(p) - \frac{1}{2} \int_p \frac{1}{B(p)} \phi_2(-p) \phi_2(p) \right. \\ & \quad \left. + S_{I,B}[\phi_1 + \phi_2] \right], \end{aligned} \quad (2.14)$$

where A and B are both non-negative functions of p^2 . By choosing

$$A(p) = K(p/\Lambda_0) - K(p/\Lambda), \quad B(p) = K(p/\Lambda), \quad (2.15)$$

we obtain (2.10) from (2.14). By applying (2.14) multiple times, we obtain

$$\begin{aligned} & \int [d\phi] \exp \left[-\frac{1}{2} \int_p \frac{1}{\sum_{i=1}^n A_i(p)} \phi(-p) \phi(p) + S_{I,B}[\phi] \right] \\ &= \int [d\phi_1] \cdots [d\phi_n] \exp \left[-\frac{1}{2} \sum_{i=1}^n \int_p \frac{1}{A_i(p)} \phi_i(-p) \phi_i(p) + S_{I,B} \left[\sum_{i=1}^n \phi_i \right] \right]. \end{aligned} \quad (2.16)$$

For example, we can choose

$$A_i(p) = K(p/\Lambda_{i-1}) - K(p/\Lambda_i), \quad (2.17)$$

where

$$\Lambda_0 > \Lambda_1 > \cdots > \Lambda_{n-1} > \Lambda_n = 0. \quad (2.18)$$

Let us now prove (2.14). It is easier to go from the right-hand side to the left. Writing

$$\phi = \phi_1 + \phi_2, \quad \phi' = \phi_2 - \frac{B}{A+B} \phi, \quad (2.19)$$

we obtain

$$\begin{aligned} & -\frac{1}{2} \int_p \frac{1}{A(p)} \phi_1(-p) \phi_1(p) - \frac{1}{2} \int_p \frac{1}{B(p)} \phi_2(-p) \phi_2(p) + S_{I,B}[\phi_1 + \phi_2] \\ &= -\frac{1}{2} \int_p \frac{1}{A+B} \phi(-p) \phi(p) - \frac{1}{2} \int_p \left(\frac{1}{A} + \frac{1}{B} \right) \phi'(-p) \phi'(p) + S_{I,B}[\phi]. \end{aligned} \quad (2.20)$$

Since the jacobian for the change of variables from ϕ_1, ϕ_2 to ϕ, ϕ' is unity, we obtain

$$\begin{aligned} & \int [d\phi] \exp \left[-\frac{1}{2} \int_p \frac{1}{A} \phi_1 \phi_1 - \frac{1}{2} \int_p \frac{1}{B} \phi_2 \phi_2 + S_{I,B}[\phi_1 + \phi_2] \right] \\ &= \int [d\phi] \exp \left[-\frac{1}{2} \int_p \frac{1}{A+B} \phi(-p) \phi(p) + S_{I,B}[\phi] \right] \\ & \quad \times \int [d\phi'] \exp \left[-\frac{1}{2} \int_p \left(\frac{1}{A} + \frac{1}{B} \right) \phi'(-p) \phi'(p) \right]. \end{aligned} \quad (2.21)$$

The gaussian integral over ϕ' can be regarded as unity

$$\int [d\phi'] \exp \left[-\frac{1}{2} \int_p \left(\frac{1}{A} + \frac{1}{B} \right) \phi'(-p) \phi'(p) \right] = 1, \quad (2.22)$$

if we disregard a constant factor, the exponential of an additive constant proportional to the volume of the entire space. Hence, we obtain (2.14), and consequently (2.10).

We now define $S_{I,\Lambda}$, the interaction part of the **Wilson action**, by

$$\exp [S_{I,\Lambda}[\phi]] \equiv \int [d\phi'] \exp \left[-\frac{1}{2} \int_p \frac{p^2 + m^2}{K_0(p) - K(p)} \phi'(p) \phi'(-p) + S_{I,B}[\phi + \phi'] \right], \quad (2.23)$$

where we have used a short hand notation

$$K_0(p) \equiv K(p/\Lambda_0), \quad K(p) \equiv K(p/\Lambda). \quad (2.24)$$

Defining the full Wilson action by

$$S_\Lambda[\phi] \equiv -\frac{1}{2} \int_p \frac{p^2 + m^2}{K(p/\Lambda)} \phi(-p) \phi(p) + S_{I,\Lambda}[\phi], \quad (2.25)$$

we obtain Z_B as the functional integral of the full Wilson action

$$Z_B = \int [d\phi] \exp [S_\Lambda[\phi]]. \quad (2.26)$$

We have thus shown that as far as the vacuum functional integral Z_B is concerned, the bare action S_B with cutoff Λ_0 can be replaced by a Wilson action S_Λ with an arbitrary cutoff $\Lambda < \Lambda_0$. In the next subsection, we wish to show that this equivalence goes further: S_Λ and S_B give the same correlation functions of the fields.

The transformation from S_B to S_Λ is called an **exact renormalization group (ERG)** transformation. The word “**exact**” is a mnemonic for no loss of information.

2.2. Functional integrals with a source

We consider the generating functional $W_B[J]$ of connected correlation functions:

$$\exp[W_B[J]] \equiv \int [d\phi] \exp \left[S_B[\phi] + \int_p J(-p)\phi(p) \right]. \quad (2.27)$$

Similarly, we consider

$$\exp[W_A[J]] \equiv \int [d\phi] \exp \left[S_A[\phi] + \int_p J(-p)\phi(p) \right] \quad (2.28)$$

for the Wilson action. In the following we wish to show that W_B and W_A are related as

$$W_B[J] = W_A \left[\frac{K_0}{K} J \right] - \frac{1}{2} \int_p \frac{K_0 - K}{p^2 + m^2} \frac{K_0}{K} J(-p)J(p), \quad (2.29)$$

where we have used the short hand notation (2.24), omitting the momentum variable p . This relation implies that the correlation functions of the bare theory can be calculated using the Wilson action.

To derive (2.29), we first define

$$S_B[\phi; J] \equiv -\frac{1}{2} \int_p \frac{p^2 + m^2}{K_0} \phi(p)\phi(-p) + S_{I,B}[\phi; J], \quad (2.30)$$

where

$$S_{I,B}[\phi; J] \equiv S_{I,B}[\phi] + \int_p J(-p)\phi(p) \quad (2.31)$$

is a bare action that has a linear coupling to the classical external field J . By definition we obtain

$$\exp[W_B[J]] = \int [d\phi] \exp[S_B[\phi; J]]. \quad (2.32)$$

Let us now call the corresponding Wilson action by $S_A[\phi; J]$ and its interaction part by $S_{I,A}[\phi; J]$. Then, from the result of the previous section, we obtain

$$\exp[W_B[J]] = \int [d\phi] \exp[S_A[\phi; J]]. \quad (2.33)$$

We obtain (2.29) by working out $S_A[\phi; J]$, as we will show below.

From the definition (2.23) of a Wilson action, we obtain

$$\begin{aligned} \exp[S_{I,A}[\phi; J]] &\equiv \int [d\phi'] \exp \left[-\frac{1}{2} \int_p \frac{p^2 + m^2}{K_0 - K} \phi' \phi' + S_{I,B}[\phi + \phi'; J] \right] \\ &= \int [d\phi'] \exp \left[-\frac{1}{2} \int_p \frac{p^2 + m^2}{K_0 - K} \phi' \phi' + S_{I,B}[\phi + \phi'] + \int J(\phi + \phi') \right]. \end{aligned} \quad (2.34)$$

We can rewrite this as

$$\begin{aligned} \exp[S_{I,A}[\phi; J]] &= \int [d\phi'] \exp \left[-\frac{1}{2} \int_p \frac{p^2 + m^2}{K_0 - K} \left(\phi' - \frac{K_0 - K}{p^2 + m^2} J \right) \left(\phi' - \frac{K_0 - K}{p^2 + m^2} J \right) \right. \\ &\quad \left. + S_{I,B}[\phi + \phi'] + \int J\phi + \frac{1}{2} \int \frac{K_0 - K}{p^2 + m^2} J J \right]. \end{aligned} \quad (2.35)$$

Now, shifting ϕ' , we obtain

$$\begin{aligned} \exp [S_{I,\Lambda}[\phi; J]] &= \int [d\phi'] \exp \left[-\frac{1}{2} \int_p \frac{p^2 + m^2}{K_0 - K} \phi' \phi' \right. \\ &\quad \left. + S_{I,B} \left[\phi + \frac{K_0 - K}{p^2 + m^2} J + \phi' \right] + \int J \phi + \frac{1}{2} \int \frac{K_0 - K}{p^2 + m^2} J J \right] \\ &= \exp \left[S_{I,\Lambda} \left[\phi + \frac{K_0 - K}{p^2 + m^2} J \right] + \int J \phi + \frac{1}{2} \int \frac{K_0 - K}{p^2 + m^2} J J \right]. \end{aligned} \quad (2.36)$$

This implies an important intermediate result

$$S_{I,\Lambda}[\phi; J] = S_{I,\Lambda} \left[\phi + \frac{K_0 - K}{p^2 + m^2} J \right] + \int_p J(-p) \phi(p) + \frac{1}{2} \int_p \frac{K_0 - K}{p^2 + m^2} J(-p) J(p), \quad (2.37)$$

that the dependence on J is given mostly by the shift of ϕ proportional to J . If J couples non-linearly to the field in the bare action, however, there is no simple formula like this for the J dependence of the Wilson action.

Now, using (2.37), we can compute $W_B[J]$ as follows:

$$\begin{aligned} \exp [W_B[J]] &= \int [d\phi] \exp [S_\Lambda[\phi; J]] = \int [d\phi] \exp \left[-\frac{1}{2} \int_p \frac{p^2 + m^2}{K} \phi \phi \right. \\ &\quad \left. + S_{I,\Lambda} \left[\phi + \frac{K_0 - K}{p^2 + m^2} J \right] + \int J \phi + \frac{1}{2} \int \frac{K_0 - K}{p^2 + m^2} J J \right]. \end{aligned} \quad (2.38)$$

Changing the variable from $\phi(p)$ to

$$\phi'(p) \equiv \phi(p) + \frac{K_0 - K}{p^2 + m^2} J(p), \quad (2.39)$$

we obtain the desired relation (2.29):

$$\begin{aligned} \exp [W_B[J]] &= \int [d\phi'] \exp \left[S_\Lambda[\phi'] + \int \frac{K_0}{K} J \phi' - \frac{1}{2} \int \frac{K_0 - K}{p^2 + m^2} \frac{K_0}{K} J J \right] \\ &= \exp \left[W_\Lambda \left[\frac{K_0}{K} J \right] - \frac{1}{2} \int \frac{K_0 - K}{p^2 + m^2} \frac{K_0}{K} J J \right]. \end{aligned} \quad (2.40)$$

In fact we can go backward: given (2.29), we can derive the definition (2.23) of the Wilson action. Thus, (2.23) and (2.29) are equivalent. We postpone the derivation of (2.23) from (2.29) till §4.

Let us now derive the consequences of the relation (2.29). W_B and W_Λ are the generating functionals of connected correlation functions, which we denote by brackets:

$$\langle \phi(p_1) \cdots \phi(p_n) \rangle_{S_B} (2\pi)^D \delta^{(D)}(p_1 + \cdots + p_n) \equiv \prod_{i=1}^n \frac{\delta}{\delta J(-p_i)} \cdot W_B[J] \Big|_{J=0}, \quad (2.41)$$

$$\langle \phi(p_1) \cdots \phi(p_n) \rangle_{S_\Lambda} (2\pi)^D \delta^{(D)}(p_1 + \cdots + p_n) \equiv \prod_{i=1}^n \frac{\delta}{\delta J(-p_i)} \cdot W_\Lambda[J] \Big|_{J=0}. \quad (2.42)$$

Then, (2.29) implies

$$\begin{cases} \langle \phi(p)\phi(-p) \rangle_{S_B} = \frac{K_0^2}{K^2} \langle \phi(p)\phi(-p) \rangle_{S_A} - \frac{K_0-K}{p^2+m^2} \frac{K_0}{K}, \\ \langle \phi(p_1) \cdots \phi(p_n) \rangle_{S_B} = \prod_{i=1}^n \frac{K_0(p_i)}{K(p_i)} \cdot \langle \phi(p_1) \cdots \phi(p_n) \rangle_{S_A}, \end{cases} \quad (2.43)$$

where $n > 2$. Hence, the correlation functions of the bare action are completely determined by those of the Wilson action. We do not miss any physics by reducing the momentum cutoff from Λ_0 to Λ , as long as $K(p/\Lambda)$ is a smooth non-vanishing function so that the inverse $1/K(p/\Lambda)$ makes sense for all momenta.

2.3. Polchinski equation

The Wilson action S_A is obtained from the bare action S_B by integrating over the momenta between Λ and Λ_0 . We wish to consider the change of the Wilson action under an infinitesimal change of Λ , and express the result as a differential equation.

We first recall the precise definition of the Wilson action (2.23):

$$\exp[S_{I,\Lambda}[\phi]] \equiv \int [d\phi'] \exp \left[-\frac{1}{2} \int_p \frac{p^2 + m^2}{K_0 - K} \phi'(p) \phi'(-p) + S_{I,B}[\phi + \phi'] \right]. \quad (2.44)$$

In deriving the Polchinski equation, it is convenient to write ϕ' for $\phi' + \phi$ to rewrite

$$\exp[S_{I,\Lambda}[\phi]] \equiv \int [d\phi'] \exp \left[-\frac{1}{2} \int_p \frac{p^2 + m^2}{K_0 - K} (\phi' - \phi)(p) (\phi' - \phi)(-p) + S_{I,B}[\phi'] \right]. \quad (2.45)$$

Differentiating the above with respect to Λ , we obtain

$$\begin{aligned} -\Lambda \frac{\partial}{\partial \Lambda} \exp[S_{I,\Lambda}[\phi]] &= \frac{1}{2} \int [d\phi'] \int_p \frac{p^2 + m^2}{K_0 - K} \frac{\Delta(p/\Lambda)}{K_0 - K} (\phi' - \phi)(p) (\phi' - \phi)(-p) \\ &\times \exp \left[-\frac{1}{2} \int_p \frac{p^2 + m^2}{K_0 - K} (\phi' - \phi)(p) (\phi' - \phi)(-p) + S_{I,B}[\phi'] \right], \end{aligned} \quad (2.46)$$

where we have defined

$$\Delta(p/\Lambda) \equiv -2p^2 \frac{\partial}{\partial p^2} K(p/\Lambda) = \Lambda \frac{\partial}{\partial \Lambda} K(p/\Lambda). \quad (2.47)$$

The sketch of $\Delta(p/\Lambda)$ is given in Fig. 2, showing a peak just above $p^2 = \Lambda^2$, and vanishing for $p^2 < \Lambda^2$ if the property 2 is assumed for K .^{*)} We now compare the Λ -derivative with the second order functional derivative with respect to ϕ . The first order derivative is

$$\begin{aligned} \frac{\delta}{\delta \phi(p)} \exp[S_{I,\Lambda}[\phi]] &= \frac{p^2 + m^2}{K_0 - K} \int [d\phi'] (\phi' - \phi)(-p) \\ &\times \exp \left[-\frac{1}{2} \int_q \frac{q^2 + m^2}{K_0 - K} (\phi' - \phi)(q) (\phi' - \phi)(-q) + S_{I,B}[\phi'] \right]. \end{aligned} \quad (2.48)$$

^{*)} If only 2' is assumed for K , the function $\Delta(p/\Lambda)$ vanishes at $p = 0$, but is non-vanishing for $p^2 < \Lambda^2$.




Fig. 2. $\Delta(p/\Lambda) \equiv -2p^2 \frac{\partial K(p/\Lambda)}{\partial p^2}$ vanishes for $p^2 < \Lambda^2$ if $K(p/\Lambda) = 1$ for $p^2 < \Lambda^2$.

In differentiating this further with respect to $\phi(-p)$, we ignore differentiating $\phi(-p)$ in front of the exponential, since it would generate merely an additive constant to $\partial S_{I,\Lambda}/\partial \Lambda$, independent of ϕ .*) We then obtain

$$\begin{aligned} \frac{\delta^2}{\delta\phi(-p)\delta\phi(p)} \exp[S_{I,\Lambda}[\phi]] &= \left(\frac{p^2 + m^2}{K_0 - K} \right)^2 \int [d\phi'] (\phi' - \phi)(-p) (\phi' - \phi)(p) \\ &\times \exp \left[-\frac{1}{2} \int_q \frac{q^2 + m^2}{K_0 - K} (\phi' - \phi)(q) (\phi' - \phi)(-q) + S_{I,B}[\phi'] \right]. \end{aligned} \quad (2.49)$$

Hence, comparing this with (2.46), we obtain the differential equation

$$-\Lambda \frac{\partial}{\partial \Lambda} \exp[S_{I,\Lambda}[\phi]] = \frac{1}{2} \int_p \frac{\Delta(p/\Lambda)}{p^2 + m^2} \frac{\delta^2}{\delta\phi(p)\delta\phi(-p)} \exp[S_{I,\Lambda}[\phi]]. \quad (2.50)$$

Rewriting this for $S_{I,\Lambda}$, we get

$$-\Lambda \frac{\partial}{\partial \Lambda} S_{I,\Lambda}[\phi] = \frac{1}{2} \int_p \frac{\Delta(p/\Lambda)}{p^2 + m^2} \left\{ \frac{\delta S_{I,\Lambda}[\phi]}{\delta\phi(-p)} \frac{\delta S_{I,\Lambda}[\phi]}{\delta\phi(p)} + \frac{\delta^2 S_{I,\Lambda}[\phi]}{\delta\phi(-p)\delta\phi(p)} \right\}. \quad (2.51)$$

This is called the **Polchinski differential equation**.⁹⁴⁾

If we assume property 2, $K(p/\Lambda) = 1$ for $p^2 < \Lambda^2$, then

$$\Delta(p/\Lambda) = 0 \quad \text{for } p^2 < \Lambda^2. \quad (2.52)$$

This implies that the deformation of the Wilson action is restricted to the momentum region $p^2 > \Lambda^2$. It is in this sense that we call the ERG transformation **local**. If only 2' is satisfied, we may still call the ERG transformation “almost local.”

In the above we have derived the Polchinski differential equation starting from the definition (2.23) of the Wilson action. Note that (2.23) in fact gives an integral formula for the solution of the differential equation under the initial condition

$$S_{I,\Lambda} \Big|_{\Lambda=\Lambda_0} = S_{I,B}. \quad (2.53)$$

Finally let us rewrite the Polchinski equation for the full action

$$S_\Lambda[\phi] = -\frac{1}{2} \int_p \frac{p^2 + m^2}{K(p/\Lambda)} \phi(-p)\phi(p) + S_{I,\Lambda}[\phi]. \quad (2.54)$$

*) The constant is proportional to the volume of the entire space $(2\pi)^D \delta^{(D)}(0)$.




Fig. 3. Sketch of $\frac{1}{K_0 - K}$. (a) If $K(p/\Lambda) = 1$ for $p^2 < \Lambda^2$, this is infinite for $p^2 < \Lambda^2$. (b) If $K(p/\Lambda) < 1$ for $p^2 < \Lambda^2$, it diverges only at $p = 0$.

It is straightforward to obtain

$$\begin{aligned}
 & -\Lambda \frac{\partial}{\partial \Lambda} S_\Lambda[\phi] \\
 &= \int_p \frac{\Delta(p/\Lambda)}{p^2 + m^2} \left[\frac{p^2 + m^2}{K(p/\Lambda)} \phi(p) \frac{\delta S_\Lambda[\phi]}{\delta \phi(p)} + \frac{1}{2} \left\{ \frac{\delta S_\Lambda[\phi]}{\delta \phi(-p)} \frac{\delta S_\Lambda[\phi]}{\delta \phi(p)} + \frac{\delta^2 S_\Lambda[\phi]}{\delta \phi(-p) \delta \phi(p)} \right\} \right].
 \end{aligned} \tag{2.55}$$

2.4. Wetterich equation

The purpose of this subsection is to derive what is commonly called the Wetterich equation,⁽⁸⁸⁾⁽¹²³⁾⁽¹²²⁾⁽⁸¹⁾⁽²⁴⁾ a counterpart of the Polchinski equation (2.51).*) Even though almost all our discussions on symmetry are done with the Wilson action that satisfies the Polchinski equation, we would like to dedicate a whole subsection on the Wetterich equation, because most non-perturbative numerical works deal with the Wetterich equation instead of the Polchinski equation.

We introduce an action

$$S_{B,\Lambda}[\phi] \equiv -\frac{1}{2} \int_p \frac{p^2 + m^2}{K_0 - K} \phi(-p) \phi(p) + S_{I,B}[\phi], \tag{2.56}$$

which is obtained from the bare action S_B by the following replacement of the gaussian term:

$$\frac{p^2 + m^2}{K(p/\Lambda_0)} \longrightarrow \frac{p^2 + m^2}{K(p/\Lambda_0) - K(p/\Lambda)}. \tag{2.57}$$

With the weight $\exp[S_B]$, $\phi(p)$ of all momenta $p^2 < \Lambda_0^2$ contribute to the functional integral. With the above replacement, however, the main contribution comes from those high momentum modes $\phi(p)$ with $\Lambda^2 < p^2 < \Lambda_0^2$. (See Fig. 3.) Hence, $S_{B,\Lambda}$ has Λ as the IR cutoff and Λ_0 as the UV cutoff. Taking the difference between $S_{B,\Lambda}$ and S_B , we can also write

$$S_{B,\Lambda}[\phi] = S_B[\phi] - \frac{1}{2} \int_p R_\Lambda(p) \phi(p) \phi(-p), \tag{2.58}$$

*) See the added note of 81) for a historical perspective on this equation.

where we define

$$R_\Lambda(p) \equiv (p^2 + m^2) \left(\frac{1}{K_0 - K} - \frac{1}{K_0} \right) = (p^2 + m^2) \frac{K}{K_0(K_0 - K)}. \quad (2.59)$$

In the limit $\Lambda \rightarrow 0+$, K and hence R_Λ vanish, and we obtain the bare action:

$$\lim_{\Lambda \rightarrow 0+} S_{B,\Lambda} = S_B. \quad (2.60)$$

Throughout §2.4, we will assume

$$K(p/\Lambda) < 1 \quad \text{for} \quad 0 < p^2 < \Lambda^2. \quad (2.61)$$

This is necessary so that

$$K(p/\Lambda_0) - K(p/\Lambda) \neq 0 \quad (2.62)$$

except at $p = 0$ and $p \rightarrow \infty$. Then the division by $K(p/\Lambda_0) - K(p/\Lambda)$ makes sense. We will do the following in the remainder of §2.4: we

1. introduce the generating functional $W_{B,\Lambda}$ of $S_{B,\Lambda}$ by (2.63), and define its Legendre transform $\Gamma_{B,\Lambda}$ by (2.65) and (2.66),
2. show $W_{B,\Lambda}$ is essentially the Wilson action S_Λ as given by (2.79), and derive the relation between S_Λ and $\Gamma_{B,\Lambda}$ as given by (2.84) and (2.85),
3. derive the Wetterich equation (2.108), giving the Λ dependence of the interaction part of $\Gamma_{B,\Lambda}$.

2.4.1. Introducing $W_{B,\Lambda}$ and $\Gamma_{B,\Lambda}$

Let us first define the generating functional $W_{B,\Lambda}$ of $S_{B,\Lambda}$ by

$$\exp[W_{B,\Lambda}[J]] \equiv \int [d\phi] \exp \left[S_{B,\Lambda}[\phi] + \int_p J(-p)\phi(p) \right]. \quad (2.63)$$

Note

$$\lim_{\Lambda \rightarrow 0} W_{B,\Lambda} = W_B. \quad (2.64)$$

For $\Lambda > 0$, $W_{B,\Lambda}$ differs from both W_B and W_Λ . Recall that W_Λ , the generating functional of the Wilson action S_Λ , is essentially the same as W_B , as has been shown in subsect. 2.2.

We then define its Legendre transform, called the **effective average action**, by

$$\Gamma_{B,\Lambda}[\Phi] \equiv W_{B,\Lambda}[J] - \int_p J(-p)\Phi(p), \quad (2.65)$$

where J is determined by the condition

$$\frac{\delta W_{B,\Lambda}[J]}{\delta J(-p)} = \Phi(p) \quad (2.66)$$

as a functional of Φ . From the general properties of Legendre transformations, we can immediately conclude the following:

(a) We can obtain $W_{B,\Lambda}$ as the inverse Legendre transform of $\Gamma_{B,\Lambda}$:

$$W_{B,\Lambda}[J] = \Gamma_{B,\Lambda}[\Phi] + \int_p J(-p)\Phi(p), \quad (2.67)$$

where Φ is determined by

$$\frac{\delta \Gamma_{B,\Lambda}[\Phi]}{\delta \Phi(p)} = -J(-p) \quad (2.68)$$

as a functional of J .

(b) (2.66) and (2.68) give

$$\frac{\delta \Phi(p)}{\delta J(q)} = \frac{\delta^2 W_{B,\Lambda}[J]}{\delta J(-p)\delta J(q)}, \quad (2.69)$$

and

$$\frac{\delta J(q)}{\delta \Phi(p)} = -\frac{\delta^2 \Gamma_{B,\Lambda}[\Phi]}{\delta \Phi(-q)\delta \Phi(p)}. \quad (2.70)$$

(c) The above are the inverse of each other, and we obtain

$$\int_r \frac{\delta^2 W_{B,\Lambda}[J]}{\delta J(-p)\delta J(r)} \frac{\delta^2 \Gamma_{B,\Lambda}[\Phi]}{\delta \Phi(-r)\delta \Phi(q)} = -(2\pi)^D \delta^{(D)}(p-q), \quad (2.71)$$

and

$$\int_r \frac{\delta^2 \Gamma_{B,\Lambda}[\Phi]}{\delta \Phi(-p)\delta \Phi(r)} \frac{\delta^2 W_{B,\Lambda}[J]}{\delta J(-r)\delta J(q)} = -(2\pi)^D \delta^{(D)}(p-q). \quad (2.72)$$

(d) Given an infinitesimal change $\Delta W_{B,\Lambda}[J]$, the corresponding infinitesimal change of $\Gamma_{B,\Lambda}[\Phi]$ satisfies

$$\Delta \Gamma_{B,\Lambda}[\Phi] = \Delta W_{B,\Lambda}[J] \quad (2.73)$$

where Φ is related to J by (2.66). (Quick proof) By definition (suppressing the subscript B,Λ), we obtain

$$(\Gamma + \Delta \Gamma)[\Phi + \Delta \Phi] = (W + \Delta W)[J] - \int J \cdot (\Phi + \Delta \Phi), \quad (2.74)$$

where

$$\frac{\delta(W + \Delta W)[J]}{\delta J(-p)} = \Phi(p) + \Delta \Phi(p). \quad (2.75)$$

Hence,

$$\Delta \Gamma[\Phi] + \int \frac{\delta \Gamma}{\delta \Phi} \Delta \Phi = \Delta W[J] - \int J \cdot \Delta \Phi. \quad (2.76)$$

From (2.68), the terms proportional to $\Delta \Phi$ cancel, and we obtain (2.73).

2.4.2. Relating $W_{B,\Lambda}$ to S_Λ , and deriving the Legendre transformation between $\Gamma_{B,\Lambda}$ and S_Λ

The relation between the Wilson action S_Λ and the effective average action $\Gamma_{B,\Lambda}$ was first derived by T. Morris.⁸¹⁾ We reproduce his results in the following.

Our first task is to derive the relation of the generating functional $W_{B,\Lambda}$ and the Wilson action S_Λ . Using (2.56) and (2.63), we obtain

$$\exp[W_{B,\Lambda}[J]] = \int [d\phi] \exp \left[-\frac{1}{2} \int_p \frac{p^2 + m^2}{K_0 - K} \phi(-p) \phi(p) + S_{I,B}[\phi] + \int_p J(-p) \phi(p) \right]. \quad (2.77)$$

From the definition (2.34) of $S_{I,\Lambda}[\phi; J]$, we find

$$\begin{aligned} W_{B,\Lambda}[J] &= S_{I,\Lambda}[0; J] \\ &= S_{I,\Lambda} \left[\frac{K_0 - K}{p^2 + m^2} J \right] + \frac{1}{2} \int_p \frac{K_0 - K}{p^2 + m^2} J(-p) J(p). \end{aligned} \quad (2.78)$$

where we have used the intermediate result (2.37). Using the definition (2.25) of the full Wilson action S_Λ , this can be rewritten as

$$W_{B,\Lambda}[J] = S_\Lambda \left[\frac{K_0 - K}{p^2 + m^2} J \right] + \frac{1}{2} \int_p \frac{K_0 - K}{p^2 + m^2} \frac{K_0}{K} J(-p) J(p). \quad (2.79)$$

Thus, we find that the Wilson action S_Λ is basically the generating functional for the action $S_{B,\Lambda}$. Especially in the limit $\Lambda \rightarrow 0+$, where K vanishes, and $W_{B,\Lambda}$ becomes W_B , we obtain

$$\lim_{\Lambda \rightarrow 0+} S_{I,\Lambda} \left[\frac{K_0}{p^2 + m^2} J \right] = W_B[J] - \frac{1}{2} \int_p \frac{K_0}{p^2 + m^2} J(-p) J(p). \quad (2.80)$$

Namely, in the $\Lambda \rightarrow 0+$ limit, the interaction part of the Wilson action becomes that of the generating functional W_B .

We now use the above result to express $\Gamma_{B,\Lambda}$ in terms of S_Λ . Substituting (2.79) into (2.65), we obtain

$$\Gamma_{B,\Lambda}[\Phi] = \frac{1}{2} \int_p \frac{K_0 - K}{p^2 + m^2} \frac{K_0}{K} J J + S_\Lambda \left[\frac{K_0 - K}{p^2 + m^2} J \right] - \int_p J \Phi, \quad (2.81)$$

where

$$\Phi(p) = \frac{K_0 - K}{p^2 + m^2} \left[\frac{K_0}{K} J(p) + \frac{\delta S_\Lambda[\phi]}{\delta \phi(-p)} \Big|_{\phi = \frac{K_0 - K}{p^2 + m^2} J} \right]. \quad (2.82)$$

It is more convenient to use

$$\phi(p) \equiv \frac{K(p/\Lambda_0) - K(p/\Lambda)}{p^2 + m^2} J(p) \quad (2.83)$$

as field variables instead of J . We then obtain⁸¹⁾

$$\Gamma_{B,\Lambda}[\Phi] = S_\Lambda[\phi] + \int_p \frac{p^2 + m^2}{K_0 - K} \left(\frac{1}{2} \frac{K_0}{K} \phi(p) \phi(-p) - \Phi(p) \phi(-p) \right), \quad (2.84)$$

where

$$\Phi(p) = \frac{K_0}{K} \phi(p) + \frac{K_0 - K}{p^2 + m^2} \frac{\delta S_\Lambda[\phi]}{\delta \phi(-p)}. \quad (2.85)$$

Conversely, by regarding $S_\Lambda[\phi]$ as the Legendre transform of $\Gamma_{B,\Lambda}[\Phi]$, we obtain

$$\phi(p) = -\frac{K_0 - K}{p^2 + m^2} \frac{\delta \Gamma_{B,\Lambda}[\Phi]}{\delta \Phi(-p)}. \quad (2.86)$$

Let us now rewrite the above three equations using $S_{I,\Lambda}[\phi]$ and the interaction part of $\Gamma_{B,\Lambda}$, defined by

$$\Gamma_{I,B,\Lambda}[\Phi] \equiv \Gamma_{B,\Lambda}[\Phi] + \frac{1}{2} \int_p \frac{p^2 + m^2}{K_0 - K} \Phi(-p) \Phi(p). \quad (2.87)$$

First, we can rewrite (2.84) as

$$\Gamma_{I,B,\Lambda}[\Phi] = S_{I,\Lambda}[\phi] + \frac{1}{2} \int_p \frac{p^2 + m^2}{K_0 - K} (\Phi - \phi)(-p) (\Phi - \phi)(p). \quad (2.88)$$

Then, (2.85) gives

$$\Phi(p) = \phi(p) + \frac{K_0 - K}{p^2 + m^2} \frac{\delta S_{I,\Lambda}[\phi]}{\delta \phi(-p)}, \quad (2.89)$$

and (2.86) gives

$$\phi(p) = \Phi(p) - \frac{K_0 - K}{p^2 + m^2} \frac{\delta \Gamma_{I,B,\Lambda}[\Phi]}{\delta \Phi(-p)}. \quad (2.90)$$

Thus, we obtain

$$\frac{\delta S_{I,\Lambda}[\phi]}{\delta \phi(-p)} = \frac{\delta \Gamma_{I,B,\Lambda}[\Phi]}{\delta \Phi(-p)}. \quad (2.91)$$

Differentiating (2.89) with respect to $\phi(q)$, we obtain

$$\frac{\delta \Phi(p)}{\delta \phi(q)} = (2\pi)^D \delta^{(D)}(p - q) + \frac{K_0 - K}{p^2 + m^2} \frac{\delta^2 S_{I,\Lambda}[\phi]}{\delta \phi(-p) \delta \phi(q)}, \quad (2.92)$$

and differentiating (2.90) with respect to $\Phi(q)$, we obtain

$$\begin{aligned} \frac{\delta \phi(p)}{\delta \Phi(q)} &= (2\pi)^D \delta^{(D)}(p - q) - \frac{K_0 - K}{p^2 + m^2} \frac{\delta^2 \Gamma_{I,B,\Lambda}[\Phi]}{\delta \Phi(-p) \delta \Phi(q)} \\ &= -\frac{K_0 - K}{p^2 + m^2} \Gamma_{B,\Lambda}^{(2)}(p, -q), \end{aligned} \quad (2.93)$$

where we define

$$\Gamma_{B,\Lambda}^{(2)}(p, -q) \equiv \frac{\delta^2 \Gamma_{B,\Lambda}[\Phi]}{\delta \Phi(-p) \delta \Phi(q)}. \quad (2.94)$$

Since

$$\int_q \frac{\delta \Phi(p)}{\delta \phi(q)} \frac{\delta \phi(q)}{\delta \Phi(r)} = (2\pi)^D \delta^{(D)}(p - r), \quad (2.95)$$

we obtain

$$\int_q \frac{\delta\Phi(p)}{\delta\phi(q)} (-) \frac{K_0 - K}{q^2 + m^2} \Gamma_{B,\Lambda}^{(2)}(q, -r) = (2\pi)^D \delta^{(D)}(p - r). \quad (2.96)$$

Hence, defining

$$\begin{aligned} \left(\Gamma_{B,\Lambda}^{(2)} \right)^{-1} (p, -q) &\equiv - \frac{\delta\Phi(p)}{\delta\phi(q)} \frac{K_0 - K}{q^2 + m^2} \\ &= - (2\pi)^D \delta^{(D)}(p - q) \frac{K_0 - K}{p^2 + m^2} \\ &\quad - \frac{K_0 - K}{p^2 + m^2} \frac{\delta^2 S_{I,\Lambda}[\phi]}{\delta\phi(-p) \delta\phi(q)} \frac{K_0 - K}{q^2 + m^2}, \end{aligned} \quad (2.97)$$

where we have used (2.92), we obtain

$$\int_q \left(\Gamma_{B,\Lambda}^{(2)} \right)^{-1} (p, -q) \Gamma_{B,\Lambda}^{(2)}(q, -r) = (2\pi)^D \delta^{(D)}(p - r). \quad (2.98)$$

This justifies our notation (2.97).

Before deriving the Wetterich equation, let us consider the two limits of $\Gamma_{I,B,\Lambda}$ as $\Lambda \rightarrow \Lambda_0 - 0$ or $\Lambda \rightarrow 0+$.

- (i) $\Lambda \rightarrow \Lambda_0 - 0$ limit — From (2.89), we obtain $\Phi = \phi$, and combining this with (2.88) gives

$$\lim_{\Lambda \rightarrow \Lambda_0} \Gamma_{I,B,\Lambda} = \lim_{\Lambda \rightarrow \Lambda_0} S_{I,\Lambda} = S_{I,B}. \quad (2.99)$$

- (ii) $\Lambda \rightarrow 0+$ limit — (2.60) implies

$$\lim_{\Lambda \rightarrow 0+} \Gamma_{B,\Lambda} = \Gamma_B, \quad (2.100)$$

where Γ_B is the effective action for the bare action.^{*)} Defining the interaction part of Γ_B such that

$$\Gamma_B[\Phi] \equiv -\frac{1}{2} \int_p \frac{p^2 + m^2}{K_0} \Phi(-p) \Phi(p) + \Gamma_{I,B}[\Phi], \quad (2.101)$$

we obtain

$$\lim_{\Lambda \rightarrow 0+} \Gamma_{I,B,\Lambda} = \Gamma_{I,B}. \quad (2.102)$$

Thus, $\Gamma_{I,B,\Lambda}$ ($\Lambda_0 > \Lambda > 0$) interpolates between the bare action $S_{I,B}$ and its effective action $\Gamma_{I,B}$. In contrast, the Wilson action $S_{I,\Lambda}$ interpolates between the bare action $S_{I,B}$ and the interaction part of the generating functional W_B , up to rescaling of the source as given in (2.80).

^{*)} Strictly speaking, (2.100) is valid only if $S_{I,\Lambda}$ has no tadpole, i.e., no interaction term proportional to $\phi(0)$ at zero momentum.

2.4.3. Deriving the Wetterich equation

We are now ready to derive the Wetterich equation. Using the property (d) of the Legendre transform, the Λ derivative of $\Gamma_{I,B,\Lambda}$ is obtained as

$$-\Lambda \frac{\partial}{\partial \Lambda} \Gamma_{I,B,\Lambda}[\Phi] = -\Lambda \frac{\partial}{\partial \Lambda} S_{I,\Lambda}[\phi] - \frac{1}{2} \int_p \frac{p^2 + m^2}{(K_0 - K)^2} \Delta(p/\Lambda) (\Phi - \phi)(-p) (\Phi - \phi)(p). \quad (2.103)$$

Using (2.51) and (2.89), we obtain

$$-\Lambda \frac{\partial}{\partial \Lambda} \Gamma_{I,B,\Lambda}[\Phi] = \frac{1}{2} \int_p \frac{\Delta(p/\Lambda)}{p^2 + m^2} \frac{\delta^2 S_{I,\Lambda}[\phi]}{\delta \phi(-p) \delta \phi(p)}. \quad (2.104)$$

We rewrite this as

$$-\Lambda \frac{\partial}{\partial \Lambda} \Gamma_{I,B,\Lambda}[\Phi] = \frac{1}{2} \int_p (p^2 + m^2) \frac{\Delta(p/\Lambda)}{(K_0 - K)^2} \cdot \frac{K_0 - K}{p^2 + m^2} \frac{\delta^2 S_{I,\Lambda}[\phi]}{\delta \phi(-p) \delta \phi(p)} \frac{K_0 - K}{p^2 + m^2}. \quad (2.105)$$

We now note

$$-\Lambda \frac{\partial}{\partial \Lambda} R_\Lambda(p) = -(p^2 + m^2) \frac{\Delta(p/\Lambda)}{(K_0 - K)^2}, \quad (2.106)$$

and (2.97)

$$\begin{aligned} & \frac{K_0 - K}{p^2 + m^2} \frac{\delta^2 S_{I,\Lambda}[\phi]}{\delta \phi(-p) \delta \phi(p)} \frac{K_0 - K}{p^2 + m^2} \\ &= - \left(\Gamma_{B,\Lambda}^{(2)} \right)^{-1} (p, -p) - (2\pi)^D \delta^{(D)}(0) \cdot \frac{K_0 - K}{p^2 + m^2}. \end{aligned} \quad (2.107)$$

Thus, ignoring a field independent constant (proportional to the space volume), we obtain the Wetterich equation⁸⁸⁾¹²³⁾¹²²⁾⁸¹⁾²⁴⁾

$$-\Lambda \frac{\partial}{\partial \Lambda} \Gamma_{I,B,\Lambda}[\Phi] = \frac{1}{2} \int_p \left(-\Lambda \frac{\partial}{\partial \Lambda} R_\Lambda(p) \right) \left(\Gamma_{B,\Lambda}^{(2)} \right)^{-1} (p, -p). \quad (2.108)$$

There is no formal integral formula for the solution of the Wetterich equation such as (2.23) for the solution of the Polchinski equation.

2.5. Recapitulation

For the reader's convenience, we would like to tabulate the functionals and their relationship, introduced thus far:

1. the bare and Wilson actions (Table I) — Given a bare action S_B with cutoff Λ_0 , we construct its Wilson action S_Λ by integrating the field over momenta between Λ and Λ_0 . The generating functionals of the connected correlation functions for S_B and S_Λ are basically the same.
2. the bare action with both UV and IR cutoffs (Table II) — Instead of the bare action S_B with an UV cutoff Λ_0 , we introduce $S_{B,\Lambda}$ with both UV and IR cutoffs Λ_0 & Λ . We can define its generating functional $W_{B,\Lambda}$ of the connected correlation functions, and the Legendre transform $\Gamma_{B,\Lambda}$. The Wilson action S_Λ is equal to $W_{B,\Lambda}$.

symbol	description
$S_B[\phi]$	the bare action with a UV cutoff Λ_0
$S_{I,B}[\phi]$	the interaction part of S_B
$W_B[J]$	the generating functional of S_B
$\Gamma_B[\Phi]$	the effective action of S_B
$S_B[\phi]$	$= -\frac{1}{2} \int \frac{p^2+m^2}{K_0} \phi\phi + S_{I,B}[\phi]$ (2.1)
$\exp[W_B[J]]$	$= \int [d\phi] \exp[S_B[\phi] + \int J\phi]$ (2.27)
$\Gamma_B[\Phi]$	$= W_B[J] - \int J\Phi$, where $\begin{cases} \frac{\delta W_B[J]}{\delta J(-p)} = \Phi(p) \\ \frac{\delta \Gamma_B[\Phi]}{\delta \Phi(p)} = -J(-p) \end{cases}$
$S_A[\phi]$	the Wilson action with a UV cutoff Λ
$S_{I,A}[\phi]$	the interaction part of S_A
$W_A[J]$	the generating functional of S_A
$S_A[\phi]$	$= -\frac{1}{2} \int \frac{p^2+m^2}{K} \phi\phi + S_{I,A}[\phi]$ (2.25)
$\exp[S_{I,A}[\phi]]$	$= \int [d\phi'] \exp[-\frac{1}{2} \int \frac{p^2+m^2}{K_0-K} \phi'\phi' + S_{I,B}[\phi + \phi']]$ (2.23)
$\exp[W_A[J]]$	$= \int [d\phi] \exp[S_A[\phi] + \int J\phi]$ (2.28)
W_B and W_A are related as	
$W_B[J] = W_A\left[\frac{K_0}{K}J\right] - \frac{1}{2} \int \frac{K_0-K}{p^2+m^2} \frac{K_0}{K} JJ$ (2.29)	

Table I. The bare and Wilson actions — (2.29) shows that the generating functionals W_B and W_A are basically the same.

symbol	description
$S_{B,\Lambda}[\phi]$	the bare action with a UV cutoff Λ_0 and IR cutoff Λ
$W_{B,\Lambda}[J]$	the generating functional of $S_{B,\Lambda}$
$\Gamma_{B,\Lambda}[\Phi]$	the effective average action; the Legendre transform of $W_{B,\Lambda}$
$\Gamma_{I,B,\Lambda}[\Phi]$	the interaction part of $\Gamma_{B,\Lambda}$
$S_{B,\Lambda}[\phi]$	$= -\frac{1}{2} \int \frac{p^2+m^2}{K_0-K} \phi\phi + S_{I,B}[\phi]$ (2.56)
$\exp[W_{B,\Lambda}[J]]$	$= \int [d\phi] \exp[S_{B,\Lambda}[\phi] + \int J\phi]$ (2.63)
$\Gamma_{B,\Lambda}[\Phi]$	$= W_{B,\Lambda}[J] - \int J\Phi$, where $\begin{cases} \frac{\delta W_{B,\Lambda}[J]}{\delta J(-p)} = \Phi(p) \\ \frac{\delta \Gamma_{B,\Lambda}[\Phi]}{\delta \Phi(p)} = -J(-p) \end{cases}$ (2.65)
$\Gamma_{B,\Lambda}[\Phi]$	$= -\frac{1}{2} \int \frac{p^2+m^2}{K_0-K} \Phi\Phi + \Gamma_{I,B,\Lambda}[\Phi]$ (2.87)
the relation between $S_{I,\Lambda}$ and W_B	
$W_{B,\Lambda}[J] = S_{I,\Lambda}\left[\frac{K_0-K}{p^2+m^2}J\right] + \frac{1}{2} \int \frac{K_0-K}{p^2+m^2} JJ$ (2.78)	
the relation between $\Gamma_{I,B,\Lambda}$ and $S_{I,\Lambda}$	
$\Gamma_{I,B,\Lambda}[\Phi] = S_{I,\Lambda}[\phi] + \frac{1}{2} \int \frac{p^2+m^2}{K_0-K} (\Phi - \phi)(\Phi - \phi)$ (2.88)	
where $\begin{cases} \Phi(p) = \phi(p) + \frac{K_0-K}{p^2+m^2} \frac{\delta S_{I,\Lambda}[\phi]}{\delta \phi(-p)} \end{cases}$ (2.89)	
$\begin{cases} \phi(p) = \Phi(p) - \frac{K_0-K}{p^2+m^2} \frac{\delta \Gamma_{I,B,\Lambda}[\Phi]}{\delta \Phi(-p)} \end{cases}$ (2.90)	


Table II. We construct $S_{B,\Lambda}$ from S_B by cutting off the IR modes with momenta below Λ . The Wilson action S_A is the generating functional of $S_{B,\Lambda}$.

- the $\Lambda \rightarrow 0+$ and $\Lambda \rightarrow \Lambda_0-$ limits (Table III) — As we take the IR cutoff Λ to zero, $S_{B,\Lambda}$ becomes S_B . As we take Λ to Λ_0 , the interaction parts of S_A and $\Gamma_{B,\Lambda}$ become the interaction part of S_B .

$\Lambda \rightarrow 0+$ limit
$\lim_{\Lambda \rightarrow 0} S_{B,\Lambda} = S_B$
$\lim_{\Lambda \rightarrow 0} \Gamma_{B,\Lambda} = \Gamma_B$
$\lim_{\Lambda \rightarrow 0} W_{B,\Lambda} = W_B$
$\Lambda \rightarrow \Lambda_0 - 0$ limit
$\lim_{\Lambda \rightarrow \Lambda_0 - 0} S_{I,\Lambda}[\phi] = \lim_{\Lambda \rightarrow \Lambda_0 - 0} \Gamma_{I,B,\Lambda}[\phi] = S_{I,B}[\phi]$

 Table III. The two limits: $\Lambda \rightarrow 0+$ and $\Lambda \rightarrow \Lambda_0 - 0$

$$S_{I,B} = \sum_n \text{diagram with } 2n \text{ legs}$$

 Fig. 4. $S_{I,B}$ gives elementary vertices.

 Fig. 5. $\mathcal{V}_{2n}$ consists of all possible connected diagrams with $2n$ legs

2.6. Diagrammatic interpretation

Although ERG is not limited to perturbation theory, it is straightforward to derive all the results of the previous subsections using Feynman diagrams.

2.6.1. $S_{I,\Lambda}$ and the Polchinski equation

Let us first examine $S_{I,\Lambda}$, the interaction part of the Wilson action. It is defined by (2.23):

$$\exp[S_{I,\Lambda}[\phi]] \equiv \int [d\phi'] \exp \left[-\frac{1}{2} \int_p \frac{p^2 + m^2}{K_0 - K} \phi'(p) \phi'(-p) + S_{I,B}[\phi + \phi'] \right]. \quad (2.109)$$

We can compute $S_{I,\Lambda}$ using the high momentum propagator

$$\frac{K_0 - K}{p^2 + m^2} \quad (2.110)$$

and the elementary vertices given by $S_{I,B}$. (Fig. 4) If we expand

$$S_{I,\Lambda}[\phi] = \sum_{n=1}^{\infty} \frac{1}{(2n)!} \int_{p_1, \dots, p_{2n}} \phi(p_1) \cdots \phi(p_{2n}) \cdot (2\pi)^D \delta^{(D)}(p_1 + \cdots + p_{2n}) \times \mathcal{V}_{2n}(\Lambda; p_1, \dots, p_{2n}), \quad (2.111)$$

in powers of fields, the vertex function $\mathcal{V}_{2n}$ is obtained as the sum of all connected Feynman diagrams with $2n$ external legs. No propagator is assigned for the external leg. (Fig. 5)






Fig. 6. The linear coupling to the external field is interpreted as an interaction vertex.

Fig. 7. A J vertex attaches to $S_{I,\Lambda}[\phi]$ through a high momentum propagator.Fig. 8. Two J vertices connected by a high momentum propagator.

Now, let us consider $S_{I,\Lambda}[\phi; J]$ for the bare action $S_{I,B}[\phi; J]$ with a source J coupled linearly to ϕ . It is defined by

$$\exp[S_{I,\Lambda}[\phi; J]] \equiv \int [d\phi'] \exp \left[-\frac{1}{2} \int \frac{p^2 + m^2}{K_0 - K} \phi' \phi' + S_{I,B}[\phi + \phi'] + \int J \cdot (\phi + \phi') \right]. \quad (2.112)$$

The source gives an extra vertex given in Fig. 6. This extra vertex can do three things:

- (i) It attaches to one of the external lines of $\mathcal{V}_{2n}$ via a high momentum propagator as in Fig. 7. This gives a shift of ϕ :

$$S_{I,\Lambda} \left[\phi + \frac{K_0 - K}{p^2 + m^2} J \right]. \quad (2.113)$$

- (ii) Two of them get connected by a single high momentum propagator as in Fig. 8. This gives

$$\frac{1}{2} \int_p J(-p) \frac{K_0 - K}{p^2 + m^2} J(p). \quad (2.114)$$

- (iii) It remains as itself, giving



$$\int_p J(-p) \phi(p). \quad (2.115)$$

Hence, altogether we obtain (2.37), which we have called an intermediate result:

$$S_{I,\Lambda}[\phi; J] = S_{I,\Lambda} \left[\phi + \frac{K_0 - K}{p^2 + m^2} J \right] + \frac{1}{2} \int J \frac{K_0 - K}{p^2 + m^2} J + \int J \cdot \phi. \quad (2.116)$$

Now, let us consider the Λ derivative of the Wilson action $S_{I,\Lambda}$. The propagators in each Feynman diagram can be classified into two types:

1. **type 1** — if cut, the diagram breaks into two separate pieces.
2. **type 2** — if cut, the diagram remains a single piece.


 Fig. 9. Λ derivative acts on a type 1 propagator.

 Fig. 10. Λ derivative acts on a type 2 propagator.

When we differentiate $\mathcal{V}_{2n}$ with respect to Λ , only the propagators are acted on, since the vertices $S_{I,B}$ are independent of Λ . When differentiation acts on a type 1 propagator, we obtain two Feynman graphs connected to each other by the differentiated propagator:

$$-\Lambda \frac{\partial}{\partial \Lambda} \frac{K_0 - K}{p^2 + m^2} = \frac{\Delta(p/\Lambda)}{p^2 + m^2}. \quad (2.117)$$

Denoting this by a broken line, we obtain two vertex functions connected by a broken line as in Fig. 9. Summing over all possibilities, we obtain

$$\frac{1}{2} \int_p \frac{\Delta(p/\Lambda)}{p^2 + m^2} \frac{\delta S_{I,\Lambda}}{\delta \phi(-p)} \frac{\delta S_{I,\Lambda}}{\delta \phi(p)}, \quad (2.118)$$

where the factor $\frac{1}{2}$ avoids overcounting.

When differentiation acts on a type 2 propagator, we get a single vertex function with a loop of the differentiated propagator as in Fig. 10. Summing over all possibilities, we obtain

$$\frac{1}{2} \int_p \frac{\Delta(p/\Lambda)}{p^2 + m^2} \frac{\delta^2 S_{I,\Lambda}}{\delta \phi(p) \delta \phi(-p)}. \quad (2.119)$$


Thus, altogether we obtain the Polchinski equation

$$-\Lambda \frac{\partial}{\partial \Lambda} S_{I,\Lambda}[\phi] = \int_p \frac{\Delta(p/\Lambda)}{p^2 + m^2} \left\{ \frac{\delta S_{I,\Lambda}}{\delta \phi(p)} \frac{\delta S_{I,\Lambda}}{\delta \phi(-p)} + \frac{\delta^2 S_{I,\Lambda}}{\delta \phi(p) \delta \phi(-p)} \right\}. \quad (2.120)$$

2.6.2. $\Gamma_{I,B,\Lambda}$ and the Wetterich equation

The average action $\Gamma_{B,\Lambda}$ is the Legendre transform of the generating functional $W_{B,\Lambda}$. Expanding the interaction part $\Gamma_{I,B,\Lambda}$ of the average action as

$$\begin{aligned} \Gamma_{I,B,\Lambda}[\Phi] &= \sum_{n=1}^{\infty} \frac{1}{(2n)!} \int_{p_1, \dots, p_{2n}} \Phi(p_1) \cdots \Phi(p_{2n}) \cdot (2\pi)^D \delta^{(D)}(p_1 + \cdots + p_{2n}) \\ &\quad \times \Gamma_{2n}(\Lambda; p_1, \dots, p_{2n}). \end{aligned} \quad (2.121)$$

Fig. 11. Γ_{2n} consists of 1PI diagrams with $2n$ legsFig. 12. A series of 1PI diagrams giving $G(p, -q)$

we find that Γ_{2n} consists of all 1PI (one particle irreducible) diagrams with $2n$ external legs. Again, no propagator is assigned to the external legs. All the internal lines are of type 2. We denote Γ_{2n} by a shaded blob as in Fig. 11.

Differentiating $\Gamma_{2n}(\Lambda)$ with respect to Λ , we do not get graphs as in Fig. 9, but only graphs as in Fig. 10. The graph after the cutting is not necessarily 1PI; it consists of multiple 1PI graphs connected in series as in Fig. 12. Hence, we obtain the graphical equation

$$-\Lambda \frac{\partial}{\partial \Lambda} \text{blob} = \text{blob} \text{---} \text{blob} \text{---} \text{blob}$$

where the broken line on the top denotes the Λ derivative of the propagator.

Consider the sum over graphs in Fig. 12, where we assign a field Φ for each external leg except for those with momentum p or $-q$. Let us call the sum $G(p, -q)$ so that

$$-\Lambda \frac{\partial}{\partial \Lambda} \Gamma_{I,B,\Lambda}[\Phi] = \int_p \frac{\Delta(p/\Lambda)}{p^2 + m^2} G(p, -p). \quad (2.122)$$

Each blob of Fig. 12 gives

$$\Gamma_{I,B,\Lambda}^{(2)}(q, -r) \equiv \frac{\delta^2 \Gamma_{I,B,\Lambda}[\Phi]}{\delta \Phi(-q) \delta \Phi(r)}. \quad (2.123)$$

Hence, $G(p, -q)$ is a geometric series given by

$$G(p, -q) = \Gamma_{I,B,\Lambda}^{(2)}(p, -q) + \int_r \Gamma_{I,B,\Lambda}^{(2)}(p, -r) \frac{K_0 - K}{r^2 + m^2} \Gamma_{I,B,\Lambda}^{(2)}(r, -q) + \cdots \quad (2.124)$$

Hence, we obtain

$$\begin{aligned} & \int_r \left((2\pi)^D \delta^{(D)}(p - r) + \frac{K_0 - K}{p^2 + m^2} G(p, -r) \right) \frac{K_0 - K}{r^2 + m^2} \\ & \times \left(-\frac{r^2 + m^2}{K_0 - K} (2\pi)^D \delta^{(D)}(r - q) + \Gamma_{I,B,\Lambda}^{(2)}(r, -q) \right) = -(2\pi)^D \delta^{(D)}(p - q). \end{aligned} \quad (2.125)$$

Therefore, from (2.97), we find

$$\left((2\pi)^D \delta^{(D)}(p - q) + \frac{K_0 - K}{p^2 + m^2} G(p, -q) \right) \frac{K_0 - K}{q^2 + m^2} = - \left(\Gamma_{B,\Lambda}^{(2)} \right)^{-1}(p, -q). \quad (2.126)$$

Thus, ignoring an additive constant proportional to $\delta^{(D)}(0)$, we obtain

$$\begin{aligned} -\Lambda \frac{\partial}{\partial \Lambda} \Gamma_{I,B,\Lambda}[\Phi] &= \int_p (p^2 + m^2) \frac{\Delta(p/\Lambda)}{(K_0 - K)^2} \left(\frac{K_0 - K}{p^2 + m^2} \right)^2 G(p, -p) \\ &= - \int_p (p^2 + m^2) \frac{\Delta(p/\Lambda)}{(K_0 - K)^2} \left(\Gamma_{B,\Lambda}^{(2)} \right)^{-1}(p, -p), \end{aligned} \quad (2.127)$$

which is the Wetterich equation.

2.6.3. Λ dependence of the generating functional

Let us understand (2.29)

$$W_B[J] = W_\Lambda \left[\frac{K_0}{K} J \right] - \frac{1}{2} \int_p \frac{K_0 - K}{p^2 + m^2} \frac{K_0}{K} J(-p) J(p), \quad (2.128)$$

using Feynman diagrams. Recall that this is an important equation implying we miss no physics by the ERG transformation.

Let us first consider $W_B[J]$. This can be calculated perturbatively using the vertices provided by $S_{I,B}$ and the source term $\int J\phi$. The propagator is given by

$$\frac{K_0}{p^2 + m^2}. \quad (2.129)$$

Note that there is a single contribution that does not involve the interaction vertex $S_{I,B}$, which is given graphically by Fig. 8 with the above propagator:

$$\frac{1}{2} \int_p J(-p) \frac{K_0}{p^2 + m^2} J(p). \quad (2.130)$$

Now, let us consider the Feynman diagrams contributing to

$$W_B[J] - \frac{1}{2} \int_p J(-p) \frac{K_0}{p^2 + m^2} J(p). \quad (2.131)$$

Each internal propagator can be decomposed as the sum of high and low momentum propagators:

$$\frac{K_0}{p^2 + m^2} = \frac{K_0 - K}{p^2 + m^2} + \frac{K}{p^2 + m^2}. \quad (2.132)$$

We substitute the above decomposition to all the internal propagators. Those parts involving only the high momentum propagators give the vertices of the Wilson action $S_{I,\Lambda}$. The low momentum propagator is to be used with the vertices $S_{I,\Lambda}$. (See Fig. 13.) For the external propagators ending with sources, we write

$$\frac{K_0}{p^2 + m^2} J(p) = \frac{K}{p^2 + m^2} \cdot \frac{K_0}{K} J(p), \quad (2.133)$$

using the low momentum propagator. This amounts to replacing J by $K_0/K \cdot J$. Hence, the decomposition (2.132) gives

$$W_B[J] - \frac{1}{2} \int_p J \frac{K_0}{p^2 + m^2} J = W_\Lambda \left[\frac{K_0}{K} J \right] - \frac{1}{2} \int_p \frac{K_0}{K} J \frac{K}{p^2 + m^2} \frac{K_0}{K} J, \quad (2.134)$$

where we have subtracted the contributions independent of interaction vertices. From this, we obtain (2.29).




Fig. 13. On the right, the blobs contain only high momentum propagators. These blobs are connected to each other by low momentum propagators.

2.7. Fermions

It is straightforward to extend ERG to include spin $\frac{1}{2}$ fields. We enumerate the main results:

1. Given a bare action

$$S_B[\psi, \bar{\psi}] \equiv - \int_p \bar{\psi}(-p) \frac{\not{p} + im}{K_0} \psi(p) + S_{I,B}[\psi, \bar{\psi}], \quad (2.135)$$

the Wilson action is defined by

$$\begin{aligned} \exp [S_{I,A}[\psi, \bar{\psi}]] &\equiv \int [d\psi' d\bar{\psi}'] \exp \left[\right. \\ &\quad \left. - \int_p \bar{\psi}'(-p) \frac{\not{p} + im}{K_0 - K} \psi'(p) + S_{I,B}[\psi + \psi', \bar{\psi} + \bar{\psi}'] \right]. \end{aligned} \quad (2.136)$$

2. The Polchinski equation is obtained as

$$\begin{aligned} -\Lambda \frac{\partial}{\partial \Lambda} S_{I,A} &= (-) \int_p \frac{\Delta(p/\Lambda)}{p^2 + m^2} \cdot \text{Tr} \left[(\not{p} - im) \right. \\ &\quad \left. \times \left\{ \frac{\overrightarrow{\delta}}{\delta \bar{\psi}(-p)} S_{I,A} \cdot S_{I,A} \frac{\overleftarrow{\delta}}{\delta \psi(p)} + \frac{\overrightarrow{\delta}}{\delta \bar{\psi}(-p)} S_{I,A} \frac{\overleftarrow{\delta}}{\delta \psi(p)} \right\} \right], \end{aligned} \quad (2.137)$$

where the minus sign $(-)$ is due to the Fermi statistics.

3. By introducing anticommuting sources to the bare action

$$+ \int_p (\bar{\xi}(-p) \psi(p) + \bar{\psi}(-p) \eta(p)), \quad (2.138)$$

the Wilson action obtains the following source dependence:

$$\begin{aligned} S_{I,A}[\psi, \bar{\psi}; \bar{\xi}, \eta] &= S_{I,A} \left[\psi(p) + \frac{K_0 - K}{\not{p} + im} \eta(p), \bar{\psi}(-p) + \bar{\xi}(-p) \frac{K_0 - K}{\not{p} + im} \right] \\ &\quad + \int (\bar{\xi} \psi + \bar{\psi} \eta) + \int \bar{\xi}(-p) \frac{K_0 - K}{\not{p} + im} \eta(p). \end{aligned} \quad (2.139)$$

4. The generating functionals of the bare and Wilson actions are related by

$$\begin{aligned} W_B[\bar{\xi}, \eta] &- \int \bar{\xi} \frac{K_0}{\not{p} + im} \eta \\ &= W_A \left[\frac{K_0}{K} \bar{\xi}, \frac{K_0}{K} \eta \right] - \int \frac{K_0}{K} \bar{\xi}(-p) \frac{K}{\not{p} + im} \frac{K_0}{K} \eta(p). \end{aligned} \quad (2.140)$$

5. For the action with both UV and IR cutoffs

$$S_{B,\Lambda}[\psi, \bar{\psi}] \equiv - \int \bar{\psi}(-p) R_\Lambda(p) \psi(p) + S_B[\psi, \bar{\psi}], \quad (2.141)$$

where

$$R_\Lambda(p) \equiv (\not{p} + im) \left(\frac{1}{K_0 - K} - \frac{1}{K_0} \right), \quad (2.142)$$

we define the generating functional by

$$\exp [W_{B,\Lambda}[\bar{\xi}, \eta]] \equiv \exp \left[S_{B,\Lambda}[\psi, \bar{\psi}] + \int_p (\bar{\xi} \psi + \bar{\psi} \eta) \right]. \quad (2.143)$$

This is given in terms of the Wilson action by

$$W_{B,\Lambda}[\bar{\xi}, \eta] = S_{I,\Lambda} \left[\bar{\xi} \frac{K_0 - K}{\not{p} + im}, \frac{K_0 - K}{\not{p} + im} \eta \right] + \int \bar{\xi}(-p) \frac{K_0 - K}{\not{p} + im} \eta(p). \quad (2.144)$$

6. The effective average action $\Gamma_{B,\Lambda}[\Psi, \bar{\Psi}]$ is defined as its Legendre transform:

$$\Gamma_{B,\Lambda}[\Psi, \bar{\Psi}] \equiv W_{B,\Lambda}[\bar{\xi}, \eta] - \int_p (\bar{\xi} \Psi + \bar{\Psi} \eta), \quad (2.145)$$

where $\bar{\xi}, \eta$ are determined by

$$\begin{cases} \Psi(p) &= \frac{\overrightarrow{\delta}}{\delta \xi(-p)} W_{B,\Lambda}[\bar{\xi}, \eta] \\ \bar{\Psi}(-p) &= W_{B,\Lambda}[\bar{\xi}, \eta] \frac{\overleftarrow{\delta}}{\delta \eta(p)} \end{cases} \quad (2.146)$$

as functionals of $\Psi, \bar{\Psi}$.

7. We define $\Gamma_{B,\Lambda}^{(2)}$ by

$$\Gamma_{B,\Lambda}^{(2)}(p, -q) \equiv \frac{\overrightarrow{\delta}}{\delta \bar{\Psi}(-p)} \Gamma_{B,\Lambda}[\Psi, \bar{\Psi}] \frac{\overleftarrow{\delta}}{\delta \Psi(q)}, \quad (2.147)$$

and its inverse by

$$\int_q \left(\Gamma_{B,\Lambda}^{(2)} \right)^{-1}(p, -q) \Gamma_{B,\Lambda}^{(2)}(q, -r) = (2\pi)^D \delta^{(D)}(p - r). \quad (2.148)$$

8. The Wetterich equation for the effective average action is given as

$$-\Lambda \frac{\partial}{\partial \Lambda} \Gamma_{I,B,\Lambda}[\Psi, \bar{\Psi}] = (-) \int_p \text{Tr} \left(-\Lambda \frac{\partial}{\partial \Lambda} R_\Lambda(p) \right) \left(\Gamma_{B,\Lambda}^{(2)} \right)^{-1}(p, -p), \quad (2.149)$$

where

$$-\Lambda \frac{\partial}{\partial \Lambda} R_\Lambda(p) = -(\not{p} + im) \frac{\Delta(p/\Lambda)}{(K_0 - K)^2}. \quad (2.150)$$

§3. Continuum limits

In §2 we have kept the UV cutoff Λ_0 finite. In this section we consider taking the continuum limit $\Lambda_0 \rightarrow \infty$. We call the theory given by a bare action S_B **renormalizable** if we can obtain a limit of the Wilson action S_A as $\Lambda_0 \rightarrow \infty$, by giving appropriate Λ_0 dependence to the parameters of S_B . Since the Wilson action S_A contains the same physics as the bare action S_B , we expect that the physics of the continuum limit is fully contained in the $\Lambda_0 \rightarrow \infty$ limit of S_A . In this section we restrict our discussion only to perturbative renormalization, even though ERG is applicable non-perturbatively.

3.1. Perturbative renormalizability

For concreteness, let us consider the ϕ^4 theory in $D = 4$. We give the bare action in the following form:

$$S_B = -\frac{1}{2} \int_p \frac{p^2 + m^2}{K(p/\Lambda_0)} \phi(-p) \phi(p) + S_{I,B}, \quad (3.1)$$

where the interaction part has three terms:

$$S_{I,B} = - \int d^4x \left[\Delta m^2 \frac{\phi^2}{2} + \Delta z \frac{1}{2} \partial_\mu \phi \partial_\mu \phi + \lambda_0 \frac{\phi^4}{4!} \right]. \quad (3.2)$$

We determine the cutoff dependence of the coefficients $\Delta m^2, \Delta z, \lambda_0$ so that the Wilson action S_A has a finite limit as $\Lambda_0 \rightarrow \infty$:

$$\bar{S}_A \equiv \lim_{\Lambda_0 \rightarrow \infty} S_A, \quad (3.3)$$

where S_A is the solution of the Polchinski equation satisfying the initial condition $S_{\Lambda_0} = S_B$.^{*)} The Λ_0 dependence of S_A must go away in the limit. (Fig. 14) In fact it was for this proof of renormalizability that Polchinski's differential equation was first introduced.^{94)**} The idea is not to look at the continuum limit of the correlation functions, but to look instead at the limit of the Wilson action. It is beyond the scope of this review, however, to explain how to use the Polchinski equation to prove perturbative renormalizability. We refer the reader to the original paper⁹⁴⁾ and the references cited in §1.

We now recall an important result (2.29) from §2. Taking the limit $\Lambda_0 \rightarrow \infty$, we obtain

$$W_\infty[J] = W_A \left[\frac{J}{K} \right] + \frac{1}{2} \int_p \frac{1 - 1/K}{p^2 + m^2} J(-p) J(p), \quad (3.4)$$

^{*)} Here, we denote the continuum limit of the Wilson action putting a bar above S_A . This notation is used only in §3, and the continuum limit is simply written as S_A in the later sections. The bar notation is adopted again in §7 to denote a Wilson action in the presence of antifields.

^{**) The first differential equation for ERG, introduced by K. Wilson,¹²⁴⁾ is somewhat different from Polchinski's, which suits perturbation theory better. In Appendix A we derive the original ERG differential equation by Wilson.}




Fig. 14. S_A is the solution of the Polchinski equation with the initial condition $S_{\Lambda_0} = S_B$. Renormalizability amounts to the existence of a limit of S_A as we take Λ_0 to infinity.

where $W_\infty[J]$ is the generating functional of connected correlation functions in the continuum limit. Equivalently, by differentiating the above with respect to J 's (or taking the limit $\Lambda_0 \rightarrow \infty$ of (2.43)), we obtain

$$\begin{cases} \langle \phi(p)\phi(-p) \rangle^\infty = \frac{1}{K^2} \langle \phi(p)\phi(-p) \rangle_{\bar{S}_A} + \frac{1-1/K}{p^2+m^2}, \\ \langle \phi(p_1) \cdots \phi(p_n) \rangle^\infty = \prod_{i=1}^n \frac{1}{K(p_i/\Lambda)} \cdot \langle \phi(p_1) \cdots \phi(p_n) \rangle_{\bar{S}_A}, \end{cases} \quad (3.5)$$

where we denote the continuum limit by a superscript ∞ . These imply that the correlation functions of the continuum limit are fully obtained from the Wilson action $\bar{S}_A$ which has a finite momentum cutoff. Hence, if we expect some symmetry in the continuum limit, it should be a symmetry of the Wilson action. The above two relations (3.4) & (3.5) play a key role justifying our approach to realization of symmetry.

In the remaining part of this subsection, let us compute the two-point vertex function of $S_{I,A}$ at 1-loop level. This is only for an illustration; it is no substitute for a general proof of renormalizability using the Polchinski equation. Let us first recall the expansion of the Wilson action in powers of fields:

$$S_A[\phi] = \sum_{n=1}^{\infty} \frac{1}{(2n)!} \int_{p_1+\cdots+p_{2n}=0} \mathcal{V}_{2n}(\Lambda; p_1, \dots, p_{2n}) \phi(p_1) \cdots \phi(p_{2n}). \quad (3.6)$$

We wish to compute $\mathcal{V}_2$ at 1-loop. Denoting the 1-loop contribution by a superscript $^{(1)}$, we obtain

$$\mathcal{V}_2^{(1)}(\Lambda; p, -p) = -\frac{\lambda}{2} \int_q \frac{K(q/\Lambda_0) - K(q/\Lambda)}{q^2 + m^2} - (\Delta m^2)^{(1)}, \quad (3.7)$$

where λ is the tree level part of λ_0 . (Fig. 15) This is independent of p . We evaluate the integral as follows:

$$\int_q \frac{K(q/\Lambda_0) - K(q/\Lambda)}{q^2 + m^2} = \int_q (K(q/\Lambda_0) - K(q/\Lambda)) \left(\frac{1}{q^2} - \frac{m^2}{q^4} + \frac{m^4}{q^4(q^2 + m^2)} \right)$$




Fig. 15. Two-point vertex at 1-loop is calculated with a high momentum propagator. The second graph denotes a counterterm $-(\Delta m^2)^{(1)}$.

$$\begin{aligned}
&= \int_q \frac{K(q/\Lambda_0) - K(q/\Lambda)}{q^2} - m^2 \int_q \frac{K(q/\Lambda_0) - K(q/\Lambda)}{q^4} \\
&\quad + \int_q (K(q/\Lambda_0) - K(q/\Lambda)) \frac{m^4}{q^4(q^2 + m^2)}. \tag{3.8}
\end{aligned}$$

The last integral has a finite limit:

$$\lim_{\Lambda_0 \rightarrow \infty} \int_q (K(q/\Lambda_0) - K(q/\Lambda)) \frac{m^4}{q^4(q^2 + m^2)} = \int_q (1 - K(q/\Lambda)) \frac{m^4}{q^4(q^2 + m^2)}. \tag{3.9}$$

The first integral gives^{*)}

$$\int_q \frac{K(q/\Lambda_0) - K(q/\Lambda)}{q^2} = (\Lambda_0^2 - \Lambda^2) \int_q \frac{K(q)}{q^2}, \tag{3.10}$$

where the integral on the right-hand side is a finite constant, dependent on the particular cutoff function chosen. It needs a little more work to calculate the second integral. Let

$$F(\Lambda_0, \Lambda) \equiv \int_q \frac{K(q/\Lambda_0) - K(q/\Lambda)}{q^4} \tag{3.11}$$

so that

$$F(\Lambda, \Lambda) = 0. \tag{3.12}$$

Differentiating F with respect to Λ_0 , we obtain

$$\Lambda_0 \frac{\partial}{\partial \Lambda_0} F(\Lambda_0, \Lambda) = \int_q \frac{\Delta(q/\Lambda_0)}{q^4} = \int_q \frac{\Delta(q)}{q^4} \tag{3.13}$$

which is a constant. This constant is independent of the choice of K due to the following general formula:

$$\int_q \frac{\Delta(q) (1 - K(q))^n}{q^4} = \frac{2}{(4\pi)^2} \frac{1}{n+1}. \tag{3.14}$$

(Proof) Using the spherical coordinates and the definition $\Delta(q) \equiv -2q^2 dK(q)/dq^2$, we calculate

$$\int_q \frac{\Delta(q) (1 - K(q))^n}{q^4} = \frac{2\pi^2}{(2\pi)^4} \int_0^\infty \frac{q^2 dq^2}{2} (-2q^2) \frac{dK}{dq^2} (1 - K)^n \frac{1}{q^4}$$

^{*)} Please note that we are not using the short-hand notation (2.24) of §2 here; $K(q)$ on the right does not stand for $K(q/\Lambda)$. The dimensionless argument q is obtained from q/Λ or q/Λ_0 by change of variables. In this section, we do not use the short-hand notation.

$$\begin{aligned}
&= \frac{2}{(4\pi)^2} \int_0^\infty dq^2 \frac{1}{n+1} \frac{d}{dq^2} (1-K)^{n+1} \\
&= \frac{2}{(4\pi)^2} \frac{1}{n+1},
\end{aligned} \tag{3-15}$$

using $K(0) = 1$ and $K(\infty) = 0$. (End of proof)

Hence, we obtain

$$\Lambda_0 \frac{\partial}{\partial \Lambda_0} F(\Lambda_0, \Lambda) = \frac{2}{(4\pi)^2}. \tag{3-16}$$

Thus,

$$F(\Lambda_0, \Lambda) = \frac{2}{(4\pi)^2} \ln \frac{\Lambda_0}{\Lambda}. \tag{3-17}$$

Altogether, we obtain

$$\begin{aligned}
\mathcal{V}_2^{(1)}(\Lambda) &= -\frac{\lambda}{2} \left[(\Lambda_0^2 - \Lambda^2) \int_q \frac{K(q)}{q^2} - m^2 \frac{2}{(4\pi)^2} \ln \frac{\Lambda_0}{\Lambda} + \frac{m^4}{\Lambda^2} \int_q \frac{1-K(q)}{q^4 (q^2 + m^2/\Lambda^2)} \right] \\
&\quad - (\Delta m^2)^{(1)}.
\end{aligned} \tag{3-18}$$

For this to have a limit as $\Lambda_0 \rightarrow \infty$, we take

$$(\Delta m^2)^{(1)} = \frac{\lambda}{2} \left(-\Lambda_0^2 \int_q \frac{K(q)}{q^2} + m^2 \frac{2}{(4\pi)^2} \ln \frac{\Lambda_0}{\mu} \right), \tag{3-19}$$

where we have introduced an arbitrary momentum scale μ to make the argument of the logarithm dimensionless. This gives the continuum limit

$$\bar{\mathcal{V}}_2^{(1)}(\Lambda) = -\frac{\lambda}{2} \left[-\Lambda^2 \int_q \frac{K(q)}{q^2} + m^2 \frac{2}{(4\pi)^2} \ln \frac{\Lambda}{\mu} + \frac{m^4}{\Lambda^2} \int_q \frac{1-K(q)}{q^4 (q^2 + m^2/\Lambda^2)} \right] \tag{3-20}$$

as the 1-loop self-energy correction from the high-momentum modes.

In addition, the contribution of the low-momentum modes is given by

$$-\frac{\lambda}{2} \int_q \frac{K(q/\Lambda)}{q^2 + m^2}. \tag{3-21}$$

Hence, the 1-loop self-energy correction is

$$-\Sigma^{(1)} \equiv \bar{\mathcal{V}}_2^{(1)}(\Lambda) - \frac{\lambda}{2} \int_q \frac{K(q/\Lambda)}{q^2 + m^2}, \tag{3-22}$$

which is independent not only of Λ , but also of the external momentum. Thus, up to 1-loop, the inverse propagator is given by

$$1/\langle \phi(p)\phi(-p) \rangle^\infty = p^2 + m^2 + \Sigma^{(1)} = p^2 + m_{\text{ph}}^2, \tag{3-23}$$

where we have defined the physical squared mass m_{ph}^2 by

$$m_{\text{ph}}^2 \equiv m^2 + \Sigma^{(1)} = m^2 - \bar{\mathcal{V}}_2^{(1)}(\Lambda) + \frac{\lambda}{2} \int_p \frac{K(p/\Lambda)}{p^2 + m^2}. \tag{3-24}$$




Fig. 16. The Wilson action $\bar{S}_\Lambda$ of the continuum limit obeys its own Polchinski differential equation and asymptotic conditions.

This is independent of Λ , but depends on μ .

To all orders in perturbation theory, we expect the following Λ_0 dependence of the counterterms:

$$\Delta m^2 = \Lambda_0^2 A_2(\ln \Lambda_0/\mu) + m^2 B_2(\ln \Lambda_0/\mu) = O(\lambda) \quad (3.25)$$

$$\Delta z = C_2(\ln \Lambda_0/\mu) = O(\lambda^2) \quad (3.26)$$

$$\lambda_0 = A_4(\ln \Lambda_0/\mu) = \lambda + O(\lambda^2) \quad (3.27)$$

The Λ_0 dependence is determined so that S_Λ , for a fixed Λ , has a limit as $\Lambda_0 \rightarrow \infty$. To fix the Λ_0 independent additive constants in B_2, C_2, A_4 , we must provide three renormalization conditions. For example, we can adopt the BPHZ condition at zero momentum so that

$$1/\langle \phi(p)\phi(-p) \rangle^\infty = m^2 + p^2 + p^2 \cdot O\left(\frac{p^2}{m^2}\right), \quad (3.28)$$

$$\langle \phi(0)\phi(0)\phi(0)\phi(0) \rangle^\infty = -\lambda. \quad (3.29)$$

At 1-loop, the BPHZ condition amounts to choosing μ so that

$$\Sigma^{(1)} = 0. \quad (3.30)$$

3.2. Asymptotic behaviors

For realization of symmetry, it is more convenient to parameterize the theory by directly specifying the “asymptotic” behavior of the continuum limit $\bar{S}_\Lambda$ as $\Lambda \rightarrow \infty$.^{(17), (91), (106)} (See Fig. 16.) To explain this parameterization, let us expand the continuum limit $\bar{S}_\Lambda$ of the Wilson action in powers of fields:

$$\bar{S}_{I,\Lambda}[\phi] = \sum_{n=1}^{\infty} \frac{1}{(2n)!} \int_{p_1+\dots+p_{2n}=0} \bar{\mathcal{V}}_{2n}(\Lambda; p_1, \dots, p_{2n}) \phi(p_1) \cdots \phi(p_{2n}). \quad (3.31)$$

The vertex functions $\bar{\mathcal{V}}_{2n}$ are local, meaning that we can expand them in powers of m^2 and the external momenta. Since $\bar{\mathcal{V}}_{2n}$ has mass dimension $4 - 2n$, we expect the

following expansions:

$$\begin{aligned}\bar{\mathcal{V}}_2(\Lambda; p, -p) &= \Lambda^2 a_2(\ln \Lambda/\mu) + m^2 b_2(\ln \Lambda/\mu) + p^2 c_2(\ln \Lambda/\mu) + \cdots, \\ \bar{\mathcal{V}}_4(\Lambda; p_1, \cdots, p_4) &= a_4(\ln \Lambda/\mu) + \cdots, \\ \bar{\mathcal{V}}_{2n}(\Lambda; p_1, \cdots, p_{2n}) &= \frac{1}{\Lambda^{2n-4}} a_{2n}(\ln \Lambda/\mu) + \cdots. \quad (2n \geq 6)\end{aligned}\tag{3.32}$$

At each order of loop expansions, the coefficients a_2 , b_2 , etc., are finite polynomials of the logarithm of Λ . The scale parameter μ is introduced to make the argument of the logarithm dimensionless, as has been done in the previous subsection. Since the Polchinski differential equation (2.51) determines only the Λ dependence, it leaves Λ independent parts undetermined. More explicitly, the undetermined Λ independent parts are additive constants in

$$b_2(\ln \Lambda/\mu), \quad c_2(\ln \Lambda/\mu), \quad a_4(\ln \Lambda/\mu).\tag{3.33}$$

We can fix these additive constants by specifying the values of b_2, c_2, a_4 at $\Lambda = \mu$. Once the additive constants are fixed, the vertex functions, including a_2 and $a_{2n \geq 6}$, are determined unambiguously by the Polchinski differential equation. Thus, we have argued that the Wilson action can be determined uniquely by imposing a convention on the three additive constants

$$b_2(0), \quad c_2(0), \quad a_4(0).\tag{3.34}$$

The simplest choice would be

$$b_2(0) = c_2(0) = 0, \quad a_4(0) = -\lambda,\tag{3.35}$$

which is an analog of the MS scheme for dimensional regularization.^{111)*)}

In lieu of expanding the vertex functions in powers of m^2 and external momenta, we often refer to the “asymptotic behavior” of the Wilson action:


$$\begin{aligned}\bar{S}_{I,\Lambda} \xrightarrow{\Lambda \rightarrow \infty} \int d^4x \left[(\Lambda^2 a_2(\ln \Lambda/\mu) + m^2 b_2(\ln \Lambda/\mu)) \frac{\phi^2}{2} \right. \\ \left. + c_2(\ln \Lambda/\mu) \frac{1}{2} \partial_\mu \phi \partial_\mu \phi + a_4(\ln \Lambda/\mu) \frac{1}{4!} \phi^4 \right].\end{aligned}\tag{3.36}$$

This is a convenient way of summarizing the leading terms of the expansions of the vertex functions in powers of m^2 and external momenta. The above “asymptotic” expansion is valid in two ways:

1. For fixed m^2 and external momenta, we take Λ asymptotically large.
2. For fixed Λ , we take m^2 and external momenta small.

Hence, calling (3.36) an asymptotic expansion is a bit imprecise. When we refer to asymptotic expansions, we often consider the second meaning. Please note that we only keep the part of the vertex functions multiplied by non-negative powers of Λ in the asymptotic behavior; we do not write a ϕ^6 term in the asymptotic behavior, since the leading term of $\bar{\mathcal{V}}_6$ is multiplied by $1/\Lambda^2$.

*) In Appendix B.1 we derive a mass independent RG equation using the choice (3.35).

Fig. 17. The s-channel contribution to $\bar{\mathcal{V}}_4^{(1)}$

For concreteness, let us consider the examples of two- and four-point vertex functions at 1-loop. First, let us consider the differential equation for the two-point vertex:

$$-\Lambda \frac{\partial}{\partial \Lambda} \bar{\mathcal{V}}_2^{(1)}(\Lambda) = -\frac{\lambda}{2} \int_q \frac{\Delta(q/\Lambda)}{q^2 + m^2} \quad (3.37)$$

Expanding the right-hand side in $\frac{m^2}{\Lambda^2}$, we obtain

$$-\Lambda \frac{\partial}{\partial \Lambda} \bar{\mathcal{V}}_2^{(1)}(\Lambda) = -\frac{\lambda}{2} \Lambda^2 \int_q \frac{\Delta(q)}{q^2} \left(1 - \frac{m^2}{\Lambda^2} \frac{1}{q^2} + \mathcal{O}\left(\frac{m^4}{\Lambda^4}\right) \right). \quad (3.38)$$

Using

$$\int_q \frac{\Delta(q)}{q^2} = 2 \int_q \frac{K(q)}{q^2}, \quad \int_q \frac{\Delta(q)}{q^4} = \frac{2}{(4\pi)^2}, \quad (3.39)$$

we obtain the asymptotic behavior

$$\bar{\mathcal{V}}_2^{(1)}(\Lambda) \xrightarrow{\Lambda \rightarrow \infty} -\frac{\lambda}{2} \Lambda^2 \left[-\frac{1}{2} \int_q \frac{\Delta(q)}{q^2} + \frac{2}{(4\pi)^2} \frac{m^2}{\Lambda^2} \ln \frac{\Lambda}{\mu} \right] + \text{const}, \quad (3.40)$$

where we have introduced $\ln \mu$ as part of the additive constant. Hence, we obtain

$$\begin{cases} a_2^{(1)} &= \frac{\lambda}{4} \int_q \frac{\Delta(q)}{q^2} = \frac{\lambda}{2} \int_q \frac{K(q)}{q^2}, \\ b_2^{(1)}(\ln \Lambda/\mu) &= -\frac{\lambda}{(4\pi)^2} \ln \frac{\Lambda}{\mu} + \text{const}. \end{cases} \quad (3.41)$$

Then, by imposing the convention

$$b_2^{(1)}(0) = 0, \quad (3.42)$$

we obtain the asymptotic behavior of the solution given by (3.20).

Let us next consider a 1-loop contribution to the four-point vertex function corresponding to the s-channel Feynman diagram in Fig. 17. In the rest of this section, we consider only the continuum limit, and for simplicity we omit a bar to denote the limit. Calling the s-channel vertex by $v_4(\Lambda; p_1 + p_2)$, we obtain the differential equation

$$-\Lambda \frac{\partial}{\partial \Lambda} v_4(\Lambda; p_1 + p_2) = \int_p \frac{\Delta(p/\Lambda)}{p^2 + m^2} v_6(\Lambda; p_1 + p_2 + q, p_3 + p_4 - q), \quad (3.43)$$

where v_6 is part of the tree-level six-point vertex function $\mathcal{V}_6^{(0)}(\Lambda; p_1, \dots, p_4, q, -q)$ corresponding to the right diagram in Fig. 18. Now, v_6 satisfies




Fig. 18. The left represents (3.43), where the broken line gives the propagator proportional to Δ . The right represents $v_6(\Lambda; p_1 + p_2 + q, p_3 + p_4 - q)$.

$$-\Lambda \frac{\partial}{\partial \Lambda} v_6(\Lambda; p, -p) = \lambda^2 \frac{\Delta(p/\Lambda)}{p^2 + m^2}, \quad (3.44)$$

and the asymptotic condition

$$v_6(\Lambda; p, -p) \xrightarrow{\Lambda \rightarrow \infty} 0, \quad (3.45)$$

since no six-point vertex survives the $\Lambda \rightarrow \infty$ limit.*) Hence, the solution is

$$v_6(\Lambda; p, -p) = \lambda^2 \frac{1 - K(p/\Lambda)}{p^2 + m^2}. \quad (3.46)$$

Substituting this into (3.43), we obtain

$$-\Lambda \frac{\partial}{\partial \Lambda} v_4(\Lambda; p) = \lambda^2 \int_p \frac{\Delta(p/\Lambda)}{p^2 + m^2} \frac{1 - K((q+p)/\Lambda)}{(q+p)^2 + m^2}. \quad (3.47)$$

Now, $v_4(\ln \Lambda/\mu; p)$ is determined by this differential equation and an additive constant in its asymptotic behavior. To find the asymptotic behavior, we consider the integral

$$\begin{aligned} \int_q \frac{\Delta(q/\Lambda)}{q^2 + m^2} \frac{1 - K((q+p)/\Lambda)}{(q+p)^2 + m^2} &= \int_q \frac{\Delta(q)}{q^2 + m^2/\Lambda^2} \frac{1 - K(q+p/\Lambda)}{(q+p/\Lambda)^2 + m^2/\Lambda^2} \\ &\xrightarrow{\Lambda \rightarrow \infty} \int_q \frac{\Delta(q)}{q^4} \frac{1 - K(q)}{q^4} = \frac{1}{(4\pi)^2}. \end{aligned} \quad (3.48)$$

Hence, the asymptotic behavior of v_4 is given by

$$v_4(\ln \Lambda/\mu; p) \xrightarrow{\Lambda \rightarrow \infty} -\frac{\lambda^2}{(4\pi)^2} \ln \frac{\Lambda}{\mu} + \text{const}. \quad (3.49)$$

This implies, after summing over three channels,

$$a_4^{(1)}(\ln \Lambda/\mu) = -\frac{3\lambda^2}{(4\pi)^2} \ln \frac{\Lambda}{\mu} + \text{const}. \quad (3.50)$$

*) Recall our terminology. Without referring to “asymptotic” behaviors, we can say the same thing as follows. Expanding $\mathcal{V}_6$ in powers of m^2 and momenta, the leading term is proportional to a negative power of Λ .

Now, we obtain

$$\begin{aligned} & -\Lambda \frac{\partial}{\partial \Lambda} \left(v_4(\ln \Lambda/\mu; p) + \frac{\lambda^2}{(4\pi)^2} \ln \frac{\Lambda}{\mu} \right) \\ &= \lambda^2 \int_q \left[\frac{\Delta(q/\Lambda)}{q^2 + m^2} \frac{1 - K((q+p)/\Lambda)}{(q+p)^2 + m^2} - \frac{\Delta(q/\Lambda)(1 - K(q/\Lambda))}{q^4} \right]. \end{aligned} \quad (3.51)$$

This is integrated to give

$$\begin{aligned} v_4(\ln \Lambda/\mu; p) &= -\frac{\lambda^2}{(4\pi)^2} \ln \frac{\Lambda}{\mu} + \text{const} \\ &+ \frac{\lambda^2}{2} \int_q \left[\frac{1 - K(q/\Lambda)}{q^2 + m^2} \frac{1 - K((q+p)/\Lambda)}{(q+p)^2 + m^2} - \frac{(1 - K(q/\Lambda))^2}{q^4} \right]. \end{aligned} \quad (3.52)$$

The integral on the right-hand side vanishes asymptotically, i.e., as we take m^2 and p to zero. The additive constant must be zero if we adopt the convention $a_4(0) = -\lambda$.

3.3. Asymptotic behaviors of the Wilson action and the effective average action

In §2.4 we have introduced the effective average action $\Gamma_{B,\Lambda}[\Phi]$ as the Legendre transform of the Wilson action $S_\Lambda[\phi]$. Their precise relation is given by (2.84) and (2.85). In the continuum limit $\Lambda_0 \rightarrow \infty$, let us write Γ_Λ for $\Gamma_{B,\Lambda}$. (We continue omitting a bar to denote the continuum limit.) Then, we can write (2.84) and (2.85) as

$$\Gamma_\Lambda[\Phi] = S_\Lambda[\phi] + \int_p \frac{p^2 + m^2}{1 - K(p/\Lambda)} \left(\frac{1}{K(p/\Lambda)} \frac{1}{2} \phi(p)\phi(-p) - \Phi(p)\phi(-p) \right) \quad (3.53)$$

and

$$\Phi(p) = \frac{1}{K(p/\Lambda)} \phi(p) + \frac{1 - K(p/\Lambda)}{p^2 + m^2} \frac{\delta S_\Lambda[\phi]}{\delta \phi(-p)}. \quad (3.54)$$

For the interaction parts defined by

$$\begin{cases} S_{I,\Lambda}[\phi] &\equiv \frac{1}{2} \int_p \frac{p^2 + m^2}{K(p/\Lambda)} \phi(-p)\phi(p) + S_\Lambda[\phi], \\ \Gamma_{I,\Lambda}[\Phi] &\equiv \frac{1}{2} \int_p \frac{p^2 + m^2}{1 - K(p/\Lambda)} \Phi(-p)\Phi(p) + \Gamma_\Lambda[\Phi], \end{cases} \quad (3.55)$$

we can write (3.53) as

$$\Gamma_{I,\Lambda}[\Phi] = S_{I,\Lambda}[\phi] + \frac{1}{2} \int_p \frac{p^2 + m^2}{1 - K(p/\Lambda)} (\Phi - \phi)(-p)(\Phi - \phi)(p). \quad (3.56)$$

Since Φ and ϕ become equal as $\Lambda \rightarrow \infty$, (3.56) implies that $\Gamma_{I,\Lambda}[\Phi]$ and $S_{I,\Lambda}[\phi]$ have the same asymptotic behaviors. In the case of the ϕ^4 theory in $D = 4$, the asymptotic behavior (3.36) of the Wilson action implies^{*)}

$$\begin{aligned} \Gamma_{I,\Lambda}[\Phi] &\xrightarrow{\Lambda \rightarrow \infty} \int d^4x \left[(\Lambda^2 a_2(\ln \Lambda/\mu) + m^2 b_2(\ln \Lambda/\mu)) \frac{\Phi^2}{2} \right. \\ &\quad \left. + c_2(\ln \Lambda/\mu) \frac{1}{2} (\partial_\mu \Phi)^2 + a_4(\ln \Lambda/\mu) \frac{\Phi^4}{4!} \right]. \end{aligned} \quad (3.57)$$

^{*)} This is the parameterization adopted in ref. 91), where μ is denoted as Λ_R .

3.4. Alternative parameterization

We have shown how to specify the Wilson action of the ϕ^4 theory in $D = 4$ by specifying $b_2(0), c_2(0), a_4(0)$ by hand. Alternatively, we can specify the behavior of the two- and four-point vertex functions at zero external momentum as an analog of the BPHZ scheme. More concretely, we can impose

$$\mathcal{V}_2(\Lambda = \mu; p = 0) = 0, \quad (3.58)$$

$$\frac{\partial}{\partial p^2} \mathcal{V}_2(\Lambda = \mu; p, -p) \Big|_{p=0} = 0, \quad (3.59)$$

$$\mathcal{V}_4(\Lambda = \mu; p_i = 0) = -\lambda, \quad (3.60)$$

for a given choice of the renormalization scale μ .

Let us see how the above conditions change the choice of $b_2(0), a_4(0)$ at 1-loop. The two-point vertex function becomes

$$\mathcal{V}_2^{(1)}(\Lambda) = -\frac{\lambda}{2} \left[(-\Lambda^2 + \mu^2) \int_q \frac{K(q)}{q^2} + m^2 \frac{2}{(4\pi)^2} \ln \frac{\Lambda}{\mu} + m^4 \int_q \frac{K(q/\mu) - K(q/\Lambda)}{q^4(q^2 + m^2)} \right]. \quad (3.61)$$

Hence,

$$b_2(0) = -\frac{\lambda}{2} \left[\frac{\mu^2}{m^2} \int_q \frac{K(q)}{q^2} + m^2 \int_q \frac{K(q/\mu) - 1}{q^4(q^2 + m^2)} \right]. \quad (3.62)$$

Note the non-trivial dependence on m^2 .

The four-point vertex function, for a single channel, becomes

$$v_4(\ln \Lambda/\mu; p) = \frac{\lambda^2}{2} \int_q \left[\frac{1 - K(q/\Lambda)}{q^2 + m^2} \frac{1 - K((q+p)/\Lambda)}{(q+p)^2 + m^2} - \left(\frac{1 - K(q/\mu)}{q^2 + m^2} \right)^2 \right], \quad (3.63)$$

where we have used

$$\int_q \left[\left(\frac{1 - K(q/\mu)}{q^2} \right)^2 - \left(\frac{1 - K(q/\Lambda)}{q^2} \right)^2 \right] = \frac{2}{(4\pi)^2} \ln \frac{\Lambda}{\mu}. \quad (3.64)$$

Hence,

$$a_4(0) = -\lambda + 3 \times \frac{\lambda^2}{2} \int_q (1 - K(q/\mu))^2 \left[\frac{1}{q^4} - \frac{1}{(q^2 + m^2)^2} \right]. \quad (3.65)$$

where the factor 3 is for three channels (s,t,u). This also has non-trivial dependence on m^2 .^{*)}

In general, a renormalization scheme corresponds to a convention for the coefficients $b_2(0), c_2(0), a_4(0)$. In Appendix B.2, we will show that changing the three coefficients amounts to changing m^2, λ , and the normalization of ϕ under the same renormalization scheme.

^{*)} The non-trivial dependence of $b_2(0), a_4(0)$ on $\frac{m^2}{\mu^2}$ implies that the beta function and anomalous dimension of the squared mass, explained in Appendix B.1, are mass dependent.

§4. Composite operators

In this section we introduce **composite operators**,¹⁷⁾ which provide the most important tool for our program of symmetry realization. By the name we do not mean any random functionals of field variables. Only when a cutoff dependent functional satisfies a particular property, we call it a composite operator.

4.1. Definition

Let ΔS_Λ be an infinitesimal deformation of a Wilson action S_Λ so that both S_Λ and $S_\Lambda + \Delta S_\Lambda$ satisfy the same Polchinski equation. Then, the difference satisfies the following linear equation:

$$-\Lambda \frac{\partial}{\partial \Lambda} \Delta S_\Lambda = \int_p \frac{\Delta(p/\Lambda)}{p^2 + m^2} \left[\frac{p^2 + m^2}{K(p/\Lambda)} \phi(p) \frac{\delta}{\delta \phi(p)} + \frac{\delta S_\Lambda}{\delta \phi(-p)} \frac{\delta}{\delta \phi(p)} + \frac{1}{2} \frac{\delta^2}{\delta \phi(-p) \delta \phi(p)} \right] \Delta S_\Lambda. \quad (4.1)$$

This can be rewritten using the interaction part $S_{I,\Lambda}$ as

$$-\Lambda \frac{\partial}{\partial \Lambda} \Delta S_\Lambda = \int_p \frac{\Delta(p/\Lambda)}{p^2 + m^2} \left[\frac{\delta S_{I,\Lambda}}{\delta \phi(-p)} \frac{\delta}{\delta \phi(p)} + \frac{1}{2} \frac{\delta^2}{\delta \phi(-p) \delta \phi(p)} \right] \Delta S_\Lambda. \quad (4.2)$$

In general we call a Λ dependent functional $\mathcal{O}_\Lambda[\phi]$ a composite operator if it satisfies the same differential equation as above:

$$\begin{aligned} -\Lambda \frac{\partial}{\partial \Lambda} \mathcal{O}_\Lambda &= \int_p \frac{\Delta(p/\Lambda)}{p^2 + m^2} \left[\frac{p^2 + m^2}{K(p/\Lambda)} \phi(p) \frac{\delta}{\delta \phi(p)} + \frac{\delta S_\Lambda}{\delta \phi(-p)} \frac{\delta}{\delta \phi(p)} + \frac{1}{2} \frac{\delta^2}{\delta \phi(-p) \delta \phi(p)} \right] \mathcal{O}_\Lambda \\ &= \int_p \frac{\Delta(p/\Lambda)}{p^2 + m^2} \left[\frac{\delta S_{I,\Lambda}}{\delta \phi(-p)} \frac{\delta}{\delta \phi(p)} + \frac{1}{2} \frac{\delta^2}{\delta \phi(-p) \delta \phi(p)} \right] \mathcal{O}_\Lambda. \end{aligned} \quad (4.3)$$

In other words, any Λ dependent functional $\mathcal{O}_\Lambda[\phi]$ is a composite operator, if

$$S_{I,\Lambda} + \epsilon \mathcal{O}_\Lambda \quad (4.4)$$

satisfies the Polchinski equation up to first order in ϵ . Note that the right-hand side of (4.3) is a linear differential operator acting on $\mathcal{O}_\Lambda$. Hence, denoting the differential operator by $\mathcal{D}$, we may express (4.3) symbolically as

$$-\Lambda \frac{\partial}{\partial \Lambda} \mathcal{O}_\Lambda = \mathcal{D} \cdot \mathcal{O}_\Lambda. \quad (4.5)$$

To determine $\mathcal{O}_\Lambda$ unambiguously, we must supply either an initial condition at an UV scale Λ_0 , or an asymptotic condition in the case of a renormalizable theory. In the former case, given $\mathcal{O}_{\Lambda_0} = \mathcal{O}_B$, we obtain an integral formula for the solution

of the differential equation (4.3) as follows:

$$\mathcal{O}_\Lambda[\phi] \exp[S_{I,\Lambda}[\phi]] = \int [d\phi'] \mathcal{O}_B[\phi + \phi'] \exp\left[-\frac{1}{2} \int_p \frac{p^2 + m^2}{K_0 - K} \phi' \phi' + S_{I,B}[\phi + \phi']\right]. \quad (4.6)$$

This is obtained by taking an infinitesimal variation of the definition (2.23) of the Wilson action:

$$\exp[S_{I,\Lambda}[\phi]] \equiv \int [d\phi'] \exp\left[-\frac{1}{2} \int \frac{p^2 + m^2}{K_0 - K} \phi' \phi' + S_{I,B}[\phi' + \phi]\right]. \quad (4.7)$$

If the theory is renormalizable, we can choose the Λ_0 dependence of $\mathcal{O}_B$ such that

$$\lim_{\Lambda_0 \rightarrow \infty} \mathcal{O}_\Lambda[\phi] \quad (4.8)$$

exists for any fixed Λ . Alternatively, we can impose an asymptotic condition such as

$$\mathcal{O}_\Lambda[\phi] \xrightarrow{\Lambda \rightarrow \infty} \sum_i \Lambda^{x-x_i} c_i(\ln \Lambda/\mu) \mathcal{O}_i, \quad (4.9)$$

where x is the scale dimension of the composite operator $\mathcal{O}_\Lambda$, and x_i is the scale dimension of $\mathcal{O}_i$, a product of derivatives of ϕ . At a given order in perturbation theory, the coefficient c_i is a polynomial of $\ln \Lambda/\mu$. If $x_i < x$, the Λ dependence of the coefficient of $\mathcal{O}_i$ is completely determined by the differential equation (4.3). For $x_i = x$, though, c_i is left ambiguous by an additive constant independent of $\ln \Lambda/\mu$. Hence, the additive constants $c_i(0)$, satisfying $x_i = x$, parameterize the composite operator. We will give three examples from the ϕ^4 theory in $D = 4$ in subsect. 4.3. It is important to note that if

$$c_i(0) = 0 \quad (4.10)$$

for all i such that $x_i = x$, the operator itself vanishes

$$\mathcal{O}_\Lambda[\phi] = 0, \quad (4.11)$$

due to the linearity of the differential equation (4.3) and uniqueness of its solution.

To understand the correlation functions of composite operators, let us introduce a source J . We recall the relation (2.29) between W_B and W_Λ , which can be written in the form of functional integrals as

$$\begin{aligned} & \int [d\phi] \exp\left[S_B[\phi] + \int J\phi\right] \\ &= \int [d\phi] \exp\left[S_\Lambda[\phi] + \int \frac{K_0}{K} J\phi - \frac{1}{2} \int_p \frac{K_0 - K}{p^2 + m^2} \frac{K_0}{K} J J\right]. \end{aligned} \quad (4.12)$$

Changing the action infinitesimally by $\Delta S_B = \epsilon \mathcal{O}_B$, the Wilson action changes by $\Delta S_\Lambda = \epsilon \mathcal{O}_\Lambda$. Hence, we obtain

$$\begin{aligned} & \int [d\phi] \mathcal{O}_B[\phi] \exp\left[S_B[\phi] + \int J\phi\right] \\ &= \int [d\phi] \mathcal{O}_\Lambda[\phi] \exp\left[S_\Lambda[\phi] + \int \frac{K_0}{K} J\phi - \frac{1}{2} \int_p \frac{K_0 - K}{p^2 + m^2} \frac{K_0}{K} J J\right]. \end{aligned} \quad (4.13)$$

Differentiating this with respect to J 's, we obtain^{*)}

$$\langle \mathcal{O}_B \phi(p_1) \cdots \phi(p_n) \rangle_{S_B} = \prod_{i=1}^n \frac{K(p_i/\Lambda_0)}{K(p_i/\Lambda)} \cdot \langle \mathcal{O}_\Lambda \phi(p_1) \cdots \phi(p_n) \rangle_{S_\Lambda} . \quad (4.14)$$

This is independent of Λ . If the theory is renormalizable, we can take $\Lambda_0 \rightarrow \infty$ to obtain the continuum limit

$$\langle \mathcal{O} \phi(p_1) \cdots \phi(p_n) \rangle^\infty = \prod_{i=1}^n \frac{1}{K(p_i/\Lambda)} \cdot \langle \mathcal{O}_\Lambda \phi(p_1) \cdots \phi(p_n) \rangle_{S_\Lambda} , \quad (4.15)$$

where $\mathcal{O}$ denotes the continuum limit of $\mathcal{O}_B$.

We now wish to show that (4.12) is equivalent to (4.7) as we have promised in §2.2. In other words, the relation (4.12), between the generating functionals W_B and W_Λ , is equivalent to the definition (4.7) of the Wilson action; taking an infinitesimal change of S_B and S_Λ , we obtain (4.13) from (4.12), and (4.6) from (4.7). Since we have already derived (4.12) from (4.7), we only need to derive (4.7) from (4.12). For this purpose we use the functional Fourier inverse:

$$\exp[S_\Lambda[\phi]] = \int [dJ][d\phi'] \exp \left[S_\Lambda[\phi'] + \int_p \frac{K_0}{K} J(-p) (\phi' - \phi)(p) \right] , \quad (4.16)$$

where $J(-p)$ (or its Fourier transform in real space) is integrated along the imaginary axis. Using (4.12) to rewrite the right-hand side, we obtain

$$\exp[S_\Lambda[\phi]] = \int [dJ][d\phi'] \exp \left[S_B[\phi'] + \int J \cdot \left(\phi' - \frac{K_0}{K} \phi \right) + \frac{1}{2} \int \frac{K_0 - K}{p^2 + m^2} \frac{K_0}{K} J J \right] . \quad (4.17)$$

The integral over J is gaussian, and we obtain

$$\begin{aligned} \exp[S_\Lambda[\phi]] &= \int [d\phi'] \exp \left[S_B[\phi'] \right. \\ &\quad \left. - \frac{1}{2} \int (p^2 + m^2) \frac{K}{K_0(K_0 - K)} \left(\phi' - \frac{K_0}{K} \phi \right) \left(\phi' - \frac{K_0}{K} \phi \right) \right] \\ &= \int [d\phi'] \exp \left[S_{I,B}[\phi'] - \int \frac{p^2 + m^2}{K_0 - K} (\phi' - \phi)(\phi' - \phi) - \frac{1}{2} \int \frac{p^2 + m^2}{K} \phi \phi \right] \\ &= \exp \left[-\frac{1}{2} \int \frac{p^2 + m^2}{K} \phi \phi \right] \int [d\phi'] \exp \left[S_{I,B}[\phi' + \phi] - \int \frac{p^2 + m^2}{K_0 - K} \phi' \phi' \right] . \end{aligned} \quad (4.18)$$

This gives (4.7).

To recapitulate, we have introduced three equivalent criteria for a functional $\mathcal{O}_\Lambda[\phi]$ to be a composite operator:

1. $\mathcal{O}_\Lambda$ satisfies the differential equation (4.3).
2. $\mathcal{O}_\Lambda$ satisfies the integral formula (4.6) for some bare functional $\mathcal{O}_B$.
3. $\mathcal{O}_\Lambda$ satisfies the integral formula (4.13) for some bare functional $\mathcal{O}_B$. (4.13) is further equivalent to (4.14).

^{*)} As usual, we only consider connected correlations. In the presence of $J(p)$ for $p \neq 0$, we do not factor out the delta function for momentum conservation from the correlation functions.

4.2. General examples

We introduce some general examples of composite operators that play important roles in realization of symmetry.

4.2.1. $[\phi]_\Lambda(p)$

First of all, note that the elementary field $\phi(p)$ is **not** a composite operator. But

$$\begin{aligned} [\phi]_\Lambda(p) &\equiv \frac{K_0}{K} \phi(p) + \frac{K_0 - K}{p^2 + m^2} \frac{\delta S_\Lambda}{\delta \phi(-p)} \\ &= \phi(p) + \frac{K_0 - K}{p^2 + m^2} \frac{\delta S_{I,\Lambda}}{\delta \phi(-p)} \end{aligned} \quad (4.19)$$

is. We can check this directly by differentiating this with respect to Λ and using the Polchinski equation. Alternatively, we differentiate (4.12) with respect to $J(-p)$ to obtain

$$\begin{aligned} \int [d\phi] \phi(p) \exp \left[S_B[\phi] + \int J\phi \right] &= \int [d\phi] \left(\frac{K_0}{K} \phi(p) - \frac{K_0 - K}{p^2 + m^2} \frac{K_0}{K} J(p) \right) \\ &\times \exp \left[S_\Lambda[\phi] + \int \frac{K_0}{K} J\phi - \frac{1}{2} \int \frac{K_0 - K}{p^2 + m^2} \frac{K_0}{K} J J \right]. \end{aligned} \quad (4.20)$$

Since

$$\begin{aligned} &\frac{K_0}{K} J(p) \int [d\phi] \exp \left[S_\Lambda[\phi] + \int \frac{K_0}{K} J\phi \right] \\ &= \int [d\phi] \exp [S_\Lambda[\phi]] \frac{\delta}{\delta \phi(-p)} \exp \left[\int \frac{K_0}{K} J\phi \right] \\ &= - \int [d\phi] \frac{\delta S_\Lambda[\phi]}{\delta \phi(-p)} \exp \left[S_\Lambda[\phi] + \int \frac{K_0}{K} J\phi \right], \end{aligned} \quad (4.21)$$

we obtain

$$\begin{aligned} &\int [d\phi] \phi(p) \exp \left[S_B[\phi] + \int J\phi \right] \\ &= \int [d\phi] \left(\frac{K_0}{K} \phi(p) + \frac{K_0 - K}{p^2 + m^2} \frac{\delta S_\Lambda}{\delta \phi(-p)} \right) \\ &\times \exp \left[S_\Lambda[\phi] + \int \frac{K_0}{K} J\phi - \frac{1}{2} \int \frac{K_0 - K}{p^2 + m^2} \frac{K_0}{K} J J \right]. \end{aligned} \quad (4.22)$$

Thus, $[\phi]_\Lambda(p)$ is a composite operator according to (4.13). Taking the limit $\Lambda_0 \rightarrow \infty$, we obtain $[\phi]_\Lambda(p)$ that has the asymptotic behavior

$$[\phi]_\Lambda(p) \xrightarrow{\Lambda \rightarrow \infty} \phi(p). \quad (4.23)$$

For any $n = 1, 2, \dots$, we find

$$\prod_{i=1}^n \frac{1}{K(p_i/\Lambda)} \cdot \langle [\phi]_\Lambda(p) \phi(p_1) \cdots \phi(p_n) \rangle_{S_\Lambda} \quad (4.24)$$

is independent of Λ .

4.2.2. $K(p/\Lambda) \frac{\delta S_\Lambda}{\delta \phi(-p)}$

Another example is

$$K(p/\Lambda) \frac{\delta S_\Lambda}{\delta \phi(-p)}. \quad (4.25)$$

To see this is a composite operator, we multiply (4.12) by $J(p)$. An analog of (4.21) gives, for S_B ,

$$J(p) \int [d\phi] \exp \left[S_B[\phi] + \int J\phi \right] = - \int [d\phi] \frac{\delta S_B}{\delta \phi(-p)} \exp \left[S_B[\phi] + \int J\phi \right]. \quad (4.26)$$

Combining this with (4.21), we obtain

$$\begin{aligned} & \int [d\phi] \frac{\delta S_B}{\delta \phi(-p)} \exp \left[S_B[\phi] + \int J\phi \right] \\ &= \frac{K}{K_0} \int [d\phi] \frac{\delta S_\Lambda[\phi]}{\delta \phi(-p)} \exp \left[S_\Lambda[\phi] + \int \frac{K_0}{K} J - \frac{1}{2} \int \frac{K_0 - K}{p^2 + m^2} \frac{K_0}{K} J J \right]. \end{aligned} \quad (4.27)$$

Hence, according to (4.13), (4.25) is a composite operator. This composite operator may be called the equation of motion operator, since

$$\begin{aligned} & \prod_{i=1}^n \frac{1}{K(p_i/\Lambda)} \cdot \left\langle K(p/\Lambda) \frac{\delta S_\Lambda}{\delta \phi(-p)} \phi(p_1) \cdots \phi(p_n) \right\rangle_{S_\Lambda} \\ &= - \sum_{i=1}^n (2\pi)^D \delta^{(D)}(p + p_i) \left\langle \phi(p_1) \cdots \widehat{\phi(p_i)} \cdots \phi(p_n) \right\rangle^\infty, \end{aligned} \quad (4.28)$$

where $\phi(p_i)$ is omitted on the right-hand side.

4.2.3. $\frac{\delta S_{I,\Lambda}}{\delta \phi(-p)}$

We find

$$K(p/\Lambda_0) \frac{\delta S_{I,\Lambda}}{\delta \phi(-p)} = (p^2 + m^2) [\phi]_\Lambda(p) + K(p/\Lambda) \frac{\delta S_\Lambda}{\delta \phi(-p)}. \quad (4.29)$$

Hence,

$$\frac{\delta S_{I,\Lambda}}{\delta \phi(-p)} \quad (4.30)$$

is a composite operator, since it is a sum of two composite operators.

4.2.4. $K(p/\Lambda) \left(\frac{\delta \mathcal{O}_\Lambda}{\delta \phi(-p)} + \mathcal{O}_\Lambda \frac{\delta S_\Lambda}{\delta \phi(-p)} \right)$

We next consider an arbitrary composite operator $\mathcal{O}_\Lambda$ and the corresponding bare operator $\mathcal{O}_B$ so that the two are related by (4.13). Multiplying this by $J(-p)$, and integrating by parts as in (4.21) and (4.26), we obtain

$$\begin{aligned} & \int [d\phi] \left(\frac{\delta \mathcal{O}_B}{\delta \phi(-p)} + \mathcal{O}_B \frac{\delta S_B}{\delta \phi(-p)} \right) \exp \left[S_B[\phi] + \int J\phi \right] \\ &= \frac{K}{K_0} \int [d\phi] \left(\frac{\delta \mathcal{O}_\Lambda}{\delta \phi(-p)} + \mathcal{O}_\Lambda \frac{\delta S_\Lambda}{\delta \phi(-p)} \right) \\ & \quad \times \exp \left[S_\Lambda[\phi] + \int \frac{K_0}{K} J - \frac{1}{2} \int \frac{K_0 - K}{p^2 + m^2} \frac{K_0}{K} J J \right]. \end{aligned} \quad (4.31)$$

Hence, according to (4.13),

$$\mathcal{O}'_\Lambda(p) \equiv K(p/\Lambda) \left(\frac{\delta \mathcal{O}_\Lambda}{\delta \phi(-p)} + \mathcal{O}_\Lambda \frac{\delta S_\Lambda}{\delta \phi(-p)} \right) \quad (4.32)$$

is a composite operator.

The correlation functions of $\mathcal{O}'_\Lambda$ is given in terms of those of $\mathcal{O}_\Lambda$:

$$\begin{aligned} & \langle \mathcal{O}'_\Lambda(p) \phi(p_1) \cdots \phi(p_n) \rangle_{S_\Lambda} \\ &= K(p/\Lambda) \int [d\phi] \phi(p_1) \cdots \phi(p_n) \frac{\delta}{\delta \phi(-p)} (\mathcal{O}_\Lambda[\phi] \exp[S_\Lambda[\phi]]) \\ &= - \sum_{i=1}^n (2\pi)^D \delta^{(D)}(p + p_i) K(p_i/\Lambda) \left\langle \mathcal{O}_\Lambda(p) \phi(p_1) \cdots \widehat{\phi(p_i)} \cdots \phi(p_n) \right\rangle_{S_\Lambda}, \end{aligned} \quad (4.33)$$

where $\phi(p_i)$ is absent on the right-hand side. Divided by $\prod_{i=1}^n K(p_i/\Lambda)$, this becomes independent of Λ .

We may call $\mathcal{O}'_\Lambda$ a generalized equation of motion operator, since $\mathcal{O}'_\Lambda = 1$ gives the equation of motion operator $K(p/\Lambda) \frac{\delta S_\Lambda}{\delta \phi(-p)}$.

4.2.5. $\frac{\delta S_\Lambda[\phi; J]}{\delta J(-p)}$

Suppose that the Wilson action depends on a classical external field J , which is not necessarily coupled linearly to the field ϕ , so that we may not have an explicit formula such as (2.37). We find that the derivative with respect to the external field

$$\frac{\delta S_\Lambda[\phi; J]}{\delta J(-p)} = \frac{\delta S_{I,\Lambda}[\phi; J]}{\delta J(-p)} \quad (4.34)$$

is a composite operator, since the derivative can be regarded as an infinitesimal deformation of the Wilson action. We indeed find that

$$\begin{aligned} & \prod_{i=1}^n \frac{1}{K(p_i/\Lambda)} \cdot \left\langle \frac{\delta S_\Lambda[\phi; J]}{\delta J(-p)} \phi(p_1) \cdots \phi(p_n) \right\rangle_{S_\Lambda} \\ &= \frac{\delta}{\delta J(-p)} \prod_{i=1}^n \frac{1}{K(p_i/\Lambda)} \cdot \langle \phi(p_1) \cdots \phi(p_n) \rangle_{S_\Lambda} \end{aligned} \quad (4.35)$$

is independent of Λ . Hence, substituting $\mathcal{O}_\Lambda = \frac{\delta S_\Lambda[\phi; J]}{\delta J(-p)}$ into (4.32), we obtain

$$K(q/\Lambda) \left\{ \frac{\delta S_\Lambda[\phi; J]}{\delta J(-p)} \frac{\delta S_\Lambda[\phi; J]}{\delta \phi(q)} + \frac{\delta^2 S_\Lambda[\phi; J]}{\delta J(-p) \delta \phi(q)} \right\} \quad (4.36)$$

is a composite operator. The composite operators Σ and $\bar{\Sigma}$ which play important roles for realization of symmetry are given in this form. (See sects. 5 & 7.)

4.2.6. $\frac{\delta \mathcal{O}_\Lambda[\phi; J]}{\delta J(-p)} + \mathcal{O}_\Lambda[\phi; J] \frac{\delta S_{I,\Lambda}[\phi; J]}{\delta J(-p)}$

Let us give one more example from a Wilson action in the presence of J . Given a composite operator satisfying

$$\mathcal{O}_\Lambda[\phi; J] \exp[S_{I,\Lambda}[\phi; J]] = \int [d\phi'] \mathcal{O}_B[\phi + \phi'; J]$$

$$\times \exp \left[-\frac{1}{2} \int_p \frac{p^2 + m^2}{K_0 - K} + S_{I,B}[\phi + \phi'; J] \right], \quad (4.37)$$

we differentiate this with respect to $J(-p)$ to find that

$$\frac{\delta \mathcal{O}_\Lambda[\phi; J]}{\delta J(-p)} + \mathcal{O}_\Lambda[\phi; J] \frac{\delta S_{I,\Lambda}[\phi; J]}{\delta J(-p)} \quad (4.38)$$

is also a composite operator according to (4.6). By choosing $\mathcal{O}_\Lambda = K(q/\Lambda) \frac{\delta S_\Lambda[\phi; J]}{\delta \phi(q)}$ as the equation of motion operator, we obtain (4.36) again.

4.2.7. Summary

To summarize, we have introduced the composite operators of the following types:

composite operator
$[\phi]_\Lambda(p)$ defined by (4.19)
$K(p/\Lambda) \delta S_\Lambda / \delta \phi(-p)$, the equation of motion
$\delta S_{I,\Lambda} / \delta \phi(-p)$, a linear combination of the above two
$K(p/\Lambda) (\delta \mathcal{O}_\Lambda / \delta \phi(-p) + \mathcal{O}_\Lambda \delta S_\Lambda / \delta \phi(-p))$, generalized equation of motion
$\delta S_\Lambda / \delta J(-p)$, for an arbitrary external source J
$\delta \mathcal{O}_\Lambda / \delta J(-p) + \mathcal{O}_\Lambda \delta S_{I,\Lambda} / \delta J(-p)$

Table IV. Various types of composite operators

4.3. Composite operators defined by their asymptotic behaviors

In this subsection, we give three concrete examples specifically from the ϕ^4 theory in $D = 4$. Since the theory is renormalizable, we can introduce composite operators by specifying their asymptotic behaviors.

- (i) $[\phi^2/2]_\Lambda$ is defined by the differential equation (4.3) and the asymptotic condition

$$\left[\frac{\phi^2}{2} \right]_\Lambda(p) \xrightarrow{\Lambda \rightarrow \infty} b(\ln \Lambda / \mu) \frac{1}{2} \int_q \phi(q) \phi(p - q). \quad (4.39)$$

The normalization is determined by the value $b(0)$. For example, we can take

$$b(0) = 1. \quad (4.40)$$

If we choose $b(0) = 0$, then the operator vanishes identically.

- (ii) The dimension 4 operator $[\phi^4/4!]_\Lambda$ with zero momentum has the following asymptotic behavior

$$\begin{aligned} \left[\frac{\phi^4}{4!} \right]_\Lambda(0) &\xrightarrow{\Lambda \rightarrow \infty} d(\ln \Lambda / \mu) \frac{1}{4!} \int_{p_1, p_2, p_3} \phi(p_1) \phi(p_2) \phi(p_3) \phi(-p_1 - p_2 - p_3) \\ &+ \frac{1}{2} \int_p \phi(p) \phi(-p) \{ \Lambda^2 a(\ln \Lambda / \mu) + m^2 b(\ln \Lambda / \mu) + p^2 c(\ln \Lambda / \mu) \}, \end{aligned} \quad (4.41)$$

where $d(0), b(0), c(0)$ parameterize the operator. As a simplest choice, we can take

$$d(0) = 1, \quad b(0) = c(0) = 0. \quad (4.42)$$

(iii) A related composite operator $[\frac{1}{2}\partial_\mu\phi\partial_\mu\phi](0)$ can be defined by

$$c(0) = 1, \quad b(0) = d(0) = 0. \quad (4.43)$$

If we choose $b(0) = c(0) = d(0) = 0$, the operator vanishes.

Any scalar composite operator $\mathcal{O}_\Lambda$ of dimension 4 with zero momentum is given as a linear combination

$$\mathcal{O}_\Lambda = c_1 m^2 \left[\frac{\phi^2}{2} \right]_\Lambda + c_2 \left[\frac{\phi^4}{4!} \right]_\Lambda + c_3 \left[\frac{1}{2} \partial_\mu \phi \partial_\mu \phi \right]_\Lambda, \quad (4.44)$$

where the coefficients $c_{1,2,3}$ are independent of Λ .

4.4. $1PI$ composite operators

We recall from §4.1 that a composite operator $\mathcal{O}_\Lambda[\phi]$ is an infinitesimal change of the interaction action $S_{I,\Lambda}$. Let $\mathcal{O}_\Lambda^{1PI}[\Phi]$ be the corresponding infinitesimal change in the effective average action $\Gamma_{B,\Lambda}[\Phi]$. Using the general relations (2.73) and (2.79), we obtain

$$\mathcal{O}_\Lambda^{1PI}[\Phi] = \mathcal{O}_\Lambda[\phi], \quad (4.45)$$

where Φ is given by (2.85)

$$\Phi(p) = \frac{K_0}{K} \phi(p) + \frac{K_0 - K}{p^2 + m^2} \frac{\delta S_\Lambda}{\delta \phi(-p)}. \quad (4.46)$$

We immediately notice that Φ , which has been introduced for the Legendre transformation, coincides with the composite operator $[\phi]_\Lambda$ defined by (4.19):

$$\Phi(p) = [\phi]_\Lambda(p). \quad (4.47)$$

Since Φ and ϕ are identical asymptotically, we find that in renormalized theories $\mathcal{O}_\Lambda[\phi]$ and $\mathcal{O}_\Lambda^{1PI}[\Phi]$ have the same asymptotic behaviors:

$$\begin{cases} \mathcal{O}_\Lambda[\phi] & \xrightarrow{\Lambda \rightarrow \infty} \sum_i \Lambda^{x-x_i} c_i (\ln \Lambda / \mu) \mathcal{O}_i[\phi], \\ \mathcal{O}_\Lambda^{1PI}[\Phi] & \xrightarrow{\Lambda \rightarrow \infty} \sum_i \Lambda^{x-x_i} c_i (\ln \Lambda / \mu) \mathcal{O}_i[\Phi], \end{cases} \quad (4.48)$$

where ϕ is fixed for $\mathcal{O}_\Lambda$, and Φ for $\mathcal{O}_\Lambda^{1PI}$.

The Λ dependence of $\mathcal{O}_\Lambda^{1PI}$ for fixed Φ can be obtained easily from the Wetterich equation (2.108). Varying the effective average action infinitesimally by $\mathcal{O}_\Lambda^{1PI}[\Phi]$, we obtain

$$\begin{aligned} -\Lambda \frac{\partial}{\partial \Lambda} \mathcal{O}_\Lambda^{1PI}[\Phi] &= -\frac{1}{2} \int_{p,q,r} \left(-\Lambda \frac{\partial}{\partial \Lambda} R_\Lambda(p) \right) \\ &\quad \times \left(\Gamma_{B,\Lambda}^{(2)} \right)^{-1}(p, -q) \frac{\delta^2 \mathcal{O}_\Lambda^{1PI}[\Phi]}{\delta \Phi(-q) \delta \Phi(r)} \left(\Gamma_{B,\Lambda}^{(2)} \right)^{-1}(r, -p), \end{aligned} \quad (4.49)$$

where we have used the following formula for an infinitesimal variation:

$$\delta \left(\Gamma_{B,\Lambda}^{(2)} \right)^{-1}(p, -p) = - \int_{q,r} \left(\Gamma_{B,\Lambda}^{(2)} \right)^{-1}(p, -q) \cdot \delta \Gamma_{B,\Lambda}^{(2)}(q, -r) \cdot \left(\Gamma_{B,\Lambda}^{(2)} \right)^{-1}(r, -p). \quad (4.50)$$

The simplest example satisfying the above differential equation is

$$\mathcal{O}_\Lambda^{1PI}[\Phi] = \Phi(p) , \tag{4.51}$$

for which the right-hand side of (4.49) vanishes. $\Phi(p)$ has no Λ dependence as a functional of Φ ; regarding this as a functional of ϕ , however, this has non-trivial Λ dependence as given by (4.46).

§5. Realization of symmetry by WT identity

5.1. WT identity in the continuum limit

In the continuum limit, the Ward-Takahashi (WT) identity for continuous symmetry is given in the following form:

$$\sum_{i=1}^n \langle \phi(p_1) \cdots \mathcal{O}(p_i) \cdots \phi(p_n) \rangle^\infty = 0, \quad (5.1)$$

where $\mathcal{O}(p)$ is a composite operator, giving an infinitesimal symmetry transformation of the elementary field $\phi(p)$.

We wish to realize the above WT identity in the ERG formalism as an invariance of the Wilson action. Denoting by $\mathcal{O}_\Lambda(p)$ a composite operator that gives $\mathcal{O}(p)$ in the continuum limit, we consider an infinitesimal transformation of the elementary field as

$$\delta\phi(p) = K(p/\Lambda) \mathcal{O}_\Lambda(p). \quad (5.2)$$

As we will explain shortly, the change of the Wilson action under the above transformation is given by the **WT composite operator**:

$$\Sigma_\Lambda \equiv \int_p K(p/\Lambda) \left(\mathcal{O}_\Lambda(p) \frac{\delta S_\Lambda}{\delta\phi(p)} + \frac{\delta\mathcal{O}_\Lambda(p)}{\delta\phi(p)} \right). \quad (5.3)$$

This is a composite operator of type (4.32), integrated over momentum p . Hence, its correlation functions are given by

$$\langle \Sigma_\Lambda \phi(p_1) \cdots \phi(p_n) \rangle^\infty = \sum_{i=1}^n \langle \phi(p_1) \cdots \mathcal{O}_\Lambda(p_i) \cdots \phi(p_n) \rangle^\infty. \quad (5.4)$$

Therefore, the WT identity is equivalent to the vanishing of the WT composite operator Σ_Λ :

$$\Sigma_\Lambda = 0. \quad (5.5)$$

The WT operator (5.3) is easy to understand. The first part of Σ_Λ

$$\int_p K(p/\Lambda) \mathcal{O}_\Lambda(p) \frac{\delta S_\Lambda}{\delta\phi(p)} \quad (5.6)$$

denotes the change of the Wilson action under the infinitesimal transformation (5.2). The second part

$$\int_p K(p/\Lambda) \frac{\delta\mathcal{O}_\Lambda(p)}{\delta\phi(p)} \quad (5.7)$$

denotes the contribution of the transformation (5.2) to the jacobian. Hence, (5.3) is the “quantum” change of the Wilson action, and (5.5) amounts to the **quantum invariance**.

Let us consider Σ_A further. This is determined by S_A and $\mathcal{O}_A$. Recall how S_A and $\mathcal{O}_A$ are determined in renormalizable theories. The interaction part $S_{I,A}$ of the Wilson action is determined by the Polchinski equation

$$-\Lambda \frac{\partial}{\partial \Lambda} S_{I,A} = \int_p \frac{\Delta(p/\Lambda)}{p^2 + m^2} \frac{1}{2} \left\{ \frac{\delta S_{I,A}}{\delta \phi(-p)} \frac{\delta S_{I,A}}{\delta \phi(p)} + \frac{\delta^2 S_{I,A}}{\delta \phi(-p) \delta \phi(p)} \right\}, \quad (5.8)$$

and its asymptotic behavior. For example, for the ϕ^4 theory in $D = 4$, the Wilson action $S_{I,A}$ is specified uniquely by the additive constants $b_2(0), c_2(0), a_4(0)$ in the asymptotic behavior:

$$S_{I,A} \xrightarrow{\Lambda \rightarrow \infty} \int d^4x \left[\left(\Lambda^2 a_2(\ln \Lambda/\mu) + m^2 b_2(\ln \Lambda/\mu) \right) \frac{1}{2} \phi^2 + c_2(\ln \Lambda/\mu) \frac{1}{2} \partial_\mu \phi \partial_\mu \phi + a_4(\ln \Lambda/\mu) \frac{1}{4!} \phi^4 \right]. \quad (5.9)$$

Similarly, the composite operator $\mathcal{O}_A$ is determined by the ERG linear differential equation:

$$-\Lambda \frac{\partial}{\partial \Lambda} \mathcal{O}_A = \int_p \frac{\Delta(p/\Lambda)}{p^2 + m^2} \left\{ \frac{\delta S_{I,A}}{\delta \phi(-p)} \frac{\delta}{\delta \phi(p)} + \frac{1}{2} \frac{\delta^2}{\delta \phi(-p) \delta \phi(p)} \right\} \mathcal{O}_A, \quad (5.10)$$

and its asymptotic behavior:

$$\mathcal{O}_A \xrightarrow{\Lambda \rightarrow \infty} \sum_i \Lambda^{x-x_i} d_i(\ln \Lambda/\mu) \mathcal{O}_i, \quad (5.11)$$

where $x \geq x_i$, and $\mathcal{O}_i$ is a local polynomial of ϕ and its derivatives.^{*)} $\mathcal{O}_A$ is completely determined by $d_i(0)$ for those i satisfying $x_i = x$.

Let $c_i(0)$ ($i = 1, \dots, N$) be the generic parameters of the Wilson action, and $d_i(0)$ ($i = 1, \dots, K$) be those of the composite operator $\mathcal{O}_A$, so that the theory has altogether $N + K$ parameters that we can adjust. We call the theory symmetric if we can adjust these parameters so that

$$\Sigma_A = 0. \quad (5.12)$$

Note that Σ_A itself is a composite operator that satisfies the linear ERG differential equation. So, if we show

$$\lim_{\Lambda \rightarrow \infty} \Sigma_A = 0, \quad (5.13)$$

then it vanishes for any finite Λ .

Let us go over perturbative construction. We expand the parameters in the number of loops:

$$c_i(0) = \sum_{l=0}^{\infty} c_i^{(l)}, \quad d_i(0) = \sum_{l=0}^{\infty} d_i^{(l)}. \quad (5.14)$$

^{*)} $\mathcal{O}_i$ may also contain a product of mass parameters such as m^2 of the ϕ^4 theory.

Let us also expand $S_{I,\Lambda}$ and Σ_Λ in loops:

$$S_{I,\Lambda} = \sum_{l=0}^{\infty} S_{I,\Lambda}^{(l)}, \quad \Sigma_\Lambda = \sum_{l=0}^{\infty} \Sigma_\Lambda^{(l)}. \quad (5.15)$$

Suppose $\Sigma_\Lambda = 0$ is made to vanish up to l -loop

$$\Sigma_\Lambda^{(0)} = \dots = \Sigma_\Lambda^{(l)} = 0 \quad (5.16)$$

by adjusting $c_i(0), d_i(0)$ also up to l -loop. Under this assumption, we consider if we can make $\Sigma_\Lambda^{(l+1)}$ vanish by choosing $c_i^{(l+1)}, d_i^{(l+1)}$ appropriately.

Due to the induction hypothesis, $\Sigma_\Lambda^{(l+1)}$ satisfies the ERG differential equation

$$-\Lambda \frac{\partial}{\partial \Lambda} \Sigma_\Lambda^{(l+1)} = \int_p \frac{\Delta(p/\Lambda)}{p^2 + m^2} \frac{\delta S_{I,\Lambda}^{(0)}}{\delta \phi(-p)} \frac{\delta \Sigma_\Lambda^{(l+1)}}{\delta \phi(p)}. \quad (5.17)$$

Since for fixed momentum p

$$\Delta(p/\Lambda) \xrightarrow{\Lambda \rightarrow \infty} 0, \quad (5.18)$$

we find that the right-hand side vanishes and that $\Sigma_\Lambda^{(l+1)}$ is independent of Λ for large Λ :

$$\Sigma_\Lambda^{(l+1)} \xrightarrow{\Lambda \rightarrow \infty} \sum_i s_i^{(l+1)} \mathcal{O}_i, \quad (5.19)$$

where $s_i^{(l+1)}$ is independent of Λ , and $\mathcal{O}_i$ is a local polynomial of ϕ and its derivatives.*) The scale dimension of $\mathcal{O}_i$ must be the same as that of Σ_Λ .

On the other hand, from the definition (5.3), we obtain

$$\begin{aligned} \Sigma_\Lambda^{(l+1)} &= \int_p K(p/\Lambda) \left(\mathcal{O}_\Lambda^{(l+1)} \frac{\delta S_{I,\Lambda}^{(0)}}{\delta \phi(p)} + \mathcal{O}_\Lambda^{(0)} \frac{\delta S_{I,\Lambda}^{(l+1)}}{\delta \phi(p)} \right) \\ &\quad + \int_p K(p/\Lambda) \left(\sum_{k=1}^l \mathcal{O}_\Lambda^{(l+1-k)} \frac{\delta S_{I,\Lambda}^{(k)}}{\delta \phi(p)} + \frac{\delta \mathcal{O}_\Lambda^{(l)}}{\delta \phi(p)} \right). \end{aligned} \quad (5.20)$$

Only the first integral depends on the $(l+1)$ -loop parameters $c_i^{(l+1)}, d_i^{(l+1)}$. The dependence is linear. Asymptotically, neither $S_{I,\Lambda}^{(0)}$ nor $\mathcal{O}_\Lambda^{(0)}$ has Λ dependence. Hence, the part of $\Sigma_\Lambda^{(l+1)}$ that depends on $c_i^{(l+1)}, d_i^{(l+1)}$ are independent of Λ . For large Λ , the Λ dependence of the first integral cancels that of the second integral, and we find that s_i of (5.19) is written as the sum

$$s_i^{(l+1)} = t_i^{(l+1)} + u_i^{(l+1)}. \quad (5.21)$$

*) $\mathcal{O}_i$ may contain a product of mass parameters such as m^2 of the ϕ^4 theory which has scale dimension 2.

Here, t_i is linear in $c^{(l+1)}$'s and $d^{(l+1)}$'s with l -independent constant coefficients:

$$t_i^{(l+1)} = \sum_{j=1}^N A_{ij} c_j^{(l+1)} + \sum_{j=1}^K B_{ij} d_j^{(l+1)}. \quad (5.22)$$

$u_i^{(l+1)}$ denotes the contribution from the second integral of (5.20), and is determined only by the lower loop coefficients.

Whether or not we can choose $c^{(l+1)}$'s and $d^{(l+1)}$'s so that

$$s_i^{(l+1)} = 0 \quad (5.23)$$

must be examined case by case, and cannot be answered in general. In §6, we discuss four concrete examples. In the cases of QED, the Wess-Zumino model, and the two dimensional $O(N)$ non-linear sigma model, we can prove the possibility of realizing (5.23). In the case of YM theories we can also satisfy (5.23), but the possibility cannot be proven within the present formalism; we must resort to the antifield formalism of §7. In general, if a symmetry present at the tree level cannot be extended to arbitrary loop levels, we call the symmetry **anomalous**. The anomaly is expressed by the non-vanishing Σ_A . We discuss how the axial and chiral anomalies arise in the ERG formalism in §9.

5.2. WT identity for the bare action

In the previous subsection we have discussed the Wilson action in the continuum limit, and shown how to satisfy the Ward-Takahashi identity by adjusting renormalized parameters. The symmetry is checked with the Wilson action, but not with the correlation functions, which correspond to the Wilson action in the limit of $\Lambda \rightarrow 0$. The advantage of our method is twofold:

1. No problem with massless theories — Low momentum modes are crucially important for the manifestation of the underlying symmetry of the theory. But they are totally irrelevant to the question of the existence of the underlying symmetry. By keeping the cutoff Λ non-vanishing, we stay away safely from the low momentum modes. Massless theories cause no problem, since we never look at the correlation functions.
2. Only a few parameters to deal with — We examine the vanishing of the composite operator Σ_A for large Λ . The existence of symmetry is a matter of adjusting a few parameters, if it is possible.

In the following we ask if it is possible to realize symmetry using the bare action S_B with a large cutoff Λ_0 . Under the symmetry transformation

$$\delta\phi(p) = K(p/\Lambda_0) \mathcal{O}_B[\phi](p) \quad (5.24)$$

the bare action changes by

$$\Sigma_B \equiv \int_p K(p/\Lambda_0) \left(\frac{\delta S_B}{\delta\phi(p)} \mathcal{O}_B[\phi](p) + \frac{\delta}{\delta\phi(p)} \mathcal{O}_B[\phi](p) \right). \quad (5.25)$$

If the symmetry exists, the above must vanish. Since we are only interested in the continuum limit, we only demand Σ_B to vanish in the continuum limit $\Lambda_0 \rightarrow \infty$. But what does it mean that Σ_B vanishes in the limit?

Unless it vanishes identically, it is very difficult to see if Σ_B vanishes or not. To see if it vanishes, we must lower the cutoff from Λ_0 to a finite Λ , and take the limit $\Lambda_0 \rightarrow \infty$ while fixing Λ . In other words, we must look at Σ_Λ , defined by

$$\Sigma_\Lambda \exp[S_{I,\Lambda}[\phi]] = \int [d\phi'] \Sigma_B[\phi + \phi'] \exp \left[-\frac{1}{2} \int_p \frac{p^2 + m^2}{K_0 - K} \phi' \phi' + S_{I,B}[\phi + \phi'] \right], \quad (5.26)$$

and check if we can make it vanish in the limit $\Lambda_0 \rightarrow \infty$.

In realizing symmetry in the continuum limit, it is difficult to use the bare action S_B and bare WT operator Σ_B . To appreciate this difficulty, let us consider a bare operator

$$\mathcal{O}_B \equiv \frac{m^2 \phi^4}{\Lambda_0^2 4!} \quad (5.27)$$

in the ϕ^4 theory in $D = 4$. Naively, this operator appears to vanish in the limit $\Lambda_0 \rightarrow \infty$. But that is false. Lowering the cutoff from Λ_0 to Λ , we obtain the following 1-loop correction to $\mathcal{O}_\Lambda$:

$$\mathcal{O}_\Lambda^{(1)} = m^2 c(\ln \Lambda_0/\mu) \frac{\phi^2}{2} + \dots, \quad (5.28)$$

where

$$c(\ln \Lambda_0/\mu) = \frac{1}{\Lambda_0^2} \int_p \frac{K_0 - K}{p^2 + m^2} \simeq \int_p \frac{K(p)}{p^2} = \text{const}. \quad (5.29)$$

This does not vanish even in the limit $\Lambda_0 \rightarrow \infty$. Hence, the small coefficient proportional to $\frac{1}{\Lambda_0^2}$ in $\mathcal{O}_B$ is misleading.

Thus, in general it is difficult to judge the existence of symmetry from the bare action and its transformation property, unless the bare action is exactly invariant, i.e., the symmetry is manifest. We must lower the cutoff from Λ_0 to a finite fixed value Λ . We have already pointed out the potential problem with choosing $\Lambda = 0$. Thus, we advocate the examination of symmetry using a Wilson action at non-vanishing Λ .

§6. Examples

In this section we give a few examples to apply the formalism introduced in the previous section and realize symmetry as the WT identity (5.5).

6.1. QED

The WT identities of QED have been discussed using the ERG formalism in various references such as 27), 92), 107) to name only a few. Our presentation here follows closely that of 107).*)

The classical action of QED is given by

$$S_{cl} \equiv \int d^4x \left[-\frac{1}{2} \partial_\mu A_\nu \partial_\mu A_\nu + \frac{1}{2} (1 - 1/\xi) (\partial_\mu A_\mu)^2 - \partial_\mu \bar{c} \partial_\mu c \right. \\ \left. - \bar{\psi} \left(\frac{1}{i} \not{\partial} + im \right) \psi + e \bar{\psi} \not{A} \psi \right], \quad (6.1)$$

where ξ is a gauge fixing parameter, and $c, \bar{c}$ are the free Faddeev-Popov ghost, antighost fields. The action is invariant under the classical BRST transformation:

$$\begin{cases} \delta_\epsilon A_\mu = \epsilon \frac{1}{i} \partial_\mu c, \\ \delta_\epsilon c = 0, \\ \delta_\epsilon \bar{c} = \epsilon \frac{1}{\xi} \frac{1}{i} \partial_\mu A_\mu, \\ \delta_\epsilon \psi = e \epsilon c \psi, \\ \delta_\epsilon \bar{\psi} = -e \epsilon c \bar{\psi}, \end{cases} \quad (6.2)$$

where ϵ is an arbitrary anticommuting constant. In the following we quantize this system using the ERG formalism.

The Wilson action of QED is given as the sum of free and interacting parts:

$$S_A = S_{F,A} + S_{I,A}, \quad (6.3)$$

where the free action is given by

$$S_{F,A} = -\frac{1}{2} \int_k \frac{1}{K(k/\Lambda)} (k^2 \delta_{\mu\nu} - (1 - 1/\xi) k_\mu k_\nu) A_\mu(-k) A_\nu(k) \\ - \int_k \frac{k^2}{K(k/\Lambda)} \bar{c}(-k) c(k) - \int_p \frac{1}{K(p/\Lambda)} \bar{\psi}(-p) (\not{p} + im) \psi(p). \quad (6.4)$$

Here, we have chosen the same cutoff function for all the fields; it is for the sake of simplicity, and by no means necessary. The cutoff propagators are given by

$$\begin{cases} \langle A_\mu(k) A_\nu(-k) \rangle_{S_{F,A}} = \frac{K(k/\Lambda)}{k^2} \left(\delta_{\mu\nu} - (1 - \xi) \frac{k_\mu k_\nu}{k^2} \right), \\ \langle c(k) \bar{c}(-k) \rangle_{S_{F,A}} = \frac{K(k/\Lambda)}{k^2}, \\ \langle \psi(p) \bar{\psi}(-p) \rangle_{S_{F,A}} = \frac{K(p/\Lambda)}{\not{p} + im}. \end{cases} \quad (6.5)$$

*) For scalar QED, the ERG formalism was first applied in 97).

The ghost and antighost fields are free, and the interaction action depends only on $A_\mu, \psi, \bar{\psi}$. The Polchinski equation is given by

$$\begin{aligned} & -\Lambda \frac{\partial}{\partial \Lambda} S_{I,\Lambda} \\ &= \int_k \frac{\Delta(k/\Lambda)}{k^2} \left(\delta_{\mu\nu} - (1-\xi) \frac{k_\mu k_\nu}{k^2} \right) \cdot \frac{1}{2} \left\{ \frac{\delta S_{I,\Lambda}}{\delta A_\mu(-k)} \frac{\delta S_{I,\Lambda}}{\delta A_\nu(k)} + \frac{\delta^2 S_{I,\Lambda}}{\delta A_\mu(-k) \delta A_\nu(k)} \right\} \\ & - \int_p \frac{\Delta(p/\Lambda)}{p^2 + m^2} \text{Tr} (\not{p} - im) \left[\frac{\vec{\delta}}{\delta \bar{\psi}(-p)} S_{I,\Lambda} \cdot S_{I,\Lambda} \frac{\overleftarrow{\delta}}{\delta \psi(p)} + \frac{\vec{\delta}}{\delta \bar{\psi}(-p)} S_{I,\Lambda} \frac{\overleftarrow{\delta}}{\delta \psi(p)} \right], \end{aligned} \quad (6.6)$$

where the minus sign before the second integral is due to the Fermi statistics. The asymptotic behavior of $S_{I,\Lambda}$ is given in the following form:

$$\begin{aligned} S_{I,\Lambda} \xrightarrow{\Lambda \rightarrow \infty} \int d^4x \left[(\Lambda^2 a_2(\ln \Lambda/\mu) + m^2 b_2(\ln \Lambda/\mu)) \frac{1}{2} A_\mu^2 \right. \\ + c_2(\ln \Lambda/\mu) \frac{1}{2} \partial_\mu A_\nu \partial_\mu A_\nu + d_2(\ln \Lambda/\mu) \frac{1}{2} (\partial_\mu A_\mu)^2 \\ + a_4(\ln \Lambda/\mu) \frac{1}{4!} (A_\mu^2)^2 \\ + a_f(\ln \Lambda/\mu) \bar{\psi} \frac{1}{i} \not{\partial} \psi + b_f(\ln \Lambda/\mu) im \bar{\psi} \psi \\ \left. + a_3(\ln \Lambda/\mu) \bar{\psi} \not{A} \psi \right]. \end{aligned} \quad (6.7)$$

Hence, the theory has 7 parameters:

$$b_2(0), c_2(0), d_2(0), a_f(0), b_f(0), a_3(0), a_4(0). \quad (6.8)$$

Out of these, three are unphysical, since they only change normalization of the fields and the mass parameter m . So, we can arbitrarily impose the normalization condition

$$c_2(0) = a_f(0) = b_f(0) = 0. \quad (6.9)$$

This leaves us four arbitrary parameters:

$$b_2(0), d_2(0), a_3(0), a_4(0). \quad (6.10)$$

We will show how to fix these for the theory to become QED.

The quantum BRST transformation is defined by

$$\begin{cases} \delta A_\mu(k) = k_\mu \epsilon c(k), \\ \delta c(k) = 0, \\ \delta \bar{c}(-k) = \epsilon \frac{1}{\xi} k_\mu [A_\mu]_\Lambda(-k), \\ \delta \psi(p) = e \int_k \epsilon c(k) [\psi]_\Lambda(p-k), \\ \delta \bar{\psi}(-p) = -e \int_k \epsilon c(k) [\bar{\psi}]_\Lambda(-p-k), \end{cases} \quad (6.11)$$

where e is the gauge coupling. We have introduced an anticommuting constant ϵ so that the transformation preserves the statistics of the fields. The composite

operators $[A_\mu]_\Lambda$, $[\psi]_\Lambda$, $[\bar{\psi}]_\Lambda$ are defined by

$$\begin{cases} [A_\mu]_\Lambda(-k) &\equiv \frac{1}{K(k/\Lambda)} A_\mu(-k) + \frac{1-K(k/\Lambda)}{k^2} \left(\delta_{\mu\nu} - (1-\xi) \frac{k_\mu k_\nu}{k^2} \right) \frac{\delta S_\Lambda}{\delta A_\nu(k)}, \\ [\psi]_\Lambda(p) &\equiv \frac{1}{K(p/\Lambda)} \psi(p) + \frac{1-K(p/\Lambda)}{\not{p}+im} \frac{\overrightarrow{\delta}}{\delta \psi(-p)} S_\Lambda, \\ [\bar{\psi}]_\Lambda(-p) &\equiv \frac{1}{K(p/\Lambda)} \bar{\psi}(-p) + S_\Lambda \frac{\overleftarrow{\delta}}{\delta \bar{\psi}(p)} \frac{1-K(p/\Lambda)}{\not{p}+im}. \end{cases} \quad (6.12)$$

The corresponding WT composite operator is

$$\begin{aligned} \Sigma_\Lambda &\equiv \int_k K(k/\Lambda) \left[\frac{\delta S_\Lambda}{\delta A_\mu(k)} \delta A_\mu(k) + \delta \bar{c}(-k) \frac{\overrightarrow{\delta}}{\delta \bar{c}(-k)} S_\Lambda \right] \\ &+ \int_p K(p/\Lambda) \left[S_\Lambda \frac{\overleftarrow{\delta}}{\delta \psi(p)} \delta \psi(p) - \text{Tr} \delta \psi(p) \frac{\overleftarrow{\delta}}{\delta \psi(p)} \right. \\ &\quad \left. + \delta \bar{\psi}(-p) \frac{\overrightarrow{\delta}}{\delta \bar{\psi}(-p)} S_\Lambda - \text{Tr} \frac{\overrightarrow{\delta}}{\delta \bar{\psi}(-p)} \delta \bar{\psi}(-p) \right]. \end{aligned} \quad (6.13)$$

For the Wilson action to be invariant under the BRST transformation (6.11), this must vanish.

Substituting the transformation (6.11) into the above, we obtain^{*)}

$$\Sigma_\Lambda = e \int_k c(k) \left\{ k_\mu \frac{\delta S_{I,\Lambda}}{\delta A_\mu(k)} - \Phi(-k) \right\}, \quad (6.14)$$

where

$$J_\mu(-k) \equiv \frac{\delta S_{I,\Lambda}}{\delta A_\mu(k)} \quad (6.15)$$

is a composite operator of type (4.30), and denotes an electric current. Φ is a composite operator of type (4.32) defined by

$$\begin{aligned} \Phi(-k) &\equiv e \int_p K(p/\Lambda) \text{Tr} \left[[\psi]_\Lambda(p-k) \cdot S_\Lambda \frac{\overleftarrow{\delta}}{\delta \psi(p)} + [\psi]_\Lambda(p-k) \frac{\overleftarrow{\delta}}{\delta \psi(p)} \right] \\ &- e \int_p K(p/\Lambda) \text{Tr} \left[\frac{\overrightarrow{\delta}}{\delta \bar{\psi}(-p)} S_\Lambda \cdot [\bar{\psi}]_\Lambda(-p-k) + \frac{\overrightarrow{\delta}}{\delta \bar{\psi}(-p)} [\bar{\psi}]_\Lambda(-p-k) \right]. \end{aligned} \quad (6.16)$$

Thus, the WT identity is equivalent to the operator identity:

$$k_\mu J_\mu(-k) = \Phi(-k) \quad (6.17)$$

for arbitrary k . This equation is valid if and only if the asymptotic behaviors of the both hand sides match.

^{*)} It is straightforward and not tedious.

The asymptotic behavior of the left-hand side is obtained from (6.7) as

$$\begin{aligned} k_\mu J_\mu(-k) &\xrightarrow{\Lambda \rightarrow \infty} \left(\Lambda^2 a_2(\ln \Lambda/\mu) + m^2 b_2(\ln \Lambda/\mu) + k^2 (c_2 + d_2)(\ln \Lambda/\mu) \right) k_\mu A_\mu(-k) \\ &\quad + a_3(\ln \Lambda/\mu) \int_p \bar{\psi}(-p-k) \not{k} \psi(p) \\ &\quad + a_4(\ln \Lambda/\mu) \frac{1}{2} \int_{k_1, k_2} A_\nu(k_1) A_\nu(k_2) k_\mu A_\mu(-k - k_1 - k_2). \end{aligned} \quad (6.18)$$

To obtain the asymptotic behavior of $\Phi(-k)$, we first note that $\Phi(-k)$ vanishes at $k = 0$ due to the fermion number conservation. For $k = 0$, substituting the definition of the composite operators, we obtain

$$\Phi(0) = e \int_p \left[-S_\Lambda \frac{\overleftarrow{\delta}}{\delta \psi(p)} \psi(p) + \bar{\psi}(-p) \frac{\overrightarrow{\delta}}{\delta \bar{\psi}(-p)} S_\Lambda \right]. \quad (6.19)$$

Since S_Λ contains the same number of ψ 's and $\bar{\psi}$'s, this vanishes:

$$\Phi(0) = 0. \quad (6.20)$$

Therefore, $\Phi(-k)$ is a total derivative. Hence, its asymptotic behavior must be proportional to k . Since Φ has dimension 4, the most general form is given by

$$\begin{aligned} \Phi(-k) &\xrightarrow{\Lambda \rightarrow \infty} \left(\Lambda^2 \bar{a}_2(\ln \Lambda/\mu) + m^2 \bar{b}_2(\ln \Lambda/\mu) + k^2 \bar{d}_2(\ln \Lambda/\mu) \right) k_\mu A_\mu(-k) \\ &\quad + \bar{a}_3(\ln \Lambda/\mu) \int_p \bar{\psi}(-p-k) \not{k} \psi(p) \\ &\quad + \bar{a}_4(\ln \Lambda/\mu) \frac{1}{2} \int_{k_1, k_2} A_\nu(k_1) A_\nu(k_2) k_\mu A_\mu(-k - k_1 - k_2). \end{aligned} \quad (6.21)$$

It is crucial that this has the same form as the asymptotic behavior of $k_\mu J_\mu(-k)$.


Thus, the WT identity is equivalent to the following four equations:

$$\begin{cases} b_2(0) = \bar{b}_2(0), \\ d_2(0) = \bar{d}_2(0), \\ a_3(0) = \bar{a}_3(0), \\ a_4(0) = \bar{a}_4(0), \end{cases} \quad (6.22)$$

where we have used the normalization condition $c_2(0) = 0$. If the right-hand sides do not depend on $b_2(0), d_2(0), a_3(0), a_4(0)$, we can use the above equations to determine these parameters.

To show that the above equations can be solved uniquely, we must examine the asymptotic behavior of $\Phi(-k)$ further. Splitting the action into the free and interaction parts, we can rewrite

$$\begin{aligned} \Phi(-k) &= e \int_p K(p/\Lambda) \left[-S_\Lambda \frac{\overleftarrow{\delta}}{\delta \psi(p)} [\psi]_\Lambda(p-k) + [\bar{\psi}]_\Lambda(-p-k) \frac{\overrightarrow{\delta}}{\delta \bar{\psi}(-p)} S_\Lambda \right] \\ &\quad + e \int_p \text{Tr} U(-p-k, p) \frac{\overrightarrow{\delta}}{\delta \bar{\psi}(-p)} S_{I, \Lambda} \frac{\overleftarrow{\delta}}{\delta \psi(p+k)}, \end{aligned} \quad (6.23)$$

Fig. 19. The broken curve gives eU (6.24).

where

$$U(-p-k, p) \equiv K(p+k/\Lambda) \frac{1 - K(p/\Lambda)}{\not{p} + im} - \frac{1 - K((p+k)/\Lambda)}{\not{p} + \not{k} + im} K(p/\Lambda) . \quad (6.24)$$

It is straightforward to obtain the asymptotic behavior of the first integral on the right-hand side:

$$\begin{aligned} e \int_p K(p/\Lambda) \left[-S_\Lambda \frac{\overleftarrow{\delta}}{\delta\psi(p)} [\psi]_\Lambda(p-k) + [\bar{\psi}]_\Lambda(-p-k) \frac{\overrightarrow{\delta}}{\delta\bar{\psi}(-p)} S_\Lambda \right] \\ \xrightarrow{\Lambda \rightarrow \infty} e(1 - a_f(\ln \Lambda/\mu)) \int_p \bar{\psi}(-p-k) \not{k} \psi(p) . \end{aligned} \quad (6.25)$$

Since $a_f(0) = 0$ by the normalization condition, we find that the above contributes e to $\bar{a}_3(0)$, and nothing at all to $\bar{b}_2(0), \bar{d}_2(0), \bar{a}_4(0)$.

We now introduce the loop expansions of the coefficients:

$$\left\{ \begin{array}{l} b_2(0) = \sum_{l=1}^{\infty} b_2^{(l)} , \\ d_2(0) = \sum_{l=1}^{\infty} d_2^{(l)} , \\ a_3(0) = e + \sum_{l=1}^{\infty} a_3^{(l)} , \\ a_4(0) = \sum_{l=1}^{\infty} a_4^{(l)} , \end{array} \right\} \quad \left\{ \begin{array}{l} \bar{b}_2(0) = \sum_{l=1}^{\infty} \bar{b}_2^{(l)} , \\ \bar{d}_2(0) = \sum_{l=1}^{\infty} \bar{d}_2^{(l)} , \\ \bar{a}_3(0) = e + \sum_{l=1}^{\infty} \bar{a}_3^{(l)} , \\ \bar{a}_4(0) = \sum_{l=1}^{\infty} \bar{a}_4^{(l)} , \end{array} \right. \quad (6.26)$$

where each term of the series is a function of e and ξ . The sum is over 1- and higher-loop contributions; we have fixed the tree level of $a_3(0)$ to satisfy the WT identity (6.22) at tree level.

Now, let us suppose we have fixed $b_2(0), d_2(0), a_3(0), a_4(0)$ up to l -loop level by the WT identity (6.22). This fixes $S_{I,\Lambda}$ up to l -loop level. Using this, we can compute (6.23) up to $(l+1)$ -loop level, where the extra loop is provided by the integral over p for the second integral. Then, using (6.22), we can determine $b_2^{(l+1)}, d_2^{(l+1)}, a_3^{(l+1)}, a_4^{(l+1)}$ unambiguously. We can repeat this procedure to all orders. This is how we can construct QED perturbatively.

We end this subsection by giving concrete 1-loop calculations. We need to evaluate the second integral on the right-hand side of (6.23). We only need the tree level vertex $e\gamma_\mu$ and the definition (6.24) of U for the calculations.

- (i) $b_2^{(1)}, d_2^{(1)}$ — The corresponding Feynman graph is given by Fig. 19.

$$\int_p \text{Tr } e\gamma_\mu eU(-p-k, p)$$




Fig. 20. The solid curve gives a high-momentum photon propagator.

$$\begin{aligned} & \xrightarrow{\Lambda \rightarrow \infty} e^2 k_\mu \left[-2\Lambda^2 \int_p \frac{1}{q^2} \Delta(q) (1 - K(q)) + \left(m^2 + \frac{k^2}{3} \right) \int_p \frac{\Delta(p)}{p^4} \right] \\ & = \left(\Lambda^2 a_2^{(1)} + m^2 b_2^{(1)} + k^2 d_2^{(1)} \right) k_\nu. \end{aligned} \quad (6.27)$$

Using the integral formula (3.14), we obtain

$$b_2^{(1)} = \frac{2e^2}{(4\pi)^2}, \quad d_2^{(1)} = \frac{2e^2}{3(4\pi)^2}. \quad (6.28)$$

We also obtain

$$a_2^{(1)} = -2e^2 \int_p \frac{\Delta(p) (1 - K(p))}{p^2}, \quad (6.29)$$

which is a constant dependent on the choice of K .

- (ii) $a_3^{(1)}$ — The corresponding Feynman graph is given by Fig. 20.

$$\begin{aligned} & (-) \int_q e\gamma_\mu eU(-q-k, q) e\gamma_\nu \\ & \quad \times \frac{1 - K((q-p)/\Lambda)}{(q-p)^2} \left(\delta_{\mu\nu} - (1-\xi) \frac{(q-p)_\mu (q-p)_\nu}{(q-p)^2} \right) \\ & \xrightarrow{\Lambda \rightarrow \infty} -e^3 \not{k} \int_q \frac{1}{q^4} \left\{ \xi K(q) (1 - K(q))^2 + \frac{3-\xi}{4} (1 - K(q)) \Delta(q) \right\} = a_3^{(1)} \not{k}, \end{aligned} \quad (6.30)$$

where the first minus sign is due to the interchange of $\psi(p)$ and $\bar{\psi}(-p-k)$. Hence, we obtain

$$a_3^{(1)} = -e^3 \left(\xi \int_q \frac{K(1-K)^2}{q^4} + \frac{3-\xi}{4(4\pi)^2} \right). \quad (6.31)$$

- (iii) $a_4^{(1)}$ — The corresponding Feynman graph is given by Fig. 21. Writing $k_3 = -k - k_1 - k_2$, we obtain

$$\begin{aligned} & \int_p \text{Tr} e\gamma_\alpha eU(-p-k, p) e\gamma_\gamma \frac{1 - K((q-k_3)/\Lambda)}{\not{q} - \not{k}_3 + im} e\gamma_\beta \frac{1 - K((q-k_2-k_3)/\Lambda)}{\not{q} - \not{k}_2 - \not{k}_3} \\ & \quad + (5 \text{ permutations}) \\ & \xrightarrow{\Lambda \rightarrow \infty} (k_\alpha \delta_{\beta\gamma} + k_\beta \delta_{\gamma\alpha} + k_\gamma \delta_{\alpha\beta}) 2e^4 \int_p \frac{\Delta(q)}{q^4} (1 - K(q))^2 \\ & = (k_\alpha \delta_{\beta\gamma} + k_\beta \delta_{\gamma\alpha} + k_\gamma \delta_{\alpha\beta}) a_4^{(1)}. \end{aligned} \quad (6.32)$$




Fig. 21. The solid line gives a high-momentum electron propagator.

Hence, we obtain

$$a_4^{(1)} = \frac{4}{3} \frac{e^4}{(4\pi)^2}. \quad (6.33)$$

It is interesting to note the presence of a_2, b_2, a_4 . The first and second are photon mass terms, and the third is a four-photon coupling. These break the invariance of the action under the classical BRST transformation (6.2). As has been emphasized in §5, the theory has full gauge (or BRST) invariance, but it requires order-by-order fine tuning, and is not realized manifestly.

6.2. Yang-Mills theories

In the ERG framework YM theories have been discussed usually in the antifield formalism explained in §7. We give relevant references at the beginning of §8.2.

The classical action of a generic YM theory is given by

$$S_{cl} = \int d^4x \left[-\frac{1}{4} (F_{\mu\nu}^a)^2 - \frac{1}{2\xi} (\partial_\mu A_\mu^a)^2 - \partial_\mu \bar{c}^a (D_\mu c)^a \right], \quad (6.34)$$

where the field strength is defined by

$$F_{\mu\nu}^a \equiv \partial_\mu A_\nu^a - \partial_\nu A_\mu^a - g f^{abc} A_\mu^b A_\nu^c, \quad (6.35)$$

and the covariant derivative is defined by

$$(D_\mu c)^a \equiv \partial_\mu c^a - g f^{abc} A_\mu^b c^c. \quad (6.36)$$

f^{abc} is a totally antisymmetric structure constant of whatever gauge group under consideration, satisfying the Jacobi identity

$$f^{abd} f^{dce} + f^{bcd} f^{dae} + f^{cad} f^{dbe} = 0. \quad (6.37)$$

The classical action is invariant under the following classical BRST transformation:

$$\begin{cases} \delta_\epsilon A_\mu^a = \epsilon \frac{1}{i} (D_\mu c)^a, \\ \delta_\epsilon c^a = \epsilon \frac{g}{2i} f^{abc} c^b c^c, \\ \delta_\epsilon \bar{c}^a = \epsilon \frac{1}{\xi} \frac{1}{i} \partial_\mu A_\mu^a, \end{cases} \quad (6.38)$$

where ϵ is an arbitrary anticommuting constant. Transforming twice, we obtain

$$\begin{cases} \delta_\eta \delta_\epsilon A_\mu^a = 0, \\ \delta_\eta \delta_\epsilon c^a = 0, \\ \delta_\eta \delta_\epsilon \bar{c}^a = -\epsilon \eta \frac{1}{\xi} \partial_\mu (D_\mu c)^a. \end{cases} \quad (6.39)$$

The last one vanishes by the ghost equation of motion

$$\partial_\mu (D_\mu c)^a = 0. \quad (6.40)$$

In the following we quantize the system using the ERG formalism.^{*)}

The Wilson action is given as the sum of free and interacting parts:

$$S_A = S_{F,A} + S_{I,A}, \quad (6.41)$$

where the free action is given by

$$\begin{aligned} S_{F,A} = & -\frac{1}{2} \int_k \frac{1}{K(k/\Lambda)} (k^2 \delta_{\mu\nu} - (1 - 1/\xi) k_\mu k_\nu) A_\mu^a(-k) A_\nu^a(k) \\ & - \int_k \frac{k^2}{K(k/\Lambda)} \bar{c}^a(-k) c^a(k). \end{aligned} \quad (6.42)$$

Hence, the cutoff propagators are given by

$$\begin{cases} \langle A_\mu^a(k) A_\nu^b(-k) \rangle_{S_{F,A}} = \delta^{ab} \frac{K(k/\Lambda)}{k^2} \left(\delta_{\mu\nu} - (1 - \xi) \frac{k_\mu k_\nu}{k^2} \right), \\ \langle c^a(k) \bar{c}^b(-k) \rangle_{S_{F,A}} = \delta^{ab} \frac{K(k/\Lambda)}{k^2}. \end{cases} \quad (6.43)$$

The Polchinski equation for the interaction action is given by

$$\begin{aligned} & -\Lambda \frac{\partial}{\partial \Lambda} S_{I,A} \\ & = \int_k \frac{\Delta(k/\Lambda)}{k^2} \left(\delta_{\mu\nu} - (1 - \xi) \frac{k_\mu k_\nu}{k^2} \right) \cdot \frac{1}{2} \left\{ \frac{\delta S_{I,A}}{\delta A_\mu^a(-k)} \frac{\delta S_{I,A}}{\delta A_\nu^a(k)} + \frac{\delta^2 S_{I,A}}{\delta A_\mu^a(-k) \delta A_\nu^a(k)} \right\} \\ & + \int_k \frac{\Delta(k/\Lambda)}{k^2} \left(S_{I,A} \frac{\overleftarrow{\delta}}{\delta c^a(k)} \cdot \frac{\overrightarrow{\delta}}{\delta \bar{c}^a(-k)} S_{I,A} - \frac{\overrightarrow{\delta}}{\delta \bar{c}^a(-k)} S_{I,A} \frac{\overleftarrow{\delta}}{\delta c^a(k)} \right). \end{aligned} \quad (6.44)$$

The possible asymptotic behavior depends on the gauge group. For simplicity we consider only the group SU(2). For SU(2), the structure constant f^{abc} becomes the familiar antisymmetric symbol ϵ^{abc} , where the indices run from 1 to 3. It satisfies

$$\epsilon^{abc} \epsilon^{cde} = \delta^{ad} \delta^{ce} - \delta^{ae} \delta^{cd}. \quad (6.45)$$

^{*)} We can make the classical BRST transformation nilpotent by introducing an auxiliary field B^a , equal to $\frac{1}{\xi} \frac{1}{i} \partial_\mu A_\mu^a$ by the equation of motion. We do not do it here because B^a does not help the perturbative construction described here and in §8.2. In §11, however, we introduce B^a in order to keep the bare action $\tilde{S}_B$ at most linear in the antifields.

The asymptotic behavior of the interaction action is now given in the following form:

$$\begin{aligned}
S_{I,\Lambda} \xrightarrow{\Lambda \rightarrow \infty} \int d^4x \left[\Lambda^2 a_0 (\ln \Lambda/\mu) \frac{1}{2} (A_\mu^a)^2 \right. \\
+ a_1 (\ln \Lambda/\mu) \frac{1}{2} (\partial_\mu A_\nu^a)^2 + a_2 (\ln \Lambda/\mu) \frac{1}{2} (\partial_\mu A_\mu^a)^2 \\
+ a_3 (\ln \Lambda/\mu) g \epsilon^{abc} \partial_\mu A_\nu^a \cdot A_\mu^b A_\nu^c \\
+ a_4 (\ln \Lambda/\mu) \frac{g^2}{4} (A_\mu^a A_\mu^a)^2 + a_5 (\ln \Lambda/\mu) \frac{g^2}{4} (A_\mu^a A_\nu^a)^2 \\
\left. - a_6 (\ln \Lambda/\mu) \partial_\mu \bar{c}^a \partial_\mu c^a - a_7 (\ln \Lambda/\mu) g \epsilon^{abc} \partial_\mu \bar{c}^a A_\mu^b c^c \right]. \quad (6.46)
\end{aligned}$$

Note that the action depends only on the derivative of the antighost field $\bar{c}^a$; this property certainly holds for the tree level action, and it is further inherited by the Wilson action, since the property is preserved by the Polchinski equation. Hence, besides the gauge coupling g , the theory has 7 parameters:

$$a_1(0), a_2(0), a_3(0), a_4(0), a_5(0), a_6(0), a_7(0). \quad (6.47)$$

At tree level, we obtain

$$a_3(0) = 1, a_4(0) = -1, a_5(0) = 1, a_7(0) = -1, \quad (6.48)$$

and the rest vanishing.

Let us now consider the quantum BRST transformation, given by

$$\begin{cases} \delta A_\mu^a(k) &= \epsilon (k_\mu [c^a](k) + \mathcal{A}_\mu^a(k)) , \\ \delta c^a(k) &= \epsilon \mathcal{C}^a(k) , \\ \delta \bar{c}^a(-k) &= \epsilon \frac{1}{\xi} (-k_\mu) [A_\mu^a](-k) , \end{cases} \quad (6.49)$$

where the composite operators $[c^a], [A_\mu^a]$ are defined as usual:

$$\begin{cases} [c^a](k) &= \frac{1}{K(k/\Lambda)} c^a(k) + \frac{1-K(k/\Lambda)}{k^2} \frac{\overrightarrow{\delta}}{\delta \bar{c}^a(-k)} S_\Lambda , \\ [A_\mu^a](-k) &= \frac{1}{K(k/\Lambda)} A_\mu^a(-k) + \frac{1-K(k/\Lambda)}{k^2} \left(\delta_{\mu\nu} - (1-\xi) \frac{k_\mu k_\nu}{k^2} \right) \frac{\delta S_\Lambda}{\delta A_\nu(k)} . \end{cases} \quad (6.50)$$

We define the composite operator $\mathcal{A}_\mu^a(k)$ so that $k_\mu \delta A_\mu^a(k)$ gives the composite operator for the ghost equation of motion:

$$k_\mu (k_\mu [c^a](k) + \mathcal{A}_\mu^a(k)) = -K(k/\Lambda) \frac{\overrightarrow{\delta}}{\delta \bar{c}^a(-k)} S_\Lambda. \quad (6.51)$$

This is equivalent to

$$k_\mu \mathcal{A}_\mu^a(k) = -\frac{\overrightarrow{\delta}}{\delta \bar{c}^a(-k)} S_{I,\Lambda}. \quad (6.52)$$

This implies that the asymptotic behavior of $\mathcal{A}_\mu^a(k)$ is determined by that of the action:

$$\mathcal{A}_\mu^a(k) \xrightarrow{\Lambda \rightarrow \infty} a_6 (\ln \Lambda/\mu) k_\mu c^a(k) + a_7 (\ln \Lambda/\mu) \frac{1}{i} g \epsilon^{abc} \int_l A_\mu^b(k-l) c^c(l). \quad (6.53)$$

On the other hand, the asymptotic behavior of the composite operator $\mathcal{C}^a(k)$ is not determined by the action, and it is given by

$$\mathcal{C}^a(k) \xrightarrow{\Lambda \rightarrow \infty} a_8(\ln \Lambda/\mu) \frac{g}{2i} \epsilon^{abc} \int_l c^b(k-l) c^c(l). \quad (6.54)$$

The parameter $a_8(0)$, which is 1 at tree level, is the normalization constant of $\mathcal{C}^a$. This makes the eighth parameter of the theory, besides the seven parameters (6.47). We wish to fix these eight by demanding the invariance of the Wilson action under the BRST transformation (6.49). Here, we note that the gauge invariance is expected to leave three parameters free to choose: $a_2(0)$, $a_6(0)$ are the normalization constants of the gauge and ghost fields, respectively, and $a_3(0)$ normalizes the coupling constant g . Whatever constraints we find among the eight parameters must leave these three parameters free.

Before going further, let us take a moment to understand the requirement (6.51). The WT identity for the two-point function is

$$\left\langle \delta A_\mu^a(k) \bar{c}^b(-k) \right\rangle^\infty + \left\langle A_\mu^a(k) \delta \bar{c}^b(-k) \right\rangle^\infty = 0, \quad (6.55)$$

which gives

$$\left\langle (k_\mu c^a(k) + \mathcal{A}_\mu^a(k)) \bar{c}^b(-k) \right\rangle^\infty - \frac{1}{\xi} \left\langle A_\mu^a(k) k_\nu A_\nu^b(-k) \right\rangle^\infty = 0. \quad (6.56)$$

Thus, (6.51) gives

$$\frac{1}{\xi} \left\langle k_\mu A_\mu^a(k) k_\nu A_\nu^b(-k) \right\rangle^\infty = \delta^{ab}, \quad (6.57)$$

implying that the longitudinal part of the gauge boson propagator does not receive any radiative correction.

The WT composite operator⁶⁶⁾ is now defined by

$$\begin{aligned} \Sigma_\Lambda \equiv \int_k K(k/\Lambda) & \left[\frac{\delta S_\Lambda}{\delta A_\mu^a(k)} \delta A_\mu^a(k) + \frac{\delta}{\delta A_\mu^a(k)} \delta A_\mu^a(k) \right. \\ & + S_\Lambda \frac{\overleftarrow{\delta}}{\delta c^a(k)} \delta c^a(k) - \delta c^a(k) \frac{\overleftarrow{\delta}}{\delta c^a(k)} \\ & \left. + \delta \bar{c}^a(-k) \frac{\overrightarrow{\delta}}{\delta \bar{c}^a(-k)} S_\Lambda - \frac{\overrightarrow{\delta}}{\delta \bar{c}^a(-k)} \delta \bar{c}^a(-k) \right]. \end{aligned} \quad (6.58)$$

This is a dimension 5 scalar composite operator with ghost number 1. Hence, for SU(2), its asymptotic behavior must have the following form:

$$\begin{aligned} \Sigma_\Lambda \xrightarrow{\Lambda \rightarrow \infty} \epsilon \int d^4x & \left[\Lambda^2 s_0 c^a \partial_\mu A_\mu^a + s_1 \frac{1}{i} c^a \partial^2 \partial_\mu A_\mu^a \right. \\ & + s_2 \frac{1}{i} g \epsilon^{abc} c^a \partial^2 A_\mu^b \cdot A_\mu^c + s_3 \frac{1}{i} g \epsilon^{abc} c^a \partial_\mu \partial_\nu A_\nu^b \cdot A_\mu^c \\ & \left. + s_4 i g^2 c^a \partial_\mu A_\mu^a \cdot A_\nu^b A_\nu^b + s_5 i g^2 c^a \partial_\mu A_\nu^a \cdot A_\mu^b A_\nu^b + s_6 i g^2 c^a A_\mu^a \partial_\nu A_\nu^b A_\mu^b \right] \end{aligned}$$

$$\begin{aligned}
& +s_7 ig^2 c^a A_\mu^a \partial_\nu A_\mu^b A_\nu^b + s_8 ig^2 c^a A_\mu^a \partial_\mu A_\nu^b A_\nu^b \\
& +s_9 g\epsilon^{abc} \partial_\mu c^a c^b \frac{1}{i} \partial_\mu \bar{c}^c + s_{10} g^2 c^a c^b \frac{1}{i} \partial_\mu \bar{c}^a A_\mu^a \Big] , \tag{6.59}
\end{aligned}$$

where the coefficients s_i ($i = 0, 1, \dots, 10$) depend on $\ln \frac{\Lambda}{\mu}$ and the gauge coupling constant g . As a composite operator, Σ_Λ has ten parameters:

$$s_i(\ln \Lambda/\mu = 0) = s_i(0) \quad (i = 1, \dots, 10). \tag{6.60}$$

We must make these vanish by arranging the eight parameters.

Substituting the BRST transformation (6.49) into (6.58), we can simplify the expression of Σ_Λ somewhat:

$$\begin{aligned}
\Sigma_\Lambda = & \epsilon \int_k K(k/\Lambda) \left(\mathcal{A}_\mu^a(k) \frac{\delta S_\Lambda}{\delta A_\mu^a(k)} + \frac{\delta}{\delta A_\mu^a(k)} \mathcal{A}_\mu^a(k) \right) \\
& - \epsilon \int_k K(k/\Lambda) \left(S_\Lambda \frac{\overleftarrow{\delta}}{\delta c^a(k)} c^a(k) - c^a(k) \frac{\overleftarrow{\delta}}{\delta c^a(k)} S_\Lambda \right) \\
& + \epsilon \int_k \left[k_\mu \frac{\delta S_{I,\Lambda}}{\delta A_\mu^a(k)} c^a(k) - \frac{1}{\xi} k_\mu A_\mu^a(-k) \frac{\overrightarrow{\delta}}{\delta \bar{c}^a(-k)} S_{I,\Lambda} \right]. \tag{6.61}
\end{aligned}$$

Each of the three integrals is a composite operator by itself.

Let us now introduce loop expansions of S_Λ and Σ_Λ :

$$S_\Lambda = \sum_{l=0}^{\infty} S_\Lambda^{(l)}, \quad \Sigma_\Lambda = \sum_{l=0}^{\infty} \Sigma_\Lambda^{(l)}. \tag{6.62}$$

Correspondingly, let us expand the parameters:

$$a_i(0) = \sum_{l=0}^{\infty} a_i^{(l)}. \quad (i = 1, \dots, 8) \tag{6.63}$$

Suppose we have made

$$\Sigma_\Lambda^{(0)} = \dots = \Sigma_\Lambda^{(l)} = 0 \tag{6.64}$$

by arranging $a_i^{(0)}, \dots, a_i^{(l)}$. Then, as s_i 's at $(l+1)$ -loop are all constants, independent of $\ln \Lambda/\mu$, as a consequence of the ERG differential equation for Σ_Λ . We ask if we can choose the $(l+1)$ -loop coefficients $a_i^{(l+1)}$ to make all s_i vanish at $(l+1)$ -loop, which is equivalent to

$$\Sigma_\Lambda^{(l+1)} = 0. \tag{6.65}$$

Examining the structure of Σ_Λ above, we find that at $(l+1)$ -loop the coefficients s_i are given as the sum:

$$s_i = t_i + u_i \quad (i = 1, \dots, 10), \tag{6.66}$$

where t 's are linear in $a^{(l+1)}$'s, and u 's are determined by the action only up to l -loop. Not all t 's are independent; only the following five are independent:

$$t_1 = a_1^{(l+1)} + a_2^{(l+1)}, \quad (6.67a)$$

$$t_2 = a_1^{(l+1)} + a_3^{(l+1)} + a_6^{(l+1)} + a_7^{(l+1)}, \quad (6.67b)$$

$$t_4 = a_3^{(l+1)} + a_4^{(l+1)} - a_6^{(l+1)} - a_7^{(l+1)}, \quad (6.67c)$$

$$t_5 = -a_3^{(l+1)} + a_5^{(l+1)} + a_6^{(l+1)} + a_7^{(l+1)}, \quad (6.67d)$$

$$t_9 = a_7^{(l+1)} + a_8^{(l+1)}. \quad (6.67e)$$

The remaining t 's are given by

$$t_3 = t_1 - t_2, \quad (6.68a)$$

$$t_6 = t_5, \quad (6.68b)$$

$$t_7 = 2t_4, \quad (6.68c)$$

$$t_8 = t_5, \quad (6.68d)$$

$$t_{10} = -t_9. \quad (6.68e)$$

It is possible to make all $s_i = 0$ by adjusting $a^{(l+1)}$'s if and only if u 's satisfy the same relations as t 's. If t 's and u 's satisfy the above relations, so do s 's:

$$s_3 = s_1 - s_2, \quad (6.69a)$$

$$s_6 = s_5, \quad (6.69b)$$

$$s_7 = 2s_4, \quad (6.69c)$$

$$s_8 = s_5, \quad (6.69d)$$

$$s_{10} = -s_9. \quad (6.69e)$$

If these relations hold, we have five linearly independent conditions on eight parameters, leaving three of them arbitrary. This is as expected, since the YM gauge symmetry does not constrain the normalization of fields and the coupling constant g . For example, we can adopt the normalization condition:

$$a_1(0) = a_6(0) = 0, \quad a_3(0) = 1. \quad (6.70)$$

It is impossible to derive the five relations (6.69) within the formalism of the WT identity. We will resort to the antifield formalism of §7 for derivation.

6.3. Wess-Zumino model

The ERG formalism has been applied to the Wess-Zumino model, for example, in 35), 93), and 102), where the supersymmetry transformation is linearized via auxiliary fields. Without the auxiliary fields, the supersymmetry transformation is non-linear, and the model was first formulated with ERG in 110).

The Wess-Zumino model¹²⁰⁾ is the simplest supersymmetric model defined in 4 dimensions. Without auxiliary fields, the classical action of the model is given as

$$S_{cl} \equiv - \int d^4x \left[\partial_\mu \bar{\phi} \partial_\mu \phi + \bar{\chi}_L \sigma_\mu \partial_\mu \chi_R \right]$$

$$\begin{aligned}
& + (m + g\phi) \frac{1}{2} \bar{\chi}_R \chi_R + (\bar{m} + \bar{g}\bar{\phi}) \frac{1}{2} \bar{\chi}_L \chi_L \\
& + \left(m\phi + g\frac{\phi^2}{2} \right) \left(\bar{m}\bar{\phi} + \bar{g}\frac{\bar{\phi}^2}{2} \right) \Big], \tag{6.71}
\end{aligned}$$

where $\chi_{R,L}$ are 2-component spinors, and we define

$$\sigma_\mu \equiv (\vec{\sigma}, -i) \quad \bar{\sigma}_\mu \equiv (\vec{\sigma}, i), \quad \bar{\chi}_R \equiv \chi_R^T \sigma_y, \quad \bar{\chi}_L \equiv \chi_L^T \sigma_y. \tag{6.72}$$

This is invariant under the following supersymmetry transformation:

$$\left\{ \begin{array}{l} \delta_{cl} \phi \equiv \bar{\xi}_R \chi_R, \\ \delta_{cl} \bar{\phi} \equiv \bar{\xi}_L \chi_L, \\ \delta_{cl} \chi_R \equiv \bar{\sigma}_\mu \xi_L \partial_\mu \phi - \left(\bar{m}\bar{\phi} + \bar{g}\frac{\bar{\phi}^2}{2} \right) \xi_R, \\ \delta_{cl} \chi_L \equiv \sigma_\mu \xi_R \partial_\mu \bar{\phi} - \left(m\phi + g\frac{\phi^2}{2} \right) \xi_L, \end{array} \right. \tag{6.73}$$

where $\xi_{R,L}$ are anticommuting constant spinors.

The above classical action has a spontaneously broken $\mathbf{Z}_2$ invariance. To make the symmetry manifest, we shift the scalar fields as follows:

$$\left\{ \begin{array}{l} \phi + \frac{m}{g} \longrightarrow \phi, \\ \bar{\phi} + \frac{\bar{m}}{\bar{g}} \longrightarrow \bar{\phi}. \end{array} \right. \tag{6.74}$$

Denoting

$$v \equiv \frac{m}{g} \quad \bar{v} \equiv \frac{\bar{m}}{\bar{g}}, \tag{6.75}$$

we can rewrite the action with manifest $\mathbf{Z}_2$ invariance:

$$\begin{aligned}
S_{cl} = - \int & \left[\partial_\mu \bar{\phi} \partial_\mu \phi + \bar{\chi}_L \sigma_\mu \partial_\mu \chi_R + \frac{g}{2} \phi \bar{\chi}_R \chi_R + \frac{\bar{g}}{2} \bar{\phi} \bar{\chi}_L \chi_L \right. \\
& \left. + \frac{g}{2} (\phi^2 - v^2) \frac{\bar{g}}{2} (\bar{\phi}^2 - \bar{v}^2) \right] \tag{6.76}
\end{aligned}$$

Under the $\mathbf{Z}_2$ transformation, the fields transform as follows:

$$\begin{aligned}
\phi & \longrightarrow -\phi, & \bar{\phi} & \longrightarrow -\bar{\phi} \\
\chi_R & \longrightarrow i\chi_R, & \chi_L & \longrightarrow \frac{1}{i}\chi_L
\end{aligned} \tag{6.77}$$

In terms of the shifted scalar fields, the supersymmetry transformation is given by

$$\left\{ \begin{array}{l} \delta_{cl} \phi \equiv \bar{\xi}_R \chi_R, \\ \delta_{cl} \bar{\phi} \equiv \bar{\xi}_L \chi_L, \\ \delta_{cl} \chi_R \equiv \bar{\sigma}_\mu \xi_L \partial_\mu \phi - \frac{\bar{g}}{2} (\bar{\phi}^2 - \bar{v}^2) \xi_R, \\ \delta_{cl} \chi_L \equiv \sigma_\mu \xi_R \partial_\mu \bar{\phi} - \frac{g}{2} (\phi^2 - v^2) \xi_L. \end{array} \right. \tag{6.78}$$

The classical action is parameterized by two complex parameters g and v . If we ask if their phases have any physical significance, we immediately see that their phase changes can be canceled by the corresponding phase changes of the fields:

1. $g \rightarrow ge^{i\alpha}$, $\bar{g} \rightarrow \bar{g}e^{-i\alpha}$ are canceled by

$$\phi \rightarrow e^{-i\alpha}\phi, \quad \bar{\phi} \rightarrow e^{i\alpha}\bar{\phi}. \quad (6.79)$$

2. $v \rightarrow ve^{i\beta}$, $\bar{v} \rightarrow \bar{v}e^{-i\beta}$ are canceled by

$$\phi \rightarrow e^{i\beta}\phi, \quad \bar{\phi} \rightarrow e^{-i\beta}\bar{\phi}, \quad \chi_R \rightarrow e^{-i\frac{\beta}{2}}\chi_R, \quad \chi_L \rightarrow e^{i\frac{\beta}{2}}\chi_L. \quad (6.80)$$

Hence, the physics of the model depends only on $|g|$ and $|v|$, and we make sure that this property is carried over upon quantization.^{*)}

We now consider the Wilson action:

$$S_A = S_{F,A} + S_{I,A}, \quad (6.81)$$

where the free action is

$$S_{F,A} \equiv - \int_p \left[\frac{1}{K_b(p/\Lambda)} p^2 \bar{\phi}(-p) \phi(p) + \frac{1}{K_f(p/\Lambda)} \bar{\chi}_L(-p) i p \cdot \sigma \chi_R(p) \right]. \quad (6.82)$$

For generality we take the cutoff functions of the bosons and fermions independent, since their equality is not demanded by supersymmetry.

The interaction action is determined by the ERG differential equation

$$\begin{aligned} -\Lambda \frac{\partial}{\partial \Lambda} S_{I,A} &= \int_p \frac{\Delta_b(p/\Lambda)}{p^2} \left\{ \frac{\delta S_I}{\delta \bar{\phi}(-p)} \frac{\delta S_I}{\delta \phi(p)} + \frac{\delta^2 S_I}{\delta \bar{\phi}(-p) \delta \phi(p)} \right\} \\ &- \int_p \frac{\Delta_f(p/\Lambda)}{p^2} \text{Tr}(-ip) \cdot \bar{\sigma} \left\{ \frac{\vec{\delta}}{\delta \bar{\chi}_L(-p)} S_I \cdot S_I \frac{\overleftarrow{\delta}}{\delta \chi_R(p)} + \frac{\vec{\delta}}{\delta \bar{\chi}_L(-p)} S_I \frac{\overleftarrow{\delta}}{\delta \chi_R(p)} \right\}, \end{aligned} \quad (6.83)$$

and the asymptotic behavior

$$\begin{aligned} S_{I,A} \xrightarrow{\Lambda \rightarrow \infty} \int d^4x &\left[\Lambda^2 a(\ln \Lambda/\mu) |\phi|^2 + c_1(\ln \Lambda/\mu) \partial_\mu \bar{\phi} \partial_\mu \phi + c_2(\ln \Lambda/\mu) \bar{\chi}_L \sigma \cdot \partial \chi_R \right. \\ &+ c_3(\ln \Lambda/\mu) \left(g \phi \frac{1}{2} \bar{\chi}_R \chi_R + \bar{g} \bar{\phi} \frac{1}{2} \bar{\chi}_L \chi_L \right) + c_4(\ln \Lambda/\mu) |g|^2 \frac{1}{4} |\phi|^4 \\ &\left. + c_5(\ln \Lambda/\mu) |g|^2 \frac{1}{4} (\bar{v}^2 \phi^2 + v^2 \bar{\phi}^2) + c_6(\ln \Lambda/\mu) \frac{1}{4} |g|^2 |v|^4 \right]. \end{aligned} \quad (6.84)$$

Hence, the action has six parameters:

$$c_i(0) \quad (i = 1, \dots, 6). \quad (6.85)$$

We have included $c_6(0)$ as a parameter, even though it corresponds to an additive constant to the action. With the constant term, the derivative of the action with respect to v^2 gives an integral of the composite operator

$$\left[\frac{\phi^2}{2} \right]_\Lambda(x) \quad (6.86)$$

^{*)} The following results in this section are taken from a recent unpublished work of H. Sonoda and K. Ülker. The use of the $\mathbf{Z}_2$ symmetry makes the construction much simpler than that given in the earlier work of the same authors.¹¹⁰⁾

over space:

$$\frac{2}{\bar{g}} \frac{\partial}{\partial \bar{v}^2} S_{I,A} = \int d^4x g \left[\frac{\phi^2}{2} \right]_A (x). \quad (6.87)$$

Similarly, the derivative with respect to v^2 gives

$$\frac{2}{g} \frac{\partial}{\partial v^2} S_{I,A} = \int d^4x \bar{g} \left[\frac{\bar{\phi}^2}{2} \right]_A (x). \quad (6.88)$$

The asymptotic behavior (6.84) implies the following asymptotic behavior of the two composite operators in coordinate space:

$$\left\{ \begin{array}{l} \left[\frac{\phi^2}{2} \right]_A \xrightarrow{\Lambda \rightarrow \infty} c_5(\ln \Lambda/\mu) \frac{\phi^2}{2} + c_6(\ln \Lambda/\mu) \frac{v^2}{2}, \\ \left[\frac{\bar{\phi}^2}{2} \right]_A \xrightarrow{\Lambda \rightarrow \infty} c_5(\ln \Lambda/\mu) \frac{\bar{\phi}^2}{2} + c_6(\ln \Lambda/\mu) \frac{\bar{v}^2}{2}. \end{array} \right. \quad (6.89)$$

We now define the WT identity for supersymmetry by

$$\begin{aligned} \Sigma_A \equiv & \int_p K_b(p/\Lambda) \left[\delta\phi(p) \frac{\delta}{\delta\phi(p)} S_A + \frac{\delta}{\delta\phi(p)} \delta\phi(p) + \delta\bar{\phi}(p) \frac{\delta}{\delta\bar{\phi}(p)} S_A + \frac{\delta}{\delta\bar{\phi}(p)} \delta\bar{\phi}(p) \right] \\ & + \int_p K_f(p/\Lambda) \left[S_A \frac{\overleftarrow{\delta}}{\delta\chi_R(p)} \delta\chi_R(p) - \text{Tr} \delta\chi_R(p) \frac{\overleftarrow{\delta}}{\delta\chi_R(p)} \right. \\ & \left. + S_A \frac{\overleftarrow{\delta}}{\delta\chi_L(p)} \delta\chi_L(p) - \text{Tr} \delta\chi_L(p) \frac{\overleftarrow{\delta}}{\delta\chi_L(p)} \right], \end{aligned}$$

where the composite operators $\delta\phi(p)$, etc., are defined by

$$\left\{ \begin{array}{l} \delta\phi(p) = \bar{\xi}_R[\chi_R](p), \\ \delta\chi_R(p) = [\phi](p) i p \cdot \bar{\sigma} \xi_L - \bar{g} \left[\frac{\bar{\phi}^2}{2} \right](p) \xi_R, \\ \delta\bar{\phi}(p) = \bar{\xi}_L[\chi_L](p), \\ \delta\chi_L(p) = [\bar{\phi}](p) i p \cdot \sigma \xi_R - g \left[\frac{\phi^2}{2} \right](p) \xi_L. \end{array} \right. \quad (6.90)$$

Under the phase changes of g, v , the composite operator Σ_A remains invariant, if we change the phases of the fields according to (6.79) & (6.80), and at the same time change the phases of $\xi_{R,L}$ according to

$$\xi_R \rightarrow e^{-i\alpha + i\frac{3}{2}\beta} \xi_R, \quad \xi_L \rightarrow e^{i\alpha - i\frac{3}{2}\beta} \xi_L. \quad (6.91)$$

In the remainder of this subsection, we wish to show that we can satisfy the WT identity

$$\Sigma_A = 0 \quad (6.92)$$

by tuning four of the six parameters, leaving $c_1(0)$ and $c_3(0)$ arbitrary. The physical meaning of the arbitrary parameters is clear:

1. $c_1(0)$ — overall normalization of the fields; the relative normalization of ϕ and χ_R is fixed by supersymmetry.

2. $c_3(0)$ — normalization of g .

Note that the relative normalization of ϕ and $\bar{g}[\bar{\phi}^2]$ is fixed by supersymmetry. Hence, the relative normalization of ϕ and gv^2 is also fixed, since $\bar{g}[\bar{\phi}^2]$ is the derivative of the action with respect to gv^2 .

We first enumerate all possible terms in the asymptotic behavior of the composite operator Σ_A :

$$\begin{aligned} \Sigma_A \xrightarrow{\Lambda \rightarrow \infty} \int d^4x \big[& \Lambda^2 b(\ln \Lambda/\mu) (\bar{\chi}_R \xi_R \bar{\phi} + \bar{\chi}_L \xi_L \phi) \\ & + s_1(\ln \Lambda/\mu) (\partial^2 \bar{\phi} \bar{\chi}_R \xi_R + \partial^2 \phi \bar{\chi}_L \xi_L) \\ & + s_2(\ln \Lambda/\mu) \left(\bar{g} \bar{\chi}_L \sigma_\mu \xi_R \partial_\mu \frac{\bar{\phi}^2}{2} + g \bar{\chi}_R \bar{\sigma}_\mu \xi_L \partial_\mu \frac{\phi^2}{2} \right) \\ & + s_3(\ln \Lambda/\mu) |g|^2 \left(\bar{\chi}_R \xi_R \phi \frac{\bar{\phi}^2}{2} + \bar{\chi}_L \xi_L \bar{\phi} \frac{\phi^2}{2} \right) \\ & + s_4(\ln \Lambda/\mu) |g|^2 \left(\frac{\bar{v}^2}{2} \phi \bar{\chi}_R \xi_R + \frac{v^2}{2} \bar{\phi} \bar{\chi}_L \xi_L \right) \big]. \end{aligned}$$

We have enumerated all terms of dimension up to 4, which are invariant under the simultaneous phase changes of the parameters and fields given by (6.79), (6.80), (6.91).

At tree level, we find that the choice

$$a = c_1 = c_2 = 0, \quad c_3 = c_4 = -1, \quad c_5 = 1, \quad c_6 = -1 \quad (6.93)$$

satisfies the tree level WT identity

$$\Sigma_A^{(0)} = 0. \quad (6.94)$$

We now assume the vanishing of Σ_A up to l -loop level:

$$\Sigma_A^{(1)} = \dots = \Sigma_A^{(l)} = 0. \quad (6.95)$$

Then, the ERG differential equation for $\Sigma_A^{(l+1)}$ becomes

$$\begin{aligned} -\Lambda \frac{\partial}{\partial \Lambda} \Sigma_A^{(l+1)} = & \int_p \frac{\Delta_b(p/\Lambda)}{p^2} \left(\frac{\delta S_{I,\Lambda}^{(0)}}{\delta \phi(p)} \frac{\delta \Sigma_A^{(l+1)}}{\delta \bar{\phi}(-p)} + \frac{\delta S_{I,\Lambda}^{(0)}}{\delta \bar{\phi}(p)} \frac{\delta \Sigma_A^{(l+1)}}{\delta \phi(-p)} \right) \\ & + \int_p \frac{\Delta_f(p/\Lambda)}{p^2} \left(S_{I,\Lambda}^{(0)} \frac{\overleftarrow{\delta}}{\delta \chi_R(p)}(-ip) \cdot \bar{\sigma} \frac{\overrightarrow{\delta}}{\delta \bar{\chi}_L(-p)} \Sigma_A^{(l+1)} \right. \\ & \left. + S_{I,\Lambda}^{(0)} \frac{\overleftarrow{\delta}}{\delta \chi_L(p)}(-ip) \cdot \sigma \frac{\overrightarrow{\delta}}{\delta \bar{\chi}_R(-p)} \Sigma_A^{(l+1)} \right). \end{aligned} \quad (6.96)$$

This vanishes as $\Lambda \rightarrow \infty$. Hence, we obtain that

$$b^{(l+1)}(\ln \Lambda/\mu) = 0, \quad (6.97)$$

and that $s_i^{(l+1)}$'s are independent of Λ . In the following we wish to show that we can choose $c_i^{(l+1)}(0)$ ($i = 1, \dots, 6$) so that

$$s_i^{(l+1)} = 0 \quad (i = 1, \dots, 4). \quad (6.98)$$

We split $s^{(l+1)}$'s into $t^{(l+1)}$'s linear in $c^{(l+1)}$'s and $u^{(l+1)}$'s independent of them:

$$s_i^{(l+1)} = t_i^{(l+1)} + u_i^{(l+1)}. \quad (6.99)$$

The coefficients $t^{(l+1)}$'s can be calculated from

$$\begin{aligned} \Sigma_A^{(l+1),t} \equiv & \int_p K_b(p/\Lambda) \left(\delta\phi^{(0)}(p) \frac{\delta S_A^{(l+1)}}{\delta\phi(p)} + \delta\bar{\phi}^{(0)}(p) \frac{\delta S_A^{(l+1)}}{\delta\bar{\phi}(p)} \right) \\ & + \int_p K_f(p/\Lambda) \left(S_A^{(l+1)} \frac{\overleftarrow{\delta}}{\delta\chi_R(p)} \delta\chi_R^{(0)}(p) + S_A^{(l+1)} \frac{\overleftarrow{\delta}}{\delta\chi_L(p)} \delta\chi_L^{(0)}(p) \right. \\ & \left. + S_A^{(0)} \frac{\overleftarrow{\delta}}{\delta\chi_R(p)} \delta\chi_R^{(l+1)}(p) + S_A^{(0)} \frac{\overleftarrow{\delta}}{\delta\chi_L(p)} \delta\chi_L^{(l+1)}(p) \right). \end{aligned} \quad (6.100)$$

We then obtain, by simple substitution of the asymptotic behavior of $S_A^{(l+1)}$ & $\delta\chi_{R,L}^{(l+1)}$ into the above, the following results:

$$\begin{cases} t_1^{(l+1)} = -c_1^{(l+1)} + c_2^{(l+1)}, \\ t_2^{(l+1)} = c_5^{(l+1)} - c_2^{(l+1)} + c_3^{(l+1)}, \\ t_3^{(l+1)} = c_5^{(l+1)} - c_3^{(l+1)} + c_4^{(l+1)}, \\ t_4^{(l+1)} = c_6^{(l+1)} + c_3^{(l+1)} + c_5^{(l+1)}. \end{cases} \quad (6.101)$$

Thus, we must choose $c^{(l+1)}$'s so that

$$\begin{cases} -c_1^{(l+1)} + c_2^{(l+1)} = -u_1^{(l+1)}, \\ c_5^{(l+1)} - c_2^{(l+1)} + c_3^{(l+1)} = -u_2^{(l+1)}, \\ c_5^{(l+1)} - c_3^{(l+1)} + c_4^{(l+1)} = -u_3^{(l+1)}, \\ c_6^{(l+1)} + c_3^{(l+1)} + c_5^{(l+1)} = -u_4^{(l+1)}. \end{cases} \quad (6.102)$$

For arbitrary $u^{(l+1)}$'s, this is solved by

$$\begin{cases} c_2^{(l+1)} = c_1^{(l+1)} - u_1^{(l+1)}, \\ c_4^{(l+1)} = -c_1^{(l+1)} + 2c_3^{(l+1)} + u_1^{(l+1)} + u_2^{(l+1)} - u_3^{(l+1)}, \\ c_5^{(l+1)} = c_1^{(l+1)} - c_3^{(l+1)} - u_1^{(l+1)} - u_2^{(l+1)}, \\ c_6^{(l+1)} = -c_1^{(l+1)} + u_1^{(l+1)} + u_2^{(l+1)} - u_4^{(l+1)}, \end{cases} \quad (6.103)$$

where $c_1^{(l+1)}$ and $c_3^{(l+1)}$ remain arbitrary. With the above choice we can make

$$\Sigma_A^{(l+1)} = 0. \quad (6.104)$$

We end this subsection by giving the results of 1-loop calculations. For simplicity, we take

$$K_b = K_f = K. \quad (6.105)$$

For the interaction action, we obtain

$$\left\{ \begin{array}{l} a^{(1)} = |g|^2 \int_q \frac{\Delta(q)}{q^2} \left(-\frac{1}{2} + K(q) \right), \\ c_1^{(1)}(\ln \Lambda/\mu) = \frac{|g|^2}{(4\pi)^2} \ln \frac{\Lambda}{\mu} + c_1^{(1)}(0), \\ c_2^{(1)}(\ln \Lambda/\mu) = \frac{|g|^2}{(4\pi)^2} \ln \frac{\Lambda}{\mu} + c_2^{(1)}(0), \\ c_3^{(1)}(\ln \Lambda/\mu) = c_3^{(1)}(0), \\ c_4^{(1)}(\ln \Lambda/\mu) = -\frac{|g|^2}{(4\pi)^2} \ln \frac{\Lambda}{\mu} + c_4^{(1)}(0), \\ c_5^{(1)}(\ln \Lambda/\mu) = \frac{|g|^2}{(4\pi)^2} \ln \frac{\Lambda}{\mu} + c_5^{(1)}(0), \\ c_6^{(1)}(\ln \Lambda/\mu) = -\frac{|g|^2}{(4\pi)^2} \ln \frac{\Lambda}{\mu} + c_6^{(1)}(0). \end{array} \right. \quad (6.106)$$

The coefficients $u_i^{(1)}$'s can be calculated from

$$\Sigma_A^{(1),u} = - \int_p K(p/\Lambda) \left[\bar{g} \left[\frac{\bar{\phi}^2}{2} \right]_A^{(0)}(p) \frac{\overleftarrow{\delta}}{\delta \chi_R(p)} \xi_R + g \left[\frac{\phi^2}{2} \right]_A^{(0)} \frac{\overleftarrow{\delta}}{\delta \chi_L(p)} \xi_L \right] \quad (6.107)$$

as follows (Fig. 22):

$$\left\{ \begin{array}{l} u_1^{(1)} = |g|^2 \int_p \frac{1}{p^4} \Delta(p) \left(\frac{1}{4} K(p) - \frac{1}{8} \Delta(p) \right), \\ u_2^{(1)} = 0, \\ u_3^{(1)} = -2|g|^2 \int_p \frac{1}{p^4} K(1-K)^2(2-K), \\ u_4^{(1)} = 0. \end{array} \right. \quad (6.108)$$

If we choose

$$c_2^{(1)}(0) = c_3^{(1)}(0) = 0, \quad (6.109)$$

then we obtain

$$\left\{ \begin{array}{l} c_1^{(1)}(0) = |g|^2 \int_p \frac{1}{p^4} \Delta \left(\frac{1}{4} K - \frac{1}{8} \Delta \right), \\ c_4^{(1)}(0) = 2|g|^2 \int_p \frac{1}{p^4} K(1-K)^2(2-K), \\ c_5^{(1)}(0) = c_6^{(1)}(0) = 0. \end{array} \right. \quad (6.110)$$

6.4. $O(N)$ non-linear sigma model

The first application of the ERG formalism to the two-dimensional $O(N)$ non-linear sigma model was done by Becchi.¹⁷⁾ The discussion here follows a recent unpublished work by Lütfüoğlu and Sonoda.^{*)}

^{*)} The ERG formalism has been applied to the supersymmetric non-linear sigma models extensively in 52), 54)–56).




Fig. 22. Feynman graphs for $u_i^{(1)}$'s — The solid curve marked K indicates the multiplication of the cutoff function K . The dot denotes the $\bar{\phi}^2$ vertex. We use a broken line for the scalar, a solid line for the spinor.

The classical action of the two-dimensional $O(N)$ non-linear sigma model is given by

$$\begin{aligned} S_{cl} &= -\frac{1}{2g} \int d^2x \left[g \partial_\mu \phi_i \partial_\mu \phi_i + \partial_\mu \sqrt{1 - g\phi^2} \partial_\mu \sqrt{1 - g\phi^2} \right] \\ &= \int d^2x \left[\frac{1}{2} \phi_i \partial^2 \phi_i - \partial^2 \left(\frac{\phi^2}{2} \right) \cdot \frac{1}{4} \ln(1 - g\phi^2) \right], \end{aligned} \quad (6.111)$$

where the index i runs from 1 to $N - 1$, the repeated indices are summed, and

$$\phi^2 \equiv \phi_i \phi_i = \sum_{i=1}^{N-1} \phi_i \phi_i. \quad (6.112)$$

The classical action is invariant under the following infinitesimal rotation:

$$\delta_\epsilon \phi_i = \epsilon_i \Phi_N, \quad (i = 1, \dots, N - 1) \quad (6.113)$$

where ϵ_i are arbitrary infinitesimal constants, and

$$\Phi_N \equiv \sqrt{1 - g\phi^2} \quad (6.114)$$

is the N -th component of an N -dimensional unit vector whose i -th component is $\sqrt{g} \phi_i$.

Note that the usual IR problem with the massless scalar does not arise here, since the Wilson action is constructed by integration of high momentum modes $p^2 > \Lambda^2$. Hence, we do not need to introduce a mass term (or an external field) to break the symmetry explicitly to $O(N-1)$.

The Wilson action is given by

$$S_\Lambda = S_{F,\Lambda} + S_{I,\Lambda}, \quad (6.115)$$

where the free action is given by

$$S_{F,\Lambda} = -\frac{1}{2} \int_p \frac{p^2}{K(p/\Lambda)} \phi_i(-p) \phi_i(p). \quad (6.116)$$

Hence, the cutoff propagator is

$$\langle \phi_i(p) \phi_j(-p) \rangle_{S_{F,\Lambda}} = \delta_{ij} \frac{K(k/\Lambda)}{k^2}. \quad (6.117)$$

The interaction action is defined by the Polchinski equation

$$-\Lambda \frac{\partial}{\partial \Lambda} S_{I,\Lambda} = \int_k \frac{\Delta(k/\Lambda)}{k^2} \frac{1}{2} \left\{ \frac{\delta S_{I,\Lambda}}{\delta \phi_i(-p)} \frac{\delta S_{I,\Lambda}}{\delta \phi_i(p)} + \frac{\delta^2 S_{I,\Lambda}}{\delta \phi_i(-p) \delta \phi_i(p)} \right\}, \quad (6.118)$$

and the asymptotic behavior

$$S_{I,\Lambda} \xrightarrow{\Lambda \rightarrow \infty} \int d^2x \left[\Lambda^2 a(\ln \Lambda/\mu; \phi^2/2) + A(\ln \Lambda/\mu; \phi^2/2) (-\partial^2) \frac{\phi^2}{2} + B(\ln \Lambda/\mu; \phi^2/2) \phi_i (-\partial^2) \phi_i \right], \quad (6.119)$$

where the coefficient functions can be expanded in powers of $\phi^2/2$ as

$$\begin{cases} a(\ln \Lambda/\mu; x) &= \sum_{n=1}^{\infty} \frac{x^n}{n!} a_n(\ln \Lambda/\mu), \\ A(\ln \Lambda/\mu; x) &= \sum_{n=1}^{\infty} \frac{x^n}{n!} A_n(\ln \Lambda/\mu), \\ B(\ln \Lambda/\mu; x) &= \sum_{n=0}^{\infty} \frac{x^n}{n!} B_n(\ln \Lambda/\mu). \end{cases} \quad (6.120)$$

The series for A starts from $n = 1$, since the $n = 0$ would correspond to a total derivative in the action. At tree level, we find

$$a(\ln \Lambda/\mu; x) = 0, \quad A(\ln \Lambda/\mu; x) = \frac{1}{4} \ln(1 - 2gx), \quad B(\ln \Lambda/\mu; x) = 0. \quad (6.121)$$

The infinitesimal transformation is given by

$$\delta \phi_i(p) = \epsilon_i [\Phi_N]_{\Lambda}(p), \quad (6.122)$$

where the composite operator $[\Phi_N]_{\Lambda}$ is the composite operator corresponding to the classical Φ_N . $[\Phi_N]_{\Lambda}$ satisfies an ERG differential equation, and is specified by the asymptotic behavior

$$\int_p e^{ipx} [\Phi_N]_{\Lambda}(p) \xrightarrow{\Lambda \rightarrow \infty} P(\ln \Lambda/\mu; \phi(x)^2/2), \quad (6.123)$$

where the function P can be expanded as

$$P(\ln \Lambda/\mu; x) = \sum_{n=0}^{\infty} \frac{x^n}{n!} P_n(\ln \Lambda/\mu). \quad (6.124)$$

At tree level, we find

$$P(\ln \Lambda/\mu; x) = \sqrt{1 - 2gx}. \quad (6.125)$$

Thus, the action is parameterized by two functions:

$$A(0; x), \quad B(0; x), \quad (6.126)$$

while the transformation is parameterized by a function

$$P(0; x). \quad (6.127)$$

Each Taylor coefficient of $A(0; x)$, $B(0; x)$ is a parameter of the Wilson action. Similarly, $P(0; x)$ parameterizes the transformation. Hence, there are an infinite number of parameters. We wish to show how the WT identity determines the three functions.

The WT composite operator is defined by

$$\Sigma_\Lambda \equiv \int_p K(p/\Lambda) \left[\frac{\delta S_\Lambda}{\delta \phi_i(p)} \delta \phi_i(p) + \frac{\delta}{\delta \phi_i(p)} \delta \phi_i(p) \right]. \quad (6.128)$$

This has the following asymptotic behavior:

$$\Sigma_\Lambda \xrightarrow{\Lambda \rightarrow \infty} \int d^2x \epsilon_i \phi_i \left[\partial_\mu \phi_j \partial_\mu \phi_j \cdot s_1(\ln \Lambda/\mu; \phi^2/2) \right. \quad (6.129)$$

$$\left. + \phi_j \partial^2 \phi_j \cdot s_2(\ln \Lambda/\mu; \phi^2/2) + (\phi_j \partial_\mu \phi_j)^2 \cdot s_3(\ln \Lambda/\mu; \phi^2/2) \right]. \quad (6.130)$$

We would like to fine-tune the three functions $A(0; x)$, $B(0; x)$, $P(0; x)$ so that

$$s_1(0; x) = s_2(0; x) = s_3(0; x) = 0. \quad (6.131)$$

We prove this possibility by mathematical induction on the number of loops.

Let us introduce loop expansions using the by now familiar notation

$$S_\Lambda = \sum_{l=0}^{\infty} S_\Lambda^{(l)}, \quad [\Phi_N]_\Lambda = \sum_{l=0}^{\infty} [\Phi_N]_\Lambda^{(l)}, \quad \Sigma_\Lambda = \sum_{l=0}^{\infty} \Sigma_\Lambda^{(l)}, \quad A(0; x) = \sum_{l=0}^{\infty} A^{(l)}(x), \dots \quad (6.132)$$

Suppose we have determined $S_\Lambda^{(0)}, \dots, S_\Lambda^{(l)}$ (or equivalently $A^{(0, \dots, l)}(x)$, $B^{(0, \dots, l)}(x)$) and $[\Phi_N]_\Lambda^{(0)}, \dots, [\Phi_N]_\Lambda^{(l)}$ (equivalently $P^{(0, \dots, l)}(x)$) so that

$$\Sigma_\Lambda^{(0)} = \dots = \Sigma_\Lambda^{(l)} = 0. \quad (6.133)$$

Then, the asymptotic behavior of $\Sigma_\Lambda^{(l+1)}$ is independent of Λ , and we can write it as

$$\Sigma_\Lambda^{(l+1)} \xrightarrow{\Lambda \rightarrow \infty} \int d^2x \epsilon_i \phi_i \left[\partial_\mu \phi_j \partial_\mu \phi_j \cdot s_1(\phi^2/2) \right. \quad (6.134)$$

$$\left. + \phi_j \partial^2 \phi_j \cdot s_2(\phi^2/2) + (\phi_j \partial_\mu \phi_j)^2 \cdot s_3(\phi^2/2) \right], \quad (6.135)$$

where $s_i(x)$ ($i = 1, 2, 3$) are functions of x , independent of $\ln \Lambda/\mu$.

The definition of Σ_Λ gives

$$\Sigma_\Lambda^{(l+1)} = \Sigma_\Lambda^{(l+1),t} + \Sigma_\Lambda^{(l+1),u}, \quad (6.136)$$

where

$$\Sigma_\Lambda^{(l+1),t} = \epsilon_i \int_p K(p/\Lambda) \left[\frac{\delta S_\Lambda^{(l+1)}}{\delta \phi_i(p)} [\Phi_N]_\Lambda^{(0)}(p) + \frac{\delta S_\Lambda^{(0)}}{\delta \phi_i(p)} [\Phi_N]_\Lambda^{(l+1)}(p) \right], \quad (6.137)$$

$$\Sigma_\Lambda^{(l+1),u} = \epsilon_i \int_p K(p/\Lambda) \left[\sum_{k=1}^l \frac{\delta S_\Lambda^{(k)}}{\delta \phi_i(p)} [\Phi_N]_\Lambda^{(l+1-k)}(p) + \frac{\delta [\Phi_N]_\Lambda^{(l)}(p)}{\delta \phi_i(p)} \right]. \quad (6.138)$$

Only $\Sigma_A^{(l+1),t}$ depends on $A^{(l+1)}(x)$, $B^{(l+1)}(x)$, $P^{(l+1)}(x)$, and $\Sigma_A^{(l+1),u}$ are determined by S_A and $[\Phi_N]_A$ up to l -loop.

Therefore, the functions $s_i(x)$ are given as the sum

$$s_i(x) = t_i(x) + u_i(x), \quad (i = 1, 2, 3) \quad (6.139)$$

where $t_i(x)$ are linear in $A^{(l+1)}(x)$, $B^{(l+1)}(x)$, $P^{(l+1)}(x)$, and $u_i(x)$ are determined by the lower loop functions. We obtain explicitly

$$t_1(x) = P^{(l+1)'} - (2A^{(l+1)'} + B^{(l+1)'})P^{(0)} - 2A^{(0)'}P^{(l+1)} - 2P^{(0)'}B^{(l+1)}, \quad (6.140)$$

$$t_2(x) = P^{(l+1)'} - 2(A^{(l+1)'} + B^{(l+1)'})P^{(0)} - 2A^{(0)'}P^{(l+1)} - 2P^{(0)'}B^{(l+1)}, \quad (6.141)$$

$$t_3(x) = P^{(l+1)''} - (A^{(l+1)''} + B^{(l+1)'})P^{(0)} - A^{(0)''}P^{(l+1)} - 2P^{(0)''}B^{(l+1)} - 2B^{(l+1)'}P^{(0)'}, \quad (6.142)$$

where the primes denote derivatives with respect to x . There is no relation among the $t(x)$'s. Thus, whatever $u(x)$'s are, we can solve the equations

$$s_i(x) = t_i(x) + u_i(x) = 0. \quad (i = 1, 2, 3) \quad (6.143)$$

Using

$$A^{(0)}(x) = \frac{1}{4} \ln(1 - 2gx), \quad B^{(0)}(x) = 0, \quad P^{(0)}(x) = \sqrt{1 - 2gx}. \quad (6.144)$$

the solution is obtained as follows:

$$B^{(l+1)}(x) = B^{(l+1)}(0) + \int_0^x dy \frac{-u_1(y) + u_2(y)}{\sqrt{1 - 2gy}}, \quad (6.145)$$

$$\begin{aligned} \frac{d}{dx} A^{(l+1)}(x) = & \frac{1}{(1 - 2gx)^2} \left[A^{(l+1)'}(0) \right. \\ & + \int_0^x dy \left\{ -2g^2 B^{(l)}(y) + (1 - 2gy) B^{(l)''}(y) \right. \\ & + g\sqrt{1 - 2gy} (-2u_1(y) + u_2(y)) + (1 - 2gy)^{\frac{3}{2}} (2u_1'(y) - u_2'(y)) \\ & \left. \left. - (1 - 2gy)^{\frac{3}{2}} u_3(y) \right\} \right], \end{aligned} \quad (6.146)$$

$$\begin{aligned} P^{(l+1)}(x) = & \sqrt{1 - 2gx} \left[P^{(l+1)}(0) + \int_0^x dy \left\{ 2A^{(l+1)'}(y) \right. \right. \\ & \left. \left. - \frac{2g}{1 - 2gy} B^{(l+1)}(y) + \frac{-2u_1(y) + u_2(y)}{\sqrt{1 - 2gy}} \right\} \right]. \end{aligned} \quad (6.147)$$

Note that

$$A^{(l+1)'}(0), \quad B^{(l+1)}(0), \quad P^{(l+1)}(0) \quad (6.148)$$

are left undetermined as constants of integration. This is expected, since $A^{(l+1)'}(0)$ normalizes the coupling g , $B^{(l+1)}(0)$ normalizes the field ϕ^i , and $P^{(l+1)}(0)$ normalizes




Fig. 23. The broken curve denotes the cutoff function K , and the solid the high-momentum propagator. Each curve follows the same $O(N)$ index $i = 1, \dots, N-1$. The left vertex is from $[\Phi_N]_A$, and the right from $S_{I,A}$.



Fig. 24. These graphs give no contribution to the u coefficients.

the composite operator $[\Phi_N]_A$. For example, we can adopt the convention

$$\left. \frac{\partial}{\partial x} A(0; x) \right|_{x=0} = -\frac{g}{2}, \quad B(0; 0) = 0, \quad P(0; 0) = 1, \quad (6.149)$$

or equivalently

$$\begin{cases} A^{(0)'}(0) = -\frac{g}{2}, & A^{(l)'}(0) = 0, & (l \geq 1) \\ B^{(l)}(0) = 0, & (l \geq 0) \\ P^{(0)}(0) = 1, & P^{(l)}(0) = 0. & (l \geq 0) \end{cases} \quad (6.150)$$

This concludes our perturbative proof of the existence of the $O(N)$ non-linear sigma model.

Before concluding this example, let us compute $B^{(1)'}(0)$ and $P^{(1)'}(0)$. (Note $A^{(1)'}(0) = 0$ by the above convention.) For this, we compute the asymptotic behavior of

$$\Sigma_A^{(1),u} = \int_p K(p/\Lambda) \frac{\delta [\Phi_N]_A^{(0)}(p)}{\delta \phi_i(p)} \quad (6.151)$$

up to third order in ϕ 's. The two relevant Feynman diagrams given in Fig. 23. Note that the two graphs in Fig. 24 give only the Λ^2 terms, but no contribution to the u coefficients. Ignoring the Λ^2 terms, we obtain

$$\begin{aligned} \Sigma_A^{(1),u} \xrightarrow{\Lambda \rightarrow \infty} & - \int d^2x \epsilon_i \phi_i \left[\partial_\mu \phi_j \partial_\mu \phi_j g^2 \left(\frac{1}{4\pi} + \int_q \frac{K(q)(1-K(q))}{q^2} \right) \right. \\ & \left. + \phi_j \partial^2 \phi_j g^2 \int_q \frac{1}{q^2} \left(\frac{1}{4} \Delta(q)^2 + 2K(q)(1-K(q)) \right) \right] \end{aligned} \quad (6.152)$$

up to third order in ϕ 's. Hence,

$$\begin{cases} u_1(0) = -g^2 \left(\frac{1}{4\pi} + \int_q \frac{1}{q^2} K(q)(1-K(q)) \right), \\ u_2(0) = -g^2 \int_q \frac{1}{q^2} \left(\frac{1}{4} \Delta(q)^2 + 2K(q)(1-K(q)) \right). \end{cases} \quad (6.153)$$

This gives

$$\begin{cases} B^{(1)'}(0) = u_2(0) - u_1(0) = g^2 \left(\frac{1}{4\pi} - \int_q \frac{1}{q^2} (K(1-K) + \frac{1}{4}\Delta^2) \right), \\ P^{(1)'}(0) = u_2(0) - 2u_1(0) = g^2 \left(\frac{1}{2\pi} - \int_q \frac{1}{q^2} \frac{1}{4}\Delta^2 \right). \end{cases} \quad (6.154)$$

6.5. The axial symmetry with a massive propagator

We consider a Wilson action for the spin $\frac{1}{2}$ field:

$$S_\Lambda = - \int_p \frac{1}{K(p/\Lambda)} \bar{\psi}(-p) (\not{p} + im) \psi(p) + S_{I,\Lambda}, \quad (6.155)$$

where the mass m is non-vanishing. The free part of the action breaks the axial symmetry explicitly. Nevertheless, the breaking can be compensated by $S_{I,\Lambda}$, and the theory can still possess the axial symmetry.

In the presence of the axial symmetry, the correlation functions in the continuum limit satisfy the following WT identity:

$$\begin{aligned} \sum_{i=1}^N [\langle \psi(p_1) \cdots \gamma_5 \psi(p_i) \cdots \psi(p_N) \bar{\psi}(-q_1) \cdots \bar{\psi}(-q_N) \rangle^\infty \\ + \langle \psi(p_1) \psi(p_i) \cdots \psi(p_N) \bar{\psi}(-q_1) \cdots \bar{\psi}(-q_i) \gamma_5 \bar{\psi}(-q_N) \rangle^\infty] = 0. \end{aligned} \quad (6.156)$$

The WT operator for the U(1) axial symmetry is given by

$$\begin{aligned} \Sigma_\Lambda \equiv \int_p K(p/\Lambda) \left[S_\Lambda \frac{\overleftarrow{\delta}}{\delta\psi(p)} \gamma_5 [\psi](p) + [\bar{\psi}](-p) \gamma_5 \frac{\overrightarrow{\delta}}{\delta\bar{\psi}(-p)} S_\Lambda \right. \\ \left. - \text{Tr} \gamma_5 \left\{ [\psi](p) \frac{\overleftarrow{\delta}}{\delta\psi(p)} + \frac{\overrightarrow{\delta}}{\delta\bar{\psi}(-p)} [\bar{\psi}](-p) \right\} \right] = 0, \end{aligned} \quad (6.157)$$

where

$$\begin{cases} [\psi](p) \equiv \frac{1}{K(p/\Lambda)} \psi(p) + \frac{1-K(p/\Lambda)}{\not{p}+im} \frac{\overrightarrow{\delta}}{\delta\psi(-p)} S_\Lambda \\ [\bar{\psi}](-p) \equiv \frac{1}{K(p/\Lambda)} \bar{\psi}(-p) + S_\Lambda \frac{\overleftarrow{\delta}}{\delta\bar{\psi}(p)} \frac{1-K(p/\Lambda)}{\not{p}+im} \end{cases} \quad (6.158)$$

are the composite operators corresponding to $\psi, \bar{\psi}$. We can further rewrite the above as

$$\begin{aligned} \Sigma_\Lambda = \int_p \left[S_\Lambda \frac{\overleftarrow{\delta}}{\delta\psi(p)} \gamma_5 \psi(p) + \bar{\psi}(-p) \gamma_5 \frac{\overrightarrow{\delta}}{\delta\bar{\psi}(-p)} S_\Lambda - \frac{K(1-K)}{p^2 + m^2} (-2im) \right. \\ \left. \times \text{Tr} \gamma_5 \left\{ \frac{\overrightarrow{\delta}}{\delta\bar{\psi}(-p)} S_\Lambda \cdot S_\Lambda \frac{\overleftarrow{\delta}}{\delta\psi(p)} + \frac{\overrightarrow{\delta}}{\delta\bar{\psi}(-p)} S_\Lambda \frac{\overleftarrow{\delta}}{\delta\psi(p)} \right\} \right]. \end{aligned} \quad (6.159)$$

This must vanish for the axial symmetry.

In the simplest case, the theory is free, and the Wilson action is quadratic:

$$S_\Lambda = - \int_p \bar{\psi}(-p) D_\Lambda(p) \psi(p) \quad (6.160)$$

Substituting this into $\Sigma_A = 0$, we obtain the Ginsparg-Wilson relation:⁴⁷⁾

$$\{D_A(p), \gamma_5\} = \frac{K(1-K)}{p^2 + m^2}(-2im)D_A(p)\gamma_5 D_A(p). \quad (6.161)$$

The right-hand side is non-vanishing only for $p^2 > \Lambda^2$ if we choose $K(p/\Lambda) = 1$ for $p^2 < \Lambda^2$. Expanding

$$D_A(p) = \not{p}A_\Lambda(p) + imB_\Lambda(p) \quad (6.162)$$

with scalar coefficients, the above relation gives

$$B_\Lambda(p) = \frac{K(1-K)}{p^2 + m^2} (p^2 A_\Lambda(p)^2 + m^2 B_\Lambda(p)^2). \quad (6.163)$$

Moreover, the Polchinski equation for S_A gives

$$\Lambda \frac{\partial}{\partial \Lambda} D_A(p) = -2 \frac{\Delta}{K} D + \frac{\Delta}{p^2 + m^2} D(\not{p} - im)D, \quad (6.164)$$

which implies

$$\Lambda \frac{\partial}{\partial \Lambda} A_\Lambda = -2 \frac{\Delta}{K} A_\Lambda + \frac{\Delta}{p^2 + m^2} (p^2 A_\Lambda^2 - m^2 B_\Lambda^2 + 2A_\Lambda B_\Lambda m^2), \quad (6.165a)$$

$$\Lambda \frac{\partial}{\partial \Lambda} B_\Lambda = -2 \frac{\Delta}{K} B_\Lambda + \frac{\Delta}{p^2 + m^2} (-p^2 A_\Lambda^2 + m^2 B_\Lambda^2 + 2A_\Lambda B_\Lambda p^2). \quad (6.165b)$$

To obtain a general solution, we cheat a little. In this case we know the general solution for the continuum limit:

$$\langle \psi(p) \bar{\psi}(-p) \rangle^\infty = \frac{c(p)}{\not{p}}, \quad (6.166)$$

where c is a scalar function, independent of Λ .^{*)} Using

$$\langle \psi(p) \bar{\psi}(-p) \rangle^\infty = \frac{1}{K(p/\Lambda)^2} \langle \psi(p) \bar{\psi}(-p) \rangle_{S_A} + \frac{1 - 1/K(p/\Lambda)}{\not{p} + im}, \quad (6.167)$$

we obtain

$$\frac{c(p)}{\not{p}} = \frac{1}{K(p/\Lambda)^2} \frac{1}{D_\Lambda(p)} + \frac{1 - 1/K(p/\Lambda)}{\not{p} + im}. \quad (6.168)$$

Hence,

$$\begin{aligned} \frac{1}{D_\Lambda(p)} &= K(p)^2 \left(\frac{c(p)}{\not{p}} - \frac{1 - 1/K(p/\Lambda)}{\not{p} + im} \right) \\ &= K^2 \{ c(\not{p} + im) - (1 - 1/K) \not{p} \} \frac{1}{\not{p}(\not{p} + im)} \\ &= K^2 \{ (c - 1 + 1/K) \not{p} + imc \} \frac{1}{\not{p}(\not{p} + im)}. \end{aligned} \quad (6.169)$$

^{*)} We assume parity.

Inverting this, we obtain

$$\begin{aligned}
D_\Lambda(p) &= \frac{1}{K^2} \frac{p^2 + im\cancel{p}}{(c-1+1/K)^2 p^2 + m^2 c^2} \{ (c-1+1/K)\cancel{p} - imc \} \\
&= \frac{1}{K^2} \frac{(p^2(c-1+1/K) + m^2 c) \cancel{p} - imcp^2 + imp^2(c-1+1/K)}{(c-1+1/K)^2 p^2 + m^2 c^2} \\
&= \frac{1}{K} \frac{(p^2(Kc+1-K) + Kcm^2) \cancel{p} + imp^2(1-K)}{(Kc+1-K)^2 p^2 + K^2 m^2 c^2}. \tag{6.170}
\end{aligned}$$

This gives

$$A_\Lambda(p) = \frac{1}{K} \frac{p^2(Kc+1-K) + m^2 Kc}{p^2(Kc+1-K)^2 + m^2 K^2 c^2} \tag{6.171a}$$

$$B_\Lambda(p) = \frac{1}{K} \frac{p^2(1-K)}{p^2(Kc+1-K)^2 + m^2 K^2 c^2} \tag{6.171b}$$

which satisfy both the Polchinski equation (6.165) and the Ginsparg-Wilson relation (6.163). For small and large momenta, $D_\Lambda(p)$ behaves as

$$D_\Lambda(p) \begin{cases} = \frac{\cancel{p}}{c(p)}, & (p^2 < \Lambda^2) \\ \simeq \frac{1}{K(p/\Lambda)} (\cancel{p} + im). & (p^2 \gg \Lambda^2) \end{cases} \tag{6.172}$$

For the simplest case

$$c(p) = 1, \tag{6.173}$$

we obtain

$$\begin{aligned}
D_\Lambda(p) &= \frac{1}{K} \frac{(p^2 + Km^2) \cancel{p} + imp^2(1-K)}{p^2 + K^2 m^2} \\
&= \frac{\cancel{p} + im}{K} + \frac{(1-K)m^2 \cancel{p} - im(p^2 + m^2 K)}{p^2 + m^2 K^2}. \tag{6.174}
\end{aligned}$$

In the above, we have seen that the Wilson action of a fully interacting theory must satisfy (6.159) (or its analog), which reduces to the GW relation (6.161) only for free theories. However, it is possible to build up a fully interacting theory starting from (6.161).^{58), 64)} Particular solutions of (6.159) that correspond to interacting theories have been constructed explicitly in (68), (69).

§7. Realization of symmetry in the antifield formalism

The purpose of this section is to introduce a method of symmetry realization more powerful than the WT identity discussed in the previous two sections. The method is an adaptation of the antifield formalism of Batalin and Vilkovisky^{16),49),51)} to the ERG formulation of field theory.^{60)–63)} We introduce only what is needed for perturbative construction of theories, postponing more general discussions till §10. Most results of this section were first obtained by Becchi.¹⁷⁾ Our presentation here follows §§6,7 of 17).*)

7.1. Quantum master equation

We introduce an external source ϕ^* in order to generate the symmetry transformation**) of ϕ as the functional derivative of the Wilson action with respect to ϕ^* . Following the general method by Batalin and Vilkovisky, we give ϕ^* the statistics opposite to that of ϕ . In what follows, we treat ϕ as a generic bosonic field (scalars and vectors) and ϕ^* as the corresponding fermionic antifield. Extension to a fermionic field and its bosonic antifield is straightforward.

In the antifield formalism, the transformation of ϕ is called the **BRST transformation**,***) and it is given as the derivative of the Wilson action with respect to the corresponding antifield:

$$\delta\phi(p) = \frac{\overrightarrow{\delta}}{\delta\phi^*(-p)} \bar{S}_A. \quad (7.1)$$

To show the presence of antifields explicitly, we use a bar above the Wilson action and other related quantities. At the vanishing antifields, the above transformation reduces to the infinitesimal transformation for the WT identity. The WT composite operator is generalized to the **quantum master operator**

$$\bar{\Sigma}_A \equiv \int_p K(p/\Lambda) \left[\frac{\delta \bar{S}_A}{\delta\phi(p)} \cdot \frac{\overrightarrow{\delta}}{\delta\phi^*(-p)} \bar{S}_A + \frac{\overrightarrow{\delta}}{\delta\phi^*(-p)} \frac{\delta \bar{S}_A}{\delta\phi(p)} \right], \quad (7.2)$$

which reduces to the WT composite operator at $\phi^* = 0$. Note that $\bar{\Sigma}_A$ is a composite operator. (Quick proof) $\frac{\overrightarrow{\delta}}{\delta\phi^*(-p)} \bar{S}_A$ is a composite operator of type (4.34). Hence, $\bar{\Sigma}_A$ is a composite operator of type (4.32).

We now replace the WT identity by the **quantum master equation**:

$$\bar{\Sigma}_A = 0, \quad (7.3)$$

which reduces to the WT identity at the vanishing antifields. The quantum master

*) For a formulation with the effective average action, see 92), for example.

**) The symmetry can be any continuous symmetry, either local or global. For the application of the BV formalism to global symmetries, see Refs. 36), 37) for example.

***) For YM theories, this BRST transformation coincides with the ordinary BRST transformation in the limit of vanishing antifields.

equation implies, for the correlation functions, the following identities:

$$\begin{aligned} & \sum_{i=1}^n \langle \phi(p_1) \cdots \delta \phi(p_i) \cdots \phi(p_n) \rangle^\infty \\ &= \sum_{i=1}^n \frac{\vec{\delta}}{\delta \phi^*(-p_i)} \left\langle \phi(p_1) \cdots \widehat{\phi(p_i)} \cdots \phi(p_n) \right\rangle^\infty = 0. \end{aligned} \quad (7.4)$$

Given a functional $\mathcal{O}_\Lambda[\phi, \phi^*]$ in general, we define its **quantum BRST transformation** by

$$\begin{aligned} \delta_Q \mathcal{O}_\Lambda \equiv \int_p K(p/\Lambda) & \left[\frac{\delta \bar{S}_\Lambda}{\delta \phi(p)} \cdot \frac{\vec{\delta}}{\delta \phi^*(-p)} \mathcal{O}_\Lambda + \frac{\vec{\delta}}{\delta \phi^*(-p)} \bar{S}_\Lambda \cdot \frac{\delta}{\delta \phi(p)} \mathcal{O}_\Lambda \right. \\ & \left. + \frac{\vec{\delta}}{\delta \phi^*(-p)} \frac{\delta}{\delta \phi(p)} \mathcal{O}_\Lambda \right]. \end{aligned} \quad (7.5)$$

This definition is obtained from $\bar{\Sigma}_\Lambda$ by changing $\bar{S}_\Lambda$ infinitesimally by $\mathcal{O}_\Lambda$. For example, we obtain

$$\delta_Q \phi(p) = K(p/\Lambda) \frac{\vec{\delta}}{\delta \phi^*(-p)} \bar{S}_\Lambda, \quad (7.6)$$

so that we can write

$$\bar{\Sigma}_\Lambda = \int_p \left[\frac{\delta \bar{S}_\Lambda}{\delta \phi(p)} \delta_Q \phi(p) + \frac{\vec{\delta}}{\delta \phi(p)} \delta_Q \phi(p) \right]. \quad (7.7)$$

The first term is the change of $\bar{S}_\Lambda$ under the infinitesimal field transformation $\delta_Q \phi$, and the second term is the jacobian.

Suppose $\mathcal{O}_\Lambda$ is a composite operator. Then,

$$\mathcal{O}'_\Lambda \equiv K(p/\Lambda) \left(\frac{\delta \bar{S}_\Lambda}{\delta \phi(p)} \mathcal{O}_\Lambda + \frac{\delta \mathcal{O}_\Lambda}{\delta \phi(p)} \right) \quad (7.8)$$

is a composite operator of type (4.32). We then note that

$$\mathcal{O}''_\Lambda \equiv \frac{\vec{\delta}}{\delta \phi^*(-p)} \bar{S}_\Lambda \cdot \mathcal{O}'_\Lambda + \frac{\vec{\delta}}{\delta \phi^*(-p)} \mathcal{O}'_\Lambda \quad (7.9)$$

is a composite operator of type (4.38). Since

$$\delta_Q \mathcal{O}_\Lambda = \int_p \mathcal{O}''_\Lambda - \bar{\Sigma}_\Lambda \cdot \mathcal{O}_\Lambda, \quad (7.10)$$

$\delta_Q \mathcal{O}_\Lambda$ is also a composite operator, if $\bar{\Sigma}_\Lambda = 0$. If $\bar{\Sigma}_\Lambda \neq 0$, $\delta_Q \mathcal{O}_\Lambda$ is not a composite operator in general.

Note also that if $\bar{\Sigma}_\Lambda = 0$, we can rewrite

$$\delta_Q \mathcal{O}_\Lambda = \exp[-\bar{S}_\Lambda] \int_p K(p/\Lambda) \frac{\delta}{\delta \phi(p)} \frac{\vec{\delta}}{\delta \phi^*(-p)} (\exp[\bar{S}_\Lambda] \mathcal{O}_\Lambda) \quad (7.11)$$

for an arbitrary functional $\mathcal{O}_\Lambda$, not necessarily a composite operator. Since

$$\int_p K(p/\Lambda) \frac{\delta}{\delta\phi(p)} \frac{\overrightarrow{\delta}}{\delta\phi^*(-p)} \quad (7.12)$$

is a fermionic differential operator acting on the functionals of ϕ and ϕ^* , its square vanishes:

$$\delta_Q \delta_Q \mathcal{O}_\Lambda = \exp[-\bar{S}_\Lambda] \left(\int_p K(p/\Lambda) \frac{\delta}{\delta\phi(p)} \frac{\overrightarrow{\delta}}{\delta\phi^*(-p)} \right)^2 (\exp[\bar{S}_\Lambda] \mathcal{O}_\Lambda) = 0. \quad (7.13)$$

Thus, we obtain the nilpotency:

$$\delta_Q^2 = 0, \quad (7.14)$$

if $\bar{S}_\Lambda = 0$.

To summarize, the quantum master equation (7.3), i.e., $\bar{S}_\Lambda = 0$, implies the following:

- (i) $\delta_Q \mathcal{O}_\Lambda$ is a composite operator, if $\mathcal{O}_\Lambda$ is.
- (ii) δ_Q is nilpotent, $\delta_Q^2 = 0$.

As an example of (i), we find, under the assumption $\bar{S}_\Lambda = 0$,

$$\delta_Q[\phi]_\Lambda(p) = \frac{\overrightarrow{\delta}}{\delta\phi^*(-p)} \bar{S}_\Lambda, \quad (7.15)$$

which is a composite operator.

Finally, we come to an important algebraic identity

$$\delta_Q \bar{S}_\Lambda = 0. \quad (7.16)$$

This simply follows from the definitions (7.2) of $\bar{S}_\Lambda$ and (7.5) of δ_Q . The identity is valid no matter what $\bar{S}_\Lambda$ is. Eq. (7.16) is the algebraic structure we have been missing in our discussion of the WT identity in §5. We will see, in §7.2, how useful this identity is for the realization of continuous symmetry in the ERG approach.

7.2. Outline of perturbative construction

We now outline a perturbative construction of the theory that satisfies the quantum master equation (7.3). Let us first introduce loop expansions:

$$\bar{S}_\Lambda = \sum_{l=0}^{\infty} \bar{S}_\Lambda^{(l)}, \quad \bar{\Sigma}_\Lambda = \sum_{l=0}^{\infty} \bar{\Sigma}_\Lambda^{(l)}. \quad (7.17)$$

If the theory is renormalizable, the asymptotic behavior of the Wilson action is given in the form

$$\bar{S}_{I,\Lambda} \xrightarrow{\Lambda \rightarrow \infty} \int d^D x \sum_i \Lambda^{D-x_i} c_i(\ln \Lambda/\mu) \mathcal{O}_i, \quad (7.18)$$

where $\mathcal{O}_i$ is a local polynomial of ϕ, ϕ^* , and their derivatives, and $x_i \leq D$ is the scale dimension of $\mathcal{O}_i$. The Wilson action is parameterized only by $c_i(0)$ that has $x_i = D$.

We introduce the loop expansion of $c_i(0)$ as

$$c_i(0) = \sum_{l=0}^{\infty} c_i^{(l)}. \quad (7.19)$$

Our starting point is the classical action $\bar{S}_{cl}$ which is independent of the cutoff Λ . Since the interaction part of the tree level action $\bar{S}_\Lambda^{(0)}$ is the sum of tree graphs consisting of the vertices of $\bar{S}_{cl}$ and high-momentum propagators, we obtain

$$\bar{S}_{cl} = \lim_{\Lambda \rightarrow \infty} \bar{S}_\Lambda^{(0)}. \quad (7.20)$$

We assume that this satisfies the classical master equation:

$$\bar{\Sigma}_{cl} \equiv \int_p \frac{\overrightarrow{\delta}}{\delta \phi^*(-p)} \bar{S}_{cl} \cdot \frac{\delta \bar{S}_{cl}}{\delta \phi(p)} = 0. \quad (7.21)$$

Since $\bar{\Sigma}_{cl}$ is the asymptotic part of $\bar{\Sigma}_\Lambda^{(0)}$, we obtain

$$\bar{\Sigma}_\Lambda^{(0)} = 0. \quad (7.22)$$

Now, our induction hypothesis is

$$\bar{\Sigma}_\Lambda^{(0)} = \dots = \bar{\Sigma}_\Lambda^{(l)} = 0. \quad (7.23)$$

For this we have adjusted the parameters of the theory up to l -loop order. In other words we have adjusted

$$c_i^{(0)}, \dots, c_i^{(l)}. \quad (x_i = D) \quad (7.24)$$

Since $\bar{\Sigma}_\Lambda$ is a composite operator satisfying the ERG differential equation, the induction hypothesis implies that $\bar{\Sigma}_\Lambda^{(l+1)}$ is independent of Λ for large Λ . In the following we wish to show that we can fine-tune $c_i^{(l+1)}$ so that

$$\lim_{\Lambda \rightarrow \infty} \bar{\Sigma}_\Lambda^{(l+1)} = 0, \quad (7.25)$$

which is equivalent to

$$\bar{\Sigma}_\Lambda^{(l+1)} = 0. \quad (7.26)$$

Hence, (7.25) completes the perturbative construction by mathematical induction.

To show the possibility of such fine-tuning, we must examine the structure of $\bar{\Sigma}_\Lambda^{(l+1)}$ further:

$$\begin{aligned} \bar{\Sigma}_\Lambda^{(l+1)} &= \int_p K(p/\Lambda) \left[\sum_{k=0}^{l+1} \frac{\overrightarrow{\delta}}{\delta \phi^*(-p)} \bar{S}_\Lambda^{(k)} \cdot \frac{\delta \bar{S}_\Lambda^{(l+1-k)}}{\delta \phi(p)} + \frac{\overrightarrow{\delta}}{\delta \phi^*(-p)} \frac{\delta \bar{S}_\Lambda^{(l)}}{\delta \phi(p)} \right] \\ &= \bar{\Sigma}_\Lambda^{(l+1),t} + \bar{\Sigma}_\Lambda^{(l+1),u}, \end{aligned} \quad (7.27)$$

where

$$\bar{\Sigma}_\Lambda^{(l+1),t} = \int_p K(p/\Lambda) \left[\frac{\vec{\delta}}{\delta\phi^*(-p)} \bar{S}_{cl} \cdot \frac{\delta \bar{S}_\Lambda^{(l+1)}}{\delta\phi(p)} + \frac{\vec{\delta}}{\delta\phi^*(-p)} \bar{S}_\Lambda^{(l+1)} \cdot \frac{\delta \bar{S}_{cl}}{\delta\phi(p)} \right], \quad (7.28)$$

$$\bar{\Sigma}_\Lambda^{(l+1),u} = \int_p K(p/\Lambda) \left[\sum_{k=1}^l \frac{\vec{\delta}}{\delta\phi^*(-p)} \bar{S}_\Lambda^{(k)} \cdot \frac{\delta \bar{S}_\Lambda^{(l+1-k)}}{\delta\phi(p)} + \frac{\vec{\delta}}{\delta\phi^*(-p)} \frac{\delta \bar{S}_\Lambda^{(l)}}{\delta\phi(p)} \right]. \quad (7.29)$$

For large Λ , $\bar{\Sigma}_\Lambda^{(l+1)}$ is independent of Λ . Hence, the Λ dependence of $\bar{\Sigma}_\Lambda^{(l+1),t}$ and $\bar{\Sigma}_\Lambda^{(l+1),u}$ must cancel for large Λ . Note that only $\bar{\Sigma}_\Lambda^{(l+1),t}$ depends on $c_i^{(l+1)}$, and $\bar{\Sigma}_\Lambda^{(l+1),u}$ is completely determined by the lower loop parameters.

For later convenience, we introduce the following notation. We first denote

$$\bar{\Sigma}_\infty^{(l+1)} \equiv \lim_{\Lambda \rightarrow \infty} \bar{\Sigma}_\Lambda^{(l+1)}. \quad (7.30)$$

We then denote the Λ independent part (namely the part that does not vanish at $\ln \Lambda/\mu = 0$) of the asymptotic behavior of $\bar{S}_\Lambda^{(l+1)}$ by $\bar{S}_\infty^{(l+1)}$ so that

$$\bar{S}_\infty^{(l+1)} = \int d^D x \sum_i c_i^{(l+1)} \mathcal{O}_i. \quad (7.31)$$

Similarly, we denote the Λ independent parts of $\bar{\Sigma}_\Lambda^{(l+1),t}$ and $\bar{\Sigma}_\Lambda^{(l+1),u}$ by $\bar{\Sigma}_\infty^{(l+1),t}$, $\bar{\Sigma}_\infty^{(l+1),u}$, respectively. In defining these, we ignore the part multiplied by positive powers of either Λ or $\ln \Lambda/\mu$. By definition, we obtain

$$\bar{\Sigma}_\infty^{(l+1)} = \bar{\Sigma}_\infty^{(l+1),t} + \bar{\Sigma}_\infty^{(l+1),u}. \quad (7.32)$$

Now, for large Λ , (7.28) gives

$$\bar{\Sigma}_\infty^{(l+1),t} = \int d^D x \left[\frac{\vec{\delta}}{\delta\phi^*} \bar{S}_{cl} \cdot \frac{\delta \bar{S}_\infty^{(l+1)}}{\delta\phi} + \frac{\vec{\delta}}{\delta\phi^*} \bar{S}_\infty^{(l+1)} \cdot \frac{\delta \bar{S}_{cl}}{\delta\phi} \right]. \quad (7.33)$$

Introducing the classical BRST transformation

$$\delta_{cl} \equiv \int d^D x \left[\frac{\vec{\delta}}{\delta\phi^*} \bar{S}_{cl} \cdot \frac{\delta}{\delta\phi} + \frac{\delta \bar{S}_{cl}}{\delta\phi} \frac{\vec{\delta}}{\delta\phi^*} \right], \quad (7.34)$$

we can rewrite the above as

$$\bar{\Sigma}_\infty^{(l+1),t} = \delta_{cl} \bar{S}_\infty^{(l+1)}. \quad (7.35)$$

The classical master equation (7.21) implies the nilpotency of the classical BRST transformation:

$$\delta_{cl}^2 = 0. \quad (7.36)$$

Hence, we obtain

$$\delta_{cl} \bar{\Sigma}_\infty^{(l+1),t} = 0. \quad (7.37)$$

We now consider the algebraic identity (7.16). At $(l+1)$ -loop, this gives

$$\int_p K(p/\Lambda) \left[\frac{\vec{\delta}}{\delta\phi^*(-p)} \bar{S}_\Lambda^{(0)} \cdot \frac{\delta}{\delta\phi(p)} \bar{\Sigma}_\Lambda^{(l+1)} + \frac{\delta}{\delta\phi(p)} \bar{S}_\Lambda^{(0)} \cdot \frac{\vec{\delta}}{\delta\phi^*(-p)} \bar{\Sigma}_\Lambda^{(l+1)} \right] = 0 \quad (7.38)$$

due to the induction hypothesis (7.23). Considering large Λ , we obtain

$$\delta_{cl} \bar{\Sigma}_\infty^{(l+1)} = 0. \quad (7.39)$$

This gives

$$\delta_{cl} \bar{\Sigma}_\infty^{(l+1),u} = 0, \quad (7.40)$$

using (7.32) and (7.37).

To summarize so far, our goal is to show that we can fine-tune the $(l+1)$ -loop parameters $c_i^{(l+1)}$ so that $\bar{\Sigma}_\infty^{(l+1)} = 0$. This condition is equivalent to solving

$$\delta_{cl} \bar{S}_\infty^{(l+1)} = -\bar{\Sigma}_\infty^{(l+1),u}, \quad (7.41)$$

where

1. δ_{cl} , defined by (7.34), satisfies nilpotency (7.36),
2. $\bar{\Sigma}_\infty^{(l+1),u}$ satisfies (7.40), i.e., closed under δ_{cl} .
3. $\bar{S}_\infty^{(l+1)}$ is given by (7.31), the most general local functional of fields with dimension D .

Thus, we have converted the problem of perturbative construction into a question of classical algebra.*) For (7.41) to have a solution, it is sufficient that $\delta_{cl}\mathcal{O} = 0$ implies $\mathcal{O} = \delta_{cl}\mathcal{O}'$; using a more proper mathematical language, the BRST cohomology defined by δ_{cl} must be trivial.

The final algebraic question (7.41) does not have any general solution. The answer depends on the set of available fields and antifields and the choice of the classical action $\bar{S}_{cl}$. But it is clear what we must do. We first consider the most general $\bar{\Sigma}_\infty^{(l+1),u}$ that satisfies (7.40), using the available set of fields and antifields. If it can be given as $\delta_{cl} \bar{S}_\infty^{(l+1)}$, the theory can be constructed. If not, the symmetry is anomalous, and we cannot construct a symmetric quantum theory based on the classical action.

For YM theories, the algebraic problem has a well-known solution. We refer the reader to 12)–15). In §8.2, we show a concrete solution of (7.41) for the SU(2) gauge theory.

*) This is called the BRST cohomology.

§8. Examples

8.1. QED

In §6.1 we have already shown how to construct QED with the help of a WT composite operator. Though we do not need the antifield formalism for assurance of the theory's existence, it is instructive to see how the formalism applies to QED.

Our starting point is the classical action:

$$\bar{S}_{cl} = S_{cl} + \int d^4x \left[A_\mu^* \frac{1}{i} \partial_\mu c + \frac{1}{\xi} \bar{c}^* \frac{1}{i} \partial_\mu A_\mu + \frac{1}{2\xi} (\bar{c}^*)^2 + e \bar{\psi}^* c \psi + e \bar{\psi} c \psi^* \right], \quad (8.1)$$

where S_{cl} is given by (6.1). Here, the fields with the superscript $*$ are antifields that have the opposite statistics to the respective conjugate fields. Let us tabulate them:

field	antifield	statistics	dimension
A_μ	A_μ^*	Fermi	2
$\bar{c}$	$\bar{c}^*$	Bose	2
ψ	$\bar{\psi}^*$	Bose	$\frac{3}{2}$
$\bar{\psi}$	ψ^*	Bose	$\frac{3}{2}$

The classical action satisfies the following classical master equation:

$$\int d^4x \left[\frac{\delta \bar{S}_{cl}}{\delta A_\mu} \frac{\overrightarrow{\delta}}{\delta A_\mu^*} \bar{S}_{cl} + \frac{\delta \bar{S}_{cl}}{\delta \bar{c}^*} \frac{\overrightarrow{\delta}}{\delta \bar{c}} \bar{S}_{cl} + \bar{S}_{cl} \frac{\overleftarrow{\delta}}{\delta \psi} \cdot \frac{\overrightarrow{\delta}}{\delta \bar{\psi}^*} \bar{S}_{cl} + \bar{S}_{cl} \frac{\overleftarrow{\delta}}{\delta \psi^*} \cdot \frac{\overrightarrow{\delta}}{\delta \bar{\psi}} \bar{S}_{cl} \right] = 0. \quad (8.2)$$

Except for the term quadratic in $\bar{c}^*$, the antifield dependence of the action is determined so that the derivative of the action with respect to the antifield gives the BRST transformation of the corresponding field:

$$\begin{cases} \delta A_\mu = \frac{\overrightarrow{\delta}}{\delta A_\mu^*} \bar{S}_{cl} = \frac{1}{i} \partial_\mu c, \\ \delta \bar{c} = \frac{\delta \bar{S}_{cl}}{\delta \bar{c}^*} = \frac{1}{\xi} \left(\frac{1}{i} \partial_\mu A_\mu + \bar{c}^* \right), \\ \delta \psi = \frac{\overrightarrow{\delta}}{\delta \bar{\psi}^*} \bar{S}_{cl} = e c \psi, \\ \delta \bar{\psi} = \bar{S}_{cl} \frac{\overleftarrow{\delta}}{\delta \psi^*} = -e c \bar{\psi}. \end{cases} \quad (8.3)$$

We note that all the antifields are coupled linearly to the dynamical fields. The $\psi^*, \bar{\psi}^*$ are coupled to the quadratic fields $\bar{\psi}c, c\psi$, respectively, but these are basically linear since the free field c acts almost like an external field.

Let us consider the dependence of the Wilson action on each antifield.

- (i) A_μ^* — This is coupled linearly to A_μ^* , contributing a term

$$\int_k A_\mu^*(-k) k_\mu c(k) \quad (8.4)$$

to the Wilson action. Since c is free, no further A_μ^* dependence is generated.

(ii) $\bar{c}^*$ — The linear coupling

$$\int_k \bar{c}^*(-k) \frac{1}{\xi} k_\mu A_\mu(k) \quad (8.5)$$

amounts to an external source

$$\frac{1}{\xi} k_\mu \bar{c}^*(-k) \quad (8.6)$$

coupled linearly to the gauge field $A_\mu(k)$. From the general result (2.37), the linear coupling generates the quadratic term:

$$\begin{aligned} & \frac{1}{2} \int_k \frac{1}{\xi} k_\mu \bar{c}^*(-k) \frac{1 - K(k/\Lambda)}{k^2} \left(\delta_{\mu\nu} - (1 - \xi) \frac{k_\mu k_\nu}{k^2} \right) \frac{1}{\xi} (-k_\nu) \bar{c}^*(k) \\ &= -\frac{1}{2\xi} \int_k (1 - K(k/\Lambda)) \bar{c}^*(-k) \bar{c}^*(k). \end{aligned} \quad (8.7)$$

Furthermore, the shift

$$\begin{aligned} A_\mu(k) &\longrightarrow A_\mu^{sh}(k) \equiv A_\mu(k) + \frac{1 - K(k/\Lambda)}{k^2} \left(\delta_{\mu\nu} - (1 - \xi) \frac{k_\mu k_\nu}{k^2} \right) \frac{1}{\xi} (-k_\nu) \bar{c}^*(k) \\ &= A_\mu(k) - \frac{1 - K(k/\Lambda)}{k^2} k_\mu \bar{c}^*(k) \end{aligned} \quad (8.8)$$

is generated in the interaction part $S_{I,\Lambda}$ of the Wilson action.

(iii) $\psi^*, \bar{\psi}^*$ — We can regard the free field c as part of the external source coupled to $\psi, \bar{\psi}$:

$$\int_p \left(e \int_k \bar{\psi}^*(-p-k) c(k) \cdot \psi(p) + \bar{\psi}(-p) \cdot e \int_k c(k) \psi^*(p-k) \right). \quad (8.9)$$

Hence, from the general result (2.37), we find first that the quadratic term

$$\int_p e \int_k \bar{\psi}^*(-p-k) c(k) \cdot \frac{1 - K(p/\Lambda)}{\not{p} + im} \cdot \int_q e \int_l c(l) \psi^*(p-l) \quad (8.10)$$

is generated, and second that the fields are shifted as

$$\begin{cases} \psi(p) &\longrightarrow \psi^{sh}(p) \equiv \psi(p) + \frac{1 - K(p/\Lambda)}{\not{p} + im} e \int_k c(k) \psi^*(p-k), \\ \bar{\psi}(-p) &\longrightarrow \bar{\psi}^{sh}(-p) \equiv \bar{\psi}(-p) + e \int_k \bar{\psi}^*(-p-k) c(k) \frac{1 - K(p/\Lambda)}{\not{p} + im}, \end{cases} \quad (8.11)$$

in $S_{I,\Lambda}$.

Altogether, we obtain the following action in the presence of antifields:⁶⁵⁾⁵³⁾

$$\begin{aligned} \bar{S}_\Lambda &= S_{F,\Lambda} [A_\mu, c, \bar{c}, \psi, \bar{\psi}] \\ &+ \int_k \left(A_\mu^*(-k) k_\mu c(k) + \bar{c}^*(-k) \frac{1}{\xi} k_\mu A_\mu(k) \right) + \frac{1}{2\xi} \int_k K(k/\Lambda) \bar{c}^*(-k) \bar{c}^*(k) \end{aligned}$$

$$\begin{aligned}
& + \int_{p,k} (e\bar{\psi}^*(-p-k)c(k) \cdot \psi(p) + \bar{\psi}(-p) \cdot ec(k)\psi^*(p-k)) \\
& + \int_{p,k,l} e\bar{\psi}^*(-p-k)c(k) \cdot \frac{1-K(p/\Lambda)}{\not{p}+im} \cdot ec(l)\psi^*(p-l) \\
& + S_{I,\Lambda} \left[A_\mu^{sh}(k), \psi^{sh}(p), \bar{\psi}^{sh}(-p) \right], \tag{8-12}
\end{aligned}$$

where $S_{F,\Lambda}$ is the free part defined by (6.4), and $S_{I,\Lambda}$ is the interaction part constructed in §6.1. The fields are shifted in the interaction part. If $S_A = S_{F,\Lambda} + S_{I,\Lambda}$, without shifts, satisfies the Polchinski equation for QED, so does $\bar{S}_A$ given above.^{*)}

We now wish to show that $\bar{S}_A$ thus constructed indeed satisfies the quantum master equation. The quantum master operator is defined by

$$\begin{aligned}
\bar{\Sigma}_A \equiv & \int_k K(k/\Lambda) \left[\frac{\vec{\delta}}{\delta A_\mu^*(-k)} \bar{S}_A \cdot \frac{\delta \bar{S}_A}{\delta A_\mu(k)} + \frac{\delta \bar{S}_A}{\delta \bar{c}^*(k)} \cdot \frac{\vec{\delta}}{\delta \bar{c}(-k)} \bar{S}_A \right] \\
& - \int_p K(p/\Lambda) \text{Tr} \left[\frac{\vec{\delta}}{\delta \bar{\psi}^*(-p)} \bar{S}_A \cdot \bar{S}_A \frac{\overleftarrow{\delta}}{\delta \psi(p)} + \frac{\vec{\delta}}{\delta \bar{\psi}^*(-p)} \bar{S}_A \frac{\overleftarrow{\delta}}{\delta \psi(p)} \right] \\
& + \int_p K(p/\Lambda) \text{Tr} \left[\frac{\vec{\delta}}{\delta \bar{\psi}(-p)} \bar{S}_A \cdot \bar{S}_A \frac{\overleftarrow{\delta}}{\delta \psi^*(p)} + \frac{\vec{\delta}}{\delta \bar{\psi}(-p)} \bar{S}_A \frac{\overleftarrow{\delta}}{\delta \psi^*(p)} \right]. \tag{8-13}
\end{aligned}$$

We now substitute (8-12) into the above to express it in terms of $S_{I,\Lambda}$. We skip the intermediate steps which are long but straightforward.^{**)} Toward the end, we use the anticommuting nature of c :

$$\int_{k,l} c(k)c(l)F(k,l) = 0 \tag{8-14}$$

for any symmetric integration kernel $F(k,l) = F(l,k)$. We finally obtain the result

$$\bar{\Sigma}_A = \Sigma_A \left[A_\mu^{sh}, c, \psi^{sh}, \bar{\psi}^{sh} \right], \tag{8-15}$$

where Σ_A is the WT composite operator of QED defined in §6.1, and the fields are shifted as (8-8) and (8-11). Since $\Sigma_A = 0$ for arbitrary field variables, we obtain

$$\bar{\Sigma}_A = 0. \tag{8-16}$$

8.2. YM theories

For YM theories, the antifield formalism is essential for proving the possibility of their construction. We recall, from §6.2, that we still have to derive the algebraic relations (6.69), necessary in order to show the possibility of realizing $\Sigma_A = 0$ by fine-tuning the parameters. The first application of the ERG formulation and antifield formalism was done by Becchi.¹⁷⁾ This work was extended subsequently in (26), (28)–

^{*)} In §11.1, we rederive this result using a functional method.

^{**)} The details of a similar calculation are given for the O(N) linear sigma model in Appendix E.

30),32) among others. Our presentation here follows closely §7 of Ref. 17) with quite a few additions for pedagogy.¹⁰⁹⁾

In the presence of the antifields, the classical action is given by

$$\begin{aligned} \bar{S}_{cl} = \int d^4x \left[-\frac{1}{4} (F_{\mu\nu}^a)^2 + \frac{1}{2\xi} \left(\bar{c}^{a*} + \frac{1}{i} \partial_\mu A_\mu^a \right)^2 \right. \\ \left. + \left(A_\mu^{a*} + \frac{1}{i} \partial_\mu \bar{c}^a \right) (D_\mu c)^a + c^{a*} \frac{g}{2i} f^{abc} c^b c^c \right]. \end{aligned} \quad (8.17)$$

For convenience, we tabulate the antifields below:

field	antifield	statistics	dimension	ghost number
A_μ^a	A_μ^{a*}	Fermi	2	-1
c^a	c^{a*}	Bose	2	-2
$\bar{c}^a$	$\bar{c}^{a*}$	Bose	2	0

The ghost numbers are assigned so that the action has zero ghost number. The classical action satisfies the classical master equation:

$$\int d^4x \left[\frac{\delta \bar{S}_{cl}}{\delta A_\mu^a} \frac{\overrightarrow{\delta}}{\delta A_\mu^{a*}} \bar{S}_{cl} + \frac{\delta \bar{S}_{cl}}{\delta \bar{c}^{a*}} \frac{\overrightarrow{\delta}}{\delta \bar{c}^a} \bar{S}_{cl} - \bar{S}_{cl} \frac{\overleftarrow{\delta}}{\delta c^a} \frac{\delta \bar{S}_{cl}}{\delta c^{a*}} \right] = 0. \quad (8.18)$$

As in the case of QED, the BRST transformation of a field is given by the functional derivative of the classical action with respect to the conjugate field:

$$\begin{cases} \delta A_\mu^a = \frac{\overrightarrow{\delta}}{\delta A_\mu^{a*}} \bar{S}_{cl} = \frac{1}{i} (D_\mu c)^a, \\ \delta \bar{c}^a = \frac{\delta \bar{S}_{cl}}{\delta \bar{c}^{a*}} = \frac{1}{\xi} \left(\frac{1}{i} \partial_\mu A_\mu^a + \bar{c}^{a*} \right), \\ \delta c^a = \frac{\delta \bar{S}_{cl}}{\delta c^{a*}} = \frac{g}{2i} f^{abc} c^b c^c. \end{cases} \quad (8.19)$$

We now consider the Wilson action $\bar{S}_A$ in the presence of antifields. As in QED, we can determine the dependence of the Wilson action on $\bar{c}^{a*}$ and A_μ^{a*} . Let us consider them one by one:

(i) $\bar{c}^{a*}$ — This antifield is coupled linearly to the gauge field as

$$\int_k \frac{1}{\xi} k_\mu \bar{c}^{a*}(k) A_\mu^a(-k). \quad (8.20)$$

This generates the quadratic term

$$-\frac{1}{2} \int_k (1 - K(k/\Lambda)) \bar{c}^{a*}(-k) \bar{c}^{a*}(k) \quad (8.21)$$

and the shift

$$A_\mu^a(k) \longrightarrow A_\mu^a(k) - \frac{1 - K(k/\Lambda)}{k^2} k_\mu \bar{c}^{a*}(k) \quad (8.22)$$

in the interaction part of the Wilson action. This dependence on $\bar{c}^{a*}$ is the same as in QED.

(ii) A_μ^{a*} — The ghost equation of motion (6.51) is now generalized to

$$k_\mu \frac{\overrightarrow{\delta}}{\delta A_\mu^{a*}(-k)} \bar{S}_\Lambda = -K(k/\Lambda) \frac{\overrightarrow{\delta}}{\delta \bar{c}^a(-k)} \bar{S}_\Lambda, \quad (8.23)$$

since the BRST transformation of A_μ^a is now given by

$$\delta A_\mu^a(k) = \frac{\overrightarrow{\delta}}{\delta A_\mu^{a*}(-k)} \bar{S}_\Lambda. \quad (8.24)$$

(8.23) implies that the dependence of $\bar{S}_\Lambda$ on A_μ^{a*} and $\bar{c}^a$ is through a linear combination^{*)}

$$A_\mu^{a*}(-k) - \frac{1}{K(k/\Lambda)} k_\mu \bar{c}^a(-k). \quad (8.25)$$

Now that we have determined the dependence on $\bar{c}^{a*}$ and A_μ^{a*} , the Wilson action has the following form:

$$\bar{S}_\Lambda = S_{F,\Lambda} + \bar{S}_{I,\Lambda}, \quad (8.26)$$

where $S_{F,\Lambda}$ is given by (6.42), and

$$\begin{aligned} \bar{S}_{I,\Lambda} = & \frac{1}{\xi} \int_k \bar{c}^{a*}(-k) k_\mu A_\mu^a(k) + \frac{1}{2\xi} \int_k K(k/\Lambda) \bar{c}^{a*}(-k) \bar{c}^{a*}(k) + \int_k A_\mu^{a*}(-k) k_\mu c^a(k) \\ & + S'_{I,\Lambda} \left[A_\mu^a(k) - \frac{1 - K(k/\Lambda)}{k^2} k_\mu \bar{c}^{a*}(k), A_\mu^{a*}(-k) - \frac{1}{K(k/\Lambda)} k_\mu \bar{c}^a(-k), \right. \\ & \left. c^a(k), c^{a*}(-k) \right]. \end{aligned} \quad (8.27)$$

$\bar{S}_\Lambda$ reduces to the Wilson action S_Λ of §6.2 for the vanishing antifields. Especially, we find

$$S'_{I,\Lambda} \left[A_\mu^a(k), -\frac{1}{K(k/\Lambda)} k_\mu \bar{c}^a(-k), c^a(k), 0 \right] = S_{I,\Lambda} [A_\mu^a(k), \bar{c}^a(-k), c^a(k)]. \quad (8.28)$$

The interaction action $\bar{S}_{I,\Lambda}$ is determined by the Polchinski equation

$$\begin{aligned} -\Lambda \frac{\partial}{\partial \Lambda} \bar{S}_{I,\Lambda} = & \int_k \frac{\Delta(k/\Lambda)}{k^2} \left[\left(\delta_{\mu\nu} - (1-\xi) \frac{k_\mu k_\nu}{k^2} \right) \right. \\ & \times \frac{1}{2} \left\{ \frac{\delta \bar{S}_{I,\Lambda}}{\delta A_\mu^a(-k)} \frac{\delta \bar{S}_{I,\Lambda}}{\delta A_\nu^a(k)} + \frac{\delta^2 \bar{S}_{I,\Lambda}}{\delta A_\mu^a(-k) \delta A_\nu^a(k)} \right\} \\ & \left. + \bar{S}_{I,\Lambda} \frac{\overleftarrow{\delta}}{\delta c^a(k)} \cdot \frac{\overrightarrow{\delta}}{\delta \bar{c}^a(-k)} \bar{S}_{I,\Lambda} - \frac{\overrightarrow{\delta}}{\delta \bar{c}^a(-k)} \bar{S}_{I,\Lambda} \frac{\overleftarrow{\delta}}{\delta c^a(k)} \right], \end{aligned} \quad (8.29)$$

^{*)} We recall that the Wilson action S_Λ of a YM theory depends on $\bar{c}^a(-k)$ only through the derivative $k_\mu \bar{c}^a(-k)$.

and an asymptotic behavior. For $SU(2)$, the latter is given in the following form:

$$\begin{aligned} \bar{S}_{I,\Lambda} \xrightarrow{\Lambda \rightarrow \infty} \int d^4x \left[\Lambda^2 a_0 (\ln \Lambda/\mu) \frac{1}{2} (A_\mu^a)^2 \right. \\ + a_1 (\ln \Lambda/\mu) \frac{1}{2} (\partial_\mu A_\nu^a)^2 + a_2 (\ln \Lambda/\mu) \frac{1}{2} (\partial_\mu A_\mu^a)^2 \\ + a_3 (\ln \Lambda/\mu) g \epsilon^{abc} \partial_\mu A_\nu^a \cdot A_\mu^b A_\nu^c \\ + a_4 (\ln \Lambda/\mu) \frac{g^2}{4} (A_\mu^a A_\mu^a)^2 + a_5 (\ln \Lambda/\mu) \frac{g^2}{4} (A_\mu^a A_\nu^a)^2 \\ + a_6 (\ln \Lambda/\mu) \left(A_\mu^{a*} + \frac{1}{i} \partial_\mu \bar{c}^a \right) \frac{1}{i} \partial_\mu c^a \\ + a_7 (\ln \Lambda/\mu) g \epsilon^{abc} \left(A_\mu^{a*} + \frac{1}{i} \partial_\mu \bar{c}^a \right) \frac{1}{i} A_\mu^b c^c \\ \left. + a_8 (\ln \Lambda/\mu) c^{a*} \frac{g}{2i} \epsilon^{abc} c^b c^c \right]. \end{aligned} \quad (8.30)$$

We note that all the coefficients are the same as in §6.2; as for $a_0, \dots, a_7$, they survive the limit of zero antifields, and as for a_8 , we must find

$$\frac{\delta \bar{S}_{I,\Lambda}}{\delta c^{a*}(-k)} \Big|_{A^*=c^*=\bar{c}^*=0} = C^a(k). \quad (8.31)$$

Hence, the introduction of the antifields does not increase the number of parameters of the theory. We still have the same eight parameters

$$a_1(0), \dots, a_8(0) \quad (8.32)$$

at our disposal.

Now, the quantum master operator is defined by

$$\begin{aligned} \bar{\Sigma}_\Lambda \equiv \int_k K(k/\Lambda) \left[\frac{\vec{\delta}}{\delta A_\mu^{a*}(-k)} \bar{S}_\Lambda \cdot \frac{\delta \bar{S}_\Lambda}{\delta A_\mu^a(k)} + \frac{\vec{\delta}}{\delta A_\mu^{a*}(-k)} \frac{\delta \bar{S}_\Lambda}{\delta A_\mu^a(k)} \right. \\ + \frac{\delta \bar{S}_\Lambda}{\delta \bar{c}^{a*}(k)} \frac{\vec{\delta}}{\delta \bar{c}^a(-k)} \bar{S}_\Lambda + \frac{\delta}{\delta \bar{c}^{a*}(k)} \frac{\vec{\delta}}{\delta \bar{c}^a(-k)} \bar{S}_\Lambda \\ \left. - \frac{\delta \bar{S}_\Lambda}{\delta c^{a*}(-k)} \cdot \bar{S}_\Lambda \frac{\overleftarrow{\delta}}{\delta c^a(k)} - \frac{\delta \bar{S}_\Lambda}{\delta c^{a*}(-k)} \frac{\overleftarrow{\delta}}{\delta c^a(k)} \right]. \end{aligned} \quad (8.33)$$

This reduces to the WT composite operator Σ_Λ (6.58) of §6.2 at the vanishing antifields. We wish to fine-tune the eight parameters (8.32) to make $\bar{\Sigma}_\Lambda$ vanish. The possibility of such fine-tuning can be proven if we can derive the algebraic constraints (6.69).

The derivation of (6.69) takes some steps. We first rewrite the quantum master operator using all we know about the antifield dependence of $\bar{S}_\Lambda$. Substituting (8.26) and (8.27) into the definition (8.33), we obtain

$$\bar{\Sigma}_\Lambda = \int_k K(k/\Lambda) \left[\left\{ \frac{\vec{\delta}}{\delta A_\mu^{a*}(-k)} \tilde{S}_\Lambda \cdot \frac{\delta \tilde{S}_\Lambda}{\delta A_\mu^a(k)} + \frac{\vec{\delta}}{\delta A_\mu^{a*}(-k)} \frac{\delta \tilde{S}_\Lambda}{\delta A_\mu^a(k)} \right\} P_{\mu\nu}(k/\Lambda) \right]$$

$$\left. -\frac{\delta \tilde{S}_\Lambda}{\delta c^{a*}(-k)} \cdot \tilde{S}_\Lambda \frac{\overleftarrow{\delta}}{\delta c^a(k)} - \frac{\delta \tilde{S}_\Lambda}{\delta c^{a*}(-k)} \frac{\overleftarrow{\delta}}{\delta c^a(k)} \right], \quad (8.34)$$

where we define

$$P_{\mu\nu}(k) \equiv \delta_{\mu\nu} + \frac{1 - K(k)}{K(k)} \frac{k_\mu k_\nu}{k^2}, \quad (8.35)$$

and

$$\begin{aligned} \tilde{S}_\Lambda &\equiv \bar{S}_\Lambda \\ &+ \frac{1}{2\xi} \int_k \frac{1}{K(k/\Lambda)} \{k_\mu A_\mu^a(k) + K(k/\Lambda) \bar{c}^{a*}(k)\} \{k_\nu A_\nu^a(-k) - K(k/\Lambda) \bar{c}^{a*}(-k)\} \\ &= -\frac{1}{2} \int_k \frac{1}{K(k/\Lambda)} A_\mu^a(k) A_\nu^a(-k) (k^2 \delta_{\mu\nu} - k_\mu k_\nu) \\ &\quad + \int_k \left\{ A_\mu^{a*}(-k) - \frac{1}{K(k/\Lambda)} k_\mu \bar{c}^a(-k) \right\} k_\mu c^a(k) \\ &\quad + S'_{I,\Lambda} \left[A_\mu^a(k) - \frac{1 - K(k/\Lambda)}{k^2} k_\mu \bar{c}^{a*}(k), A_\mu^{a*}(-k) - \frac{1}{K(k/\Lambda)} k_\mu \bar{c}^a(-k), \right. \\ &\quad \left. c^a(k), c^{a*}(-k) \right]. \end{aligned} \quad (8.36)$$

Note that the quadratic kinetic term of the gauge field in $\tilde{S}_\Lambda$ is purely transverse. Hence, $\tilde{S}_\Lambda$ depends only on the following combinations of the field variables:

$$A_\mu^a(k) - \frac{1 - K(k/\Lambda)}{k^2} k_\mu \bar{c}^{a*}(k), A_\mu^{a*}(-k) - \frac{1}{K(k/\Lambda)} k_\mu \bar{c}^a(-k), c^a(k), c^{a*}(-k). \quad (8.37)$$

Therefore, shifting A_μ^a and its antifield by terms proportional to k_μ , we define a modified quantum master operator by

$$\begin{aligned} &\hat{\Sigma}_\Lambda [A_\mu^a, A_\mu^{a*}, c^a, c^{a*}] \\ &\equiv \int_k K(k/\Lambda) \left[P_{\mu\nu}(k/\Lambda) \left\{ \frac{\overrightarrow{\delta}}{\delta A_\mu^{a*}(-k)} \hat{S}_\Lambda \cdot \frac{\delta \hat{S}_\Lambda}{\delta A_\nu^a(k)} + \frac{\overrightarrow{\delta}}{\delta A_\mu^{a*}(-k)} \frac{\delta \hat{S}_\Lambda}{\delta A_\nu^a(k)} \right\} \right. \\ &\quad \left. - \frac{\delta \hat{S}_\Lambda}{\delta c^{a*}(-k)} \cdot \hat{S}_\Lambda \frac{\overleftarrow{\delta}}{\delta c^a(k)} - \frac{\delta \hat{S}_\Lambda}{\delta c^{a*}(-k)} \frac{\overleftarrow{\delta}}{\delta c^a(k)} \right], \end{aligned} \quad (8.38)$$

where $\hat{S}_\Lambda$ is $\tilde{S}_\Lambda$ for the shifted variables, and defined by

$$\begin{aligned} &\hat{S}_\Lambda [A_\mu^a(k), A_\mu^{a*}(-k), c^a(k), c^{a*}(-k)] \\ &\equiv -\frac{1}{2} \int_k A_\mu^a(k) A_\nu^a(-k) \frac{1}{K(k/\Lambda)} (k^2 \delta_{\mu\nu} - k_\mu k_\nu) + \int_k A_\mu^{a*}(-k) k_\mu c^a(k) \\ &\quad + S'_{I,\Lambda} [A_\mu^a(k), A_\mu^{a*}(-k), c^a(k), c^{a*}(-k)]. \end{aligned} \quad (8.39)$$

Then, the quantum master equation

$$\bar{\Sigma}_\Lambda = 0 \quad (8.40)$$

is equivalent to the modified quantum master equation

$$\hat{\Sigma}_A = 0. \quad (8.41)$$

It is the modified action $\hat{S}_A$ (8.39) and the corresponding quantum master operator $\hat{\Sigma}_A$ (8.38) that we apply the general formalism developed in §7. Let us identify the classical algebraic problem corresponding to (7.41):

$$\hat{\delta}_{cl} \hat{S}_\infty^{(l+1)} = -\hat{\Sigma}_\infty^{(l+1),u}. \quad (8.42)$$

First, the classical action is given by

$$\hat{S}_{cl} \equiv \int d^4x \left[-\frac{1}{4} (F_{\mu\nu}^a)^2 + A_\mu^{a*} (D_\mu c)^a + c^{a*} \frac{g}{2i} \epsilon^{abc} c^b c^c \right], \quad (8.43)$$

which satisfies the classical master equation

$$\int d^4x \left[\frac{\delta \hat{S}_{cl}}{\delta A_\mu^a} \cdot \frac{\overrightarrow{\delta}}{\delta A_\mu^{a*}} \hat{S}_{cl} + \frac{\delta \hat{S}_{cl}}{\delta c^{a*}} \cdot \frac{\overrightarrow{\delta}}{\delta c^a} \hat{S}_{cl} \right] = 0. \quad (8.44)$$

Then, the classical BRST transformation is defined by

$$\hat{\delta}_{cl} \equiv \int d^4x \left[\frac{\delta \hat{S}_{cl}}{\delta A_\mu^a} \frac{\overrightarrow{\delta}}{\delta A_\mu^{a*}} + \frac{\overrightarrow{\delta}}{\delta A_\mu^{a*}} \hat{S}_{cl} \frac{\delta}{\delta A_\mu^a} + \frac{\delta \hat{S}_{cl}}{\delta c^{a*}} \frac{\overrightarrow{\delta}}{\delta c^a} - \hat{S}_{cl} \frac{\overleftarrow{\delta}}{\delta c^a} \frac{\delta}{\delta c^{a*}} \right], \quad (8.45)$$

which is nilpotent

$$\hat{\delta}_{cl}^2 = 0 \quad (8.46)$$

due to the classical master equation. The most general form of $\hat{S}_\infty^{(l+1)}$, for $SU(2)$, is given by

$$\begin{aligned} \hat{S}_\infty^{(l+1)} = \int d^4x & \left[a_1^{(l+1)} \frac{1}{2} (\partial_\mu A_\nu^a)^2 + a_2^{(l+1)} \frac{1}{2} (\partial_\mu A_\mu^a)^2 \right. \\ & + a_3^{(l+1)} g \epsilon^{abc} \partial_\mu A_\nu^a \cdot A_\mu^b A_\nu^c \\ & + a_4^{(l+1)} \frac{g^2}{4} (A_\mu^a A_\mu^a)^2 + a_5^{(l+1)} \frac{g^2}{4} (A_\mu^a A_\nu^a)^2 \\ & + a_6^{(l+1)} A_\mu^{a*} \frac{1}{i} \partial_\mu c^a + a_7^{(l+1)} g \epsilon^{abc} A_\mu^{a*} \frac{1}{i} A_\mu^b c^c \\ & \left. + a_8^{(l+1)} c^{a*} \frac{g}{2i} \epsilon^{abc} c^b c^c \right], \end{aligned} \quad (8.47)$$

where we use the familiar notation for the loop expansion

$$a_i(0) = \sum_{l=0}^{\infty} a_i^{(l)}. \quad (8.48)$$

Now, $\hat{\Sigma}_A$ is a dimension 5 fermionic scalar composite operator of ghost number +1. Hence, the most general form of $\hat{\Sigma}_\infty^{(l+1)}$ is given by

$$\hat{\Sigma}_\infty^{(l+1)} = \int d^4x \left[s_1 \frac{1}{i} c^a \partial^2 \partial_\mu A_\mu^a \right]$$

$$\begin{aligned}
& +s_2 \frac{1}{i} g \epsilon^{abc} c^a \partial^2 A_\mu^b \cdot A_\mu^c + s_3 \frac{1}{i} g \epsilon^{abc} c^a \partial_\mu \partial_\nu A_\nu^b \cdot A_\mu^c \\
& +s_4 i g^2 c^a \partial_\mu A_\mu^a \cdot A_\nu^b A_\nu^b + s_5 i g^2 c^a \partial_\mu A_\nu^a \cdot A_\mu^b A_\nu^b + s_6 i g^2 c^a A_\mu^a \partial_\nu A_\nu^b A_\mu^b \\
& +s_7 i g^2 c^a A_\mu^a \partial_\nu A_\mu^b A_\nu^b + s_8 i g^2 c^a A_\mu^a \partial_\mu A_\nu^b A_\nu^b \\
& +s_9 g \epsilon^{abc} \partial_\mu c^a c^b A_\mu^{c*} + s_{10} g^2 c^a c^b A_\mu^{a*} A_\mu^a \Big] , \tag{8.49}
\end{aligned}$$

where $s_1, \dots, s_{10}$ are constants. Replacing A_μ^{a*} by $\frac{1}{i} \partial_\mu \bar{c}^a$, this is the same as the asymptotic behavior of $\Sigma_A^{(l+1)}$ in the absence of the antifields. (See (6.59).) Similarly, $\hat{\Sigma}_\infty^{(l+1),t}$ and $\hat{\Sigma}_\infty^{(l+1),u}$ are parameterized by t_i and u_i , respectively, instead of s_i . Then,

$$\hat{\Sigma}_\infty^{(l+1)} = \hat{\Sigma}_\infty^{(l+1),t} + \hat{\Sigma}_\infty^{(l+1),u} \tag{8.50}$$

implies

$$s_i = t_i + u_i \quad (i = 1, \dots, 10) \tag{8.51}$$

Since

$$\hat{\Sigma}_\infty^{(l+1),t} = \hat{\delta}_{cl} \hat{S}_\infty^{(l+1)}, \tag{8.52}$$

t_i is determined by $a^{(l+1)}$'s. But u_i is determined only by the lower loop parameters.

Let us consider t_i 's. (8.52) gives

$$t_1 = a_1^{(l+1)} + a_2^{(l+1)} \tag{8.53a}$$

$$t_2 = a_1^{(l+1)} + a_3^{(l+1)} + a_6^{(l+1)} + a_7^{(l+1)} \tag{8.53b}$$

$$t_4 = a_3^{(l+1)} + a_4^{(l+1)} - a_6^{(l+1)} - a_7^{(l+1)} \tag{8.53c}$$

$$t_5 = -a_3^{(l+1)} + a_5^{(l+1)} + a_6^{(l+1)} + a_7^{(l+1)} \tag{8.53d}$$

$$t_9 = a_7^{(l+1)} + a_8^{(l+1)}, \tag{8.53e}$$

and

$$t_3 = t_1 - t_2, \tag{8.54a}$$

$$t_6 = t_5, \tag{8.54b}$$

$$t_7 = 2t_4, \tag{8.54c}$$

$$t_8 = t_5, \tag{8.54d}$$

$$t_{10} = -t_9. \tag{8.54e}$$

These are the same results as (6.67) & (6.68) that we have found in §6.2.

Next, we consider u_i 's. The algebraic constraint

$$\hat{\delta}_{cl} \hat{\Sigma}_\infty^{(l+1),u} = 0, \tag{8.55}$$

implies

$$u_3 = u_1 - u_2, \tag{8.56a}$$

$$u_6 = u_5 , \quad (8.56b)$$

$$u_7 = 2u_4 , \quad (8.56c)$$

$$u_8 = u_5 , \quad (8.56d)$$

$$u_{10} = -u_9 . \quad (8.56e)$$

These are the relations that were missing in §6.2.

Thus, $\hat{\Sigma}_\infty^{(l+1)} = 0$ gives only five independent equations

$$s_1 = s_2 = s_4 = s_5 = s_9 = 0 . \quad (8.57)$$

The solution is given by

$$a_2^{(l+1)} = -u_1 - a_1^{(l+1)} \quad (8.58a)$$

$$a_4^{(l+1)} = -u_2 - \left(a_1^{(l+1)} + a_3^{(l+1)} + a_6^{(l+1)} \right) \quad (8.58b)$$

$$a_5^{(l+1)} = -u_2 - u_4 - a_1^{(l+1)} - 2a_3^{(l+1)} \quad (8.58c)$$

$$a_7^{(l+1)} = u_2 - u_5 + a_1^{(l+1)} + 2a_3^{(l+1)} \quad (8.58d)$$

$$a_8^{(l+1)} = u_2 - u_9 + a_1^{(l+1)} + a_3^{(l+1)} + a_6^{(l+1)} , \quad (8.58e)$$

where we have treated the normalization constants $a_{1,3,6}^{(l+1)}$ as independent parameters. The simplest choice is to set them zero for $l \geq 0$.

§9. Axial and chiral anomalies

In this section we formulate the axial and chiral anomalies in the ERG formalism. We restrict our discussions only to abelian anomalies, and it suffices to use only the WT identity of §5. The anomalies have been computed in the ERG formalism in Refs. 25), 34), 92), but here we follow an unpublished work by Igarashi, Itoh, and Sonoda. For references, we would like to mention that the chiral anomalies have been studied using the antifield formalism (but not ERG) in 114), 38), and using the BRST cohomology (but not ERG) in 12)–15).

9.1. Axial anomaly in QED

In the following we wish to consider the axial vector current $J_{5,\mu}$ in QED, and compute its divergence

$$k_\mu J_{5,\mu}(-k). \quad (9.1)$$

To define the axial vector current precisely, it is convenient to couple an external field B_μ to the axial vector current. In the Wilson action, we only consider B_μ up to first order. B_μ introduces the following extra terms into the asymptotic behavior of the action:

$$S_{I,\Lambda} \xrightarrow{\Lambda \rightarrow \infty} \int d^4x \left[\cdots + a'_3(\ln \Lambda/\mu) \bar{\psi} \gamma_5 \not{B} \psi + a_5(\ln \Lambda/\mu) \epsilon_{\alpha\beta\gamma\delta} B_\alpha A_\beta \frac{1}{i} \partial_\gamma A_\delta \right], \quad (9.2)$$

where we have suppressed the terms independent of B_μ . We define the axial vector current as the composite operator:

$$J_{5,\mu}(-k) \equiv \frac{\delta S_{I,\Lambda}}{\delta B_\mu(k)}. \quad (9.3)$$

This has the asymptotic behavior

$$\begin{aligned} J_{5,\mu}(-k) &\xrightarrow{\Lambda \rightarrow \infty} a'_3(\ln \Lambda/\mu) \int_p \bar{\psi}(-p-k) \gamma_5 \gamma_\mu \psi(p) \\ &\quad + a_5(\ln \Lambda/\mu) \epsilon_{\mu\alpha\beta\gamma} \int_{k_1+k_2=-k} A_\alpha(k_1) k_{2\beta} A_\gamma(k_2). \end{aligned} \quad (9.4)$$

The parameter $a'_3(0)$ normalizes the axial vector current, and the ratio $a_5(0)/a'_3(0)$ is determined by the WT identity of QED as we will see shortly. The above asymptotic behavior implies

$$\begin{aligned} k_\mu J_{5,\mu}(-k) &\xrightarrow{\Lambda \rightarrow \infty} a'_3(\ln \Lambda/\mu) \int_p \bar{\psi}(-p-k) \gamma_5 \not{k} \psi(p) \\ &\quad + a_5(\ln \Lambda/\mu) \epsilon_{\alpha\beta\gamma\delta} \int_{k_1+k_2=-k} k_{1\alpha} k_{2\beta} A_\gamma(k_1) A_\delta(k_2). \end{aligned} \quad (9.5)$$

To determine $a_5(0)$, we consider the WT identity in the presence of B_μ . In §6.1 we have found that the WT identity is given by

$$k_\mu J_\mu(-k) = \Phi(-k). \quad (9.6)$$

The two composite operators are defined by

$$J_\mu(-k) \equiv \frac{\delta S_{I,\Lambda}}{\delta A_\mu(k)}, \quad (9.7)$$

and

$$\begin{aligned} \Phi(-k) \equiv e \int_p K(p/\Lambda) & \left[-S_\Lambda \frac{\overleftarrow{\delta}}{\delta \psi(p)} [\psi]_\Lambda(p-k) + [\bar{\psi}]_\Lambda(-p-k) \frac{\overrightarrow{\delta}}{\delta \bar{\psi}(-p)} S_\Lambda \right. \\ & \left. + \text{Tr} [\psi]_\Lambda(p-k) \frac{\overleftarrow{\delta}}{\delta \psi(p)} - \text{Tr} \frac{\overrightarrow{\delta}}{\delta \bar{\psi}(-p)} [\bar{\psi}]_\Lambda(-p-k) \right]. \end{aligned} \quad (9.8)$$

In the presence of B_μ , the asymptotic behaviors of both $k_\mu J_\mu(-k)$ and $\Phi(-k)$ get an extra term linear in B_μ :

$$k_\mu J_\mu(-k) \xrightarrow{\Lambda \rightarrow \infty} \dots - a_5(\ln \Lambda/\mu) \epsilon_{\alpha\beta\gamma\delta} \int_{k_1+k_2=-k} k_{1\gamma} k_{2\delta} B_\alpha(k_1) A_\beta(k_2), \quad (9.9)$$

$$\Phi(-k) \xrightarrow{\Lambda \rightarrow \infty} \dots + b(\ln \Lambda/\mu) \epsilon_{\alpha\beta\gamma\delta} \int_{k_1+k_2=-k} k_{1\gamma} k_{2\delta} B_\alpha(k_1) A_\beta(k_2), \quad (9.10)$$

where the terms independent of B_μ are suppressed. Thus, the gauge invariance requires

$$a_5(0) = -b(0). \quad (9.11)$$

We leave open the convention for the normalization constant $a'_3(0)$ of the axial vector current for later convenience.

We now consider the composite operator of type (4.32) at $B_\mu = 0$:

$$\begin{aligned} \Phi_5(-k) \equiv \int_p K(p/\Lambda) & \left[S_\Lambda \frac{\overleftarrow{\delta}}{\delta \psi(p)} \gamma_5 [\psi]_\Lambda(p-k) + [\bar{\psi}]_\Lambda(p-k) \gamma_5 \frac{\overrightarrow{\delta}}{\delta \bar{\psi}(-p)} S_\Lambda \right. \\ & \left. - \text{Tr} \gamma_5 [\psi]_\Lambda(p-k) \frac{\overleftarrow{\delta}}{\delta \psi(p)} - \text{Tr} \frac{\overrightarrow{\delta}}{\delta \bar{\psi}(-p)} [\bar{\psi}]_\Lambda(p-k) \gamma_5 \right]. \end{aligned} \quad (9.12)$$

This has the correlation functions:

$$\begin{aligned} & \langle \Phi_5(-k) A_{\mu_1}(k_1) \dots A_{\mu_M}(k_M) \psi(p_1) \dots \psi(p_N) \bar{\psi}(-q_1) \dots \bar{\psi}(-q_N) \rangle^\infty \\ & = - \sum_{i=1}^N \left\{ \langle \dots \gamma_5 \psi(p_i - k) \dots \rangle^\infty + \langle \dots \bar{\psi}(-q_i - k) \gamma_5 \dots \rangle^\infty \right\}. \end{aligned} \quad (9.13)$$

The asymptotic behavior of $\Phi_5(-k)$ has the following form:*)

$$\Phi_5(-k) \xrightarrow{\Lambda \rightarrow \infty} a'(\ln \Lambda/\mu) \int_p \bar{\psi}(-p-k) \gamma_5 \not{k} \psi(p)$$

*) Φ_5 is a pseudo scalar even under charge conjugation. Hence, the dimension 4 pseudo scalars $\bar{\psi} \gamma_5 \not{A} \psi$, $\bar{\psi} \gamma_5 \overleftrightarrow{\not{D}} \psi$ are forbidden.

$$\begin{aligned}
& +b'(\ln \Lambda/\mu)\epsilon_{\alpha\beta\gamma\delta} \int_{k_1+k_2=-k} k_{1\alpha}k_{2\beta}A_\gamma(k_1)A_\delta(k_2) \\
& +c'(\ln \Lambda/\mu)im \int_p \bar{\psi}(-p-k)\gamma_5\psi(p). \tag{9.14}
\end{aligned}$$

To rewrite Φ_5 , we introduce two composite operators:

(i) $[\frac{1}{2}F\tilde{F}]_\Lambda$ with the asymptotic behavior

$$\begin{aligned}
\left[F\tilde{F}/2\right]_\Lambda(-k) & \xrightarrow{\Lambda\rightarrow\infty} f_1(\ln \Lambda/\mu)\epsilon_{\alpha\beta\gamma\delta} \int_{k_1+k_2=-k} k_{1\alpha}k_{2\beta}A_\gamma(k_1)A_\delta(k_2) \\
& +f_2(\ln \Lambda/\mu) \int_p \bar{\psi}(-p-k)\gamma_5\not{k}\psi(p), \tag{9.15}
\end{aligned}$$

where we choose the convention

$$f_1(0) = 1, \quad f_2(0) = 0. \tag{9.16}$$

(ii) J_5 with the asymptotic behavior

$$J_5(-k) \xrightarrow{\Lambda\rightarrow\infty} j(\ln \Lambda/\mu) \int_p \bar{\psi}(-p-k)\gamma_5\psi(p). \tag{9.17}$$

We will make a convenient choice for $j(0)$ shortly.

We can now expand

$$\Phi_5(-k) = s k_\mu J_{5,\mu}(-k) + t \left[F\tilde{F}/2\right]_\Lambda + u \cdot im J_5(-k), \tag{9.18}$$

where s, t, u are Λ independent constants, dependent only on e^2 and ξ . The right-hand side has the asymptotic behavior

$$\begin{aligned}
(\text{RHS}) & \xrightarrow{\Lambda\rightarrow\infty} (s a'_3(\ln \Lambda/\mu) + t f_2(\ln \Lambda/\mu)) \int_p \bar{\psi}(-p-k)\gamma_5\not{k}\psi(p) \\
& + (s a'_5(\ln \Lambda/\mu) + t f_1(\ln \Lambda/\mu)) \epsilon_{\alpha\beta\gamma\delta} \int_{k_1+k_2=-k} k_{1\alpha}k_{2\beta}A_\gamma(k_1)A_\delta(k_2) \\
& + u j(\ln \Lambda/\mu)im \int_p \bar{\psi}(-p-k)\gamma_5\psi(p). \tag{9.19}
\end{aligned}$$

Comparing this with (9.14), we obtain

$$\begin{cases} a'(\ln \Lambda/\mu) &= s a'_3(\ln \Lambda/\mu) + t f_2(\ln \Lambda/\mu), \\ b'(\ln \Lambda/\mu) &= -s b(\ln \Lambda/\mu) + t f_1(\ln \Lambda/\mu), \\ c'(\ln \Lambda/\mu) &= u j(\ln \Lambda/\mu), \end{cases} \tag{9.20}$$

where we have used (9.11). By adopting the convention

$$a'_3(0) = a'(0), \tag{9.21}$$

$$j(0) = c'(0), \tag{9.22}$$

we obtain

$$s = 1, \quad (9.23)$$

$$t = b(0) + b'(0), \quad (9.24)$$

$$u = 1. \quad (9.25)$$

Hence, we obtain

$$\Phi_5(-k) = k_\mu J_{5,\mu}(-k) + im J_5(-k) + (b(0) + b'(0)) \left[F\tilde{F}/2 \right] (-k). \quad (9.26)$$

The last term gives the axial anomaly. We will show the 1-loop calculations of the axial anomaly in the next subsection. From (9.48, 9.50), setting $e' = 1$, we obtain

$$b(0) = e^2 \frac{8}{3} \frac{1}{(4\pi)^2}, \quad b'(0) = e^2 \frac{4}{3} \frac{1}{(4\pi)^2}, \quad (9.27)$$

so that the coefficient of the axial anomaly is

$$b(0) + b'(0) = \frac{e^2}{4\pi^2}. \quad (9.28)$$

9.2. Chiral anomaly

9.2.1. $U(1)_V \times U(1)_A$ gauge theory

Let us try to construct a $U(1)_V \times U(1)_A$ gauge theory with massless fermions. Its classical action is given by

$$S_{cl} = \int d^4x \left[-\frac{1}{4} F_{\mu\nu}^2 - \frac{1}{2\xi} (\partial_\mu A_\mu)^2 - \frac{1}{4} G_{\mu\nu}^2 - \frac{1}{2\xi'} (\partial_\mu B_\mu)^2 \right. \\ \left. - \partial_\mu \bar{c} \partial_\mu c - \partial_\mu \bar{c}' \partial_\mu c' + \bar{\psi} \left(-\frac{1}{i} \not{D} + e \not{A} + e' \gamma_5 \not{B} \right) \psi \right], \quad (9.29)$$

where

$$F_{\mu\nu} \equiv \partial_\mu A_\nu - \partial_\nu A_\mu, \quad G_{\mu\nu} \equiv \partial_\mu B_\nu - \partial_\nu B_\mu. \quad (9.30)$$

This action is invariant under the following classical BRST transformation:

$$\begin{cases} \delta_\epsilon A_\mu = \epsilon \frac{1}{i} \partial_\mu c, & \delta_\epsilon B_\mu = \epsilon \frac{1}{i} \partial_\mu c', \\ \delta_\epsilon c = 0, & \delta_\epsilon c' = 0, \\ \delta_\epsilon \bar{c} = \epsilon \frac{1}{\xi} \frac{1}{i} \partial_\mu A_\mu, & \delta_\epsilon \bar{c}' = \epsilon \frac{1}{\xi'} \frac{1}{i} \partial_\mu B_\mu, \\ \delta_\epsilon \psi = \epsilon (ec - e'c' \gamma_5) \psi, & \delta_\epsilon \bar{\psi} = \bar{\psi} \epsilon (-ec - e'c' \gamma_5), \end{cases} \quad (9.31)$$

where ϵ is an arbitrary anticommuting constant.

The quantization follows the same line as for QED. We omit writing down the Polchinski equation for the interaction action $S_{I,\Lambda}$. Its asymptotic behavior is given in the following form:

$$S_{I,\Lambda} \xrightarrow{\Lambda \rightarrow \infty} \int d^4x \left[\Lambda^2 a_2 (\ln \Lambda / \mu) \frac{1}{2} A_\mu^2 + \Lambda^2 a'_2 (\ln \Lambda / \mu) \frac{1}{2} B_\mu^2 \right]$$

$$\begin{aligned}
& +c_2(\ln \Lambda/\mu)\frac{1}{2}\partial_\mu A_\nu\partial_\mu A_\nu + d_2(\ln \Lambda/\mu)\frac{1}{2}(\partial_\mu A_\mu)^2 \\
& +c'_2(\ln \Lambda/\mu)\frac{1}{2}\partial_\mu B_\nu\partial_\mu B_\nu + d'_2(\ln \Lambda/\mu)\frac{1}{2}(\partial_\mu B_\mu)^2 \\
& +a_4(\ln \Lambda/\mu)\frac{1}{8}(A_\mu^2)^2 + a'_4(\ln \Lambda/\mu)\frac{1}{8}(B_\mu^2)^2 + a''_4(\ln \Lambda/\mu)\frac{1}{4}A_\mu^2 B_\mu^2 \\
& +a_5(\ln \Lambda/\mu)\epsilon_{\alpha\beta\gamma\delta}B_\alpha A_\beta \frac{1}{i}\partial_\gamma A_\delta \\
& +\bar{\psi}\left\{a_f(\ln \Lambda/\mu)\frac{1}{i}\not{\partial} + a_3(\ln \Lambda/\mu)\not{A} + a'_3(\ln \Lambda/\mu)\gamma_5\not{B}\right\}\psi\Bigg]. \quad (9.32)
\end{aligned}$$

Here the a''_4, a_5 terms are new features, not present in QED. Given A_μ is a vector, and B_μ is an axial vector, we find that the a''_4, a_5 terms are allowed by parity. (Note $\epsilon_{\alpha\beta\gamma\delta}A_\alpha B_\beta \partial_\gamma B_\delta$ and $B_\mu A_\mu A_\nu^2$ are forbidden by parity.)

We define the quantum BRST transformation by

$$\begin{aligned}
\delta A_\mu(k) &= k_\mu \epsilon c(k), & \delta B_\mu(k) &= k_\mu \epsilon' c'(k), \\
\delta c(k) &= 0, & \delta c'(k) &= 0, \\
\delta \bar{c}(-k) &= \epsilon \frac{1}{\xi} k_\mu [A_\mu]_\Lambda(-k), & \delta \bar{c}'(-k) &= \epsilon \frac{1}{\xi'} k_\mu [B_\mu]_\Lambda(-k), \\
\delta \psi(p) &= \int_k (e \epsilon c(k) - e' \epsilon' c'(k) \gamma_5) [\psi]_\Lambda(p-k), \\
\delta \bar{\psi}(-p) &= \int_k [\bar{\psi}]_\Lambda(-p-k) (-e \epsilon c(k) - e' \epsilon' c'(k) \gamma_5), \quad (9.33)
\end{aligned}$$

where the composite operators in the square brackets are defined as usual.

The corresponding WT composite operator is given by

$$\begin{aligned}
\Sigma_\Lambda &\equiv \int_k K(k/\Lambda) \left[\frac{\delta S_\Lambda}{\delta A_\mu(k)} \delta A_\mu(k) + \delta \bar{c}(-k) \frac{\overrightarrow{\delta}}{\delta \bar{c}(-k)} S_\Lambda \right. \\
&\quad \left. + \frac{\delta S_\Lambda}{\delta B_\mu(k)} \delta B_\mu(k) + \delta \bar{c}'(-k) \frac{\overrightarrow{\delta}}{\delta \bar{c}'(-k)} S_\Lambda \right] \\
&\quad + \int_p K(p/\Lambda) \left[S_\Lambda \frac{\overleftarrow{\delta}}{\delta \psi(p)} \delta \psi(p) - \text{Tr} \delta \psi(p) \frac{\overleftarrow{\delta}}{\delta \psi(p)} \right. \\
&\quad \left. + \delta \bar{\psi}(-p) \frac{\overrightarrow{\delta}}{\delta \bar{\psi}(-p)} S_\Lambda - \text{Tr} \frac{\overrightarrow{\delta}}{\delta \bar{\psi}(-p)} \delta \bar{\psi}(-p) \right]. \quad (9.34)
\end{aligned}$$

Substituting the above BRST transformation, we can rewrite this as

$$\Sigma_\Lambda = \epsilon \int_k c(k) (k_\mu J_\mu(-k) - \Phi(-k)) + \epsilon \int_k c'(k) (k_\mu J_{5,\mu}(-k) - \Phi_5(-k)), \quad (9.35)$$

where the composite operators are defined by

$$J_\mu(-k) \equiv \frac{\delta S_{I,\Lambda}}{\delta A_\mu(k)}, \quad J_{5,\mu}(-k) \equiv \frac{\delta S_{I,\Lambda}}{\delta B_\mu(k)}, \quad (9.36)$$

and

$$\Phi(-k) \equiv e \int_p K(p/\Lambda) \left[-S_\Lambda \frac{\overleftarrow{\delta}}{\delta \psi(p)} [\psi]_\Lambda(p-k) + \text{Tr} [\psi]_\Lambda(p-k) \frac{\overleftarrow{\delta}}{\delta \psi(p)} \right]$$

$$+e \int_p \text{Tr} U(-p-k, p) \frac{\vec{\delta}}{\delta \bar{\psi}(-p)} S_{I,\Lambda} \frac{\overleftarrow{\delta}}{\delta \psi(p+k)}, \quad (9.37)$$

$$\begin{aligned} \Phi_5(-k) \equiv e' \int_p K(p/\Lambda) & \left[S_\Lambda \frac{\overleftarrow{\delta}}{\delta \bar{\psi}(p)} \gamma_5 [\psi]_\Lambda(p-k) - \text{Tr} \gamma_5 [\psi]_\Lambda(p-k) \frac{\overleftarrow{\delta}}{\delta \bar{\psi}(p)} \right] \\ & + e' \int_p \text{Tr} U(-p-k, p) \gamma_5 \frac{\vec{\delta}}{\delta \bar{\psi}(-p)} S_{I,\Lambda} \frac{\overleftarrow{\delta}}{\delta \psi(p+k)}, \end{aligned} \quad (9.38)$$

where the matrix U is defined by (6.24) with $m = 0$:

$$U(-p-k, p) \equiv K((p+k)/\Lambda) \frac{1 - K(p/\Lambda)}{\not{p}} - \frac{1 - K((p+k)/\Lambda)}{\not{p} + \not{k}} K(p/\Lambda). \quad (9.39)$$

In the following we examine the asymptotic behavior of the WT composite operator Σ_Λ . We will restrict our discussion to the part relevant to chiral anomalies, i.e., the part proportional to the ϵ tensor. The asymptotic behavior (9.32) implies

$$\begin{cases} k_\mu J_\mu(-k) \xrightarrow{\Lambda \rightarrow \infty} -a_5 \epsilon_{\alpha\beta\gamma\delta} \int_{k_1+k_2=-k} k_{1\gamma} k_{2\delta} B_\alpha(k_1) A_\beta(k_2), \\ k_\mu J_{5,\mu}(-k) \xrightarrow{\Lambda \rightarrow \infty} +a_5 \epsilon_{\alpha\beta\gamma\delta} \int_{k_1+k_2=-k} k_{1\gamma} k_{2\delta} A_\alpha(k_1) A_\beta(k_2). \end{cases} \quad (9.40)$$

We have only one parameter $a_5(0)$ at our disposal.

On the other hand, assuming parity, the asymptotic behaviors (only the part proportional to the ϵ tensor) of Φ, Φ_5 are expected to be of the following form:

$$\begin{cases} \Phi(-k) \xrightarrow{\Lambda \rightarrow \infty} \epsilon_{\alpha\beta\gamma\delta} \int_{k_1+k_2=-k} k_{1\gamma} k_{2\delta} b B_\alpha(k_1) A_\beta(k_2), \\ \Phi_5(-k) \xrightarrow{\Lambda \rightarrow \infty} \epsilon_{\alpha\beta\gamma\delta} \int_{k_1+k_2=-k} k_{1\gamma} k_{2\delta} (b' A_\alpha(k_1) A_\beta(k_2) + b'' B_\alpha(k_1) B_\beta(k_2)). \end{cases} \quad (9.41)$$

Thus, for $\Sigma_\Lambda = 0$, we need to find

$$b'' = 0, \text{ and } b + b' = 0. \quad (9.42)$$

If this is the case, we can make Σ_Λ vanish by choosing

$$a_5 = -b. \quad (9.43)$$

Let us compute b, b', b'' at 1-loop. Only the second integrals of (9.37) and (9.38) contribute.

- (i) b — The BA term of Φ is given by the Feynman diagram in Fig. 25 and the one with B_α and A_β interchanged. The graph gives the integral

$$T_{\alpha\beta}(k_1, k_2) \equiv \int_p \text{Tr} eU(-p-k, p) e' \gamma_5 \gamma_\alpha \frac{1 - K((p-k_1)/\Lambda)}{\not{p} - \not{k}_1} e \gamma_\beta, \quad (9.44)$$

where U is defined by (9.39). Its $\Lambda \rightarrow \infty$ limit can be calculated as follows:

$$T_{\alpha\beta}(k_1, k_2) = e' e^2 \int_p U(-p+k_2, p+k_1) \gamma_5 \gamma_\alpha \frac{1 - K(p/\Lambda)}{\not{p}} \gamma_\beta$$




Fig. 25. A Feynman diagram contributing to $\Phi(-k)$. The broken line gives eU .

$$\begin{aligned}
&= e'e^2 \int_p \text{Tr} \left[\gamma_5 \gamma_\alpha \frac{1 - K(p/\Lambda)}{\not{p}} \gamma_\beta \cdot \left\{ K((p - k_2)/\Lambda) \frac{1 - K((p + k_1)/\Lambda)}{\not{p} + \not{k}_1} \right. \right. \\
&\quad \left. \left. - \frac{1 - K((p - k_2)/\Lambda)}{\not{p} - \not{k}_2} K((p + k_1)/\Lambda) \right\} \right] \\
&= e'e^2 \int_p \frac{1 - K(p/\Lambda)}{p^2} \left[\frac{K((p - k_2)/\Lambda) (1 - K((p + k_1)/\Lambda))}{(p + k_1)^2} \text{Tr} \gamma_5 \gamma_\alpha \not{p} \gamma_\beta \not{k}_1 \right. \\
&\quad \left. - \frac{K((p + k_1)/\Lambda) (1 - K((p - k_2)/\Lambda))}{(p - k_2)^2} \text{Tr} \gamma_5 \gamma_\alpha \not{p} \gamma_\beta (-\not{k}_2) \right] \\
&= 4e'e^2 \int_p \frac{1 - K(p/\Lambda)}{p^2} \left[\frac{K((p - k_2)/\Lambda) (1 - K((p + k_1)/\Lambda))}{(p + k_1)^2} \epsilon_{\alpha\gamma\beta\delta} p_\gamma k_{1\delta} \right. \\
&\quad \left. - \frac{K((p + k_1)/\Lambda) (1 - K((p - k_2)/\Lambda))}{(p - k_2)^2} \epsilon_{\alpha\gamma\beta\delta} p_\gamma (-k_{2\delta}) \right]. \quad (9.45)
\end{aligned}$$

Finally, using

$$\begin{aligned}
&\int_p \frac{(1 - K(p/\Lambda))^2}{p^4} K((p - k_2)/\Lambda) p_\gamma \\
&= \Lambda \int_p \frac{(1 - K(p))^2}{p^4} K\left(p - \frac{k_2}{\Lambda}\right) p_\gamma \\
&\xrightarrow{\Lambda \rightarrow \infty} \int_p \frac{(1 - K(p))^2}{p^4} \Delta(p) \frac{(pk_2)p_\gamma}{p^2} = \frac{1}{6(4\pi)^2} k_{2\gamma}, \quad (9.46)
\end{aligned}$$

we obtain


$$T_{\alpha\beta}(k_1, k_2) \xrightarrow{\Lambda \rightarrow \infty} e'e^2 \frac{4}{3} \frac{1}{(4\pi)^2} \epsilon_{\alpha\beta\gamma\delta} k_{1\gamma} k_{2\delta}. \quad (9.47)$$

The other graph gives the same contribution, and altogether we obtain

$$b = e'e^2 \frac{8}{3} \frac{1}{(4\pi)^2}. \quad (9.48)$$

- (ii) b' — The AA term of Φ_5 is given by the diagram in Fig. 26, similar to the previous diagram. The difference is in the use of $U\gamma_5$ instead of U , and replacement of B by A . There is one more graph obtained by interchanging $A_\alpha(k_1)$ and $A_\beta(k_2)$. The graph in Fig. 26 gives

$$\int_p \text{Tr} e'U(-p - k, p) \gamma_5 e \gamma_\alpha \frac{1 - K((p - k_1)/\Lambda)}{\not{p} - \not{k}_1} e \gamma_\beta$$


 Fig. 26. The broken line gives $e'U\gamma_5$.

$$= T_{\alpha\beta}(k_1, k_2) \xrightarrow{\Lambda \rightarrow \infty} e' e^2 \frac{4}{3} \frac{1}{(4\pi)^2} \epsilon_{\alpha\beta\gamma\delta} k_{1\gamma} k_{2\delta}. \quad (9.49)$$

The other graph gives the same. Hence, we obtain

$$b' = e' e^2 \frac{4}{3} \frac{1}{(4\pi)^2} = \frac{b}{2}. \quad (9.50)$$

(iii) b'' — The BB term of Φ_5 can be calculated similarly. We obtain

$$b'' = (e')^3 \frac{4}{3} \frac{1}{(4\pi)^2}. \quad (9.51)$$

Thus, we find

$$b'' \neq 0, \quad b + b' \neq 0. \quad (9.52)$$

No matter what we choose as a_5 , we cannot satisfy $\Sigma_A = 0$. At best, we can save the vector gauge invariance by choosing

$$a_5 = -b = -e' e^2 \frac{8}{3} \frac{1}{(4\pi)^2}. \quad (9.53)$$

This leaves us the following chiral anomaly:

$$\begin{aligned} \Sigma_A \xrightarrow{\Lambda \rightarrow \infty} & -\frac{1}{(4\pi)^2} \int_{k_1, k_2} \epsilon_{\alpha\beta\gamma\delta} k_{1\gamma} k_{2\delta} c'(-k_1 - k_2) \\ & \times \left(4e' e^2 A_\alpha(k_1) A_\beta(k_2) + \frac{4}{3} (e')^3 B_\alpha(k_1) B_\beta(k_2) \right). \end{aligned} \quad (9.54)$$

9.2.2. $U(1)_R$ gauge theory

We construct a quantum theory whose classical action is given by

$$S_{cl} = \int d^4x \left[-\frac{1}{4} F_{\mu\nu}^2 + \frac{1}{2\xi} (\partial_\mu A_\mu)^2 - \partial_\mu \bar{c} \partial_\mu c + \bar{\psi}_R \left(-\frac{1}{i} \not{\partial} + e \not{A} \right) \psi_R \right], \quad (9.55)$$

where ψ_R and $\bar{\psi}_R$ are chiral:

$$\frac{1 + \gamma_5}{2} \psi_R = \psi_R, \quad \bar{\psi}_R \frac{1 - \gamma_5}{2} = \bar{\psi}_R. \quad (9.56)$$

For the quantum theory, the WT composite operator is given by

$$\Sigma_A = \epsilon \int_k c(k) \left(k_\mu \frac{\delta S_{I,A}}{\delta A_\mu(k)} - \Phi_R(-k) \right), \quad (9.57)$$

where

$$\begin{aligned} \Phi_R(-k) \equiv e \int_p K(p/\Lambda) & \left[-S_\Lambda \frac{\overleftarrow{\delta}}{\delta\psi_R(p)} [\psi_R]_\Lambda(p-k) + [\bar{\psi}_R]_\Lambda(-p-k) \frac{\overrightarrow{\delta}}{\delta\bar{\psi}(-p)} S_\Lambda \right. \\ & \left. + \text{Tr} \frac{1+\gamma_5}{2} U(-p-k, p) \frac{\overrightarrow{\delta}}{\delta\bar{\psi}_R(-p)} S_{I,\Lambda} \frac{\overleftarrow{\delta}}{\delta\psi_R(p+k)} \right], \end{aligned} \quad (9.58)$$

where U is defined by (9.39). Φ_R has the asymptotic behavior

$$\begin{aligned} \Phi_R(-k) & \xrightarrow{\Lambda \rightarrow \infty} (\Lambda^2 \bar{a}_2(\ln \Lambda/\mu) + k^2 \bar{d}_2(\ln \Lambda/\mu)) k_\mu A_\mu(-k) \\ & + \bar{a}_3(\ln \Lambda/\mu) \int_p \bar{\psi}_R(-p-k) \not{k} \psi_R(p) \\ & + b(\ln \Lambda/\mu) \epsilon_{\alpha\beta\gamma\delta} \int_{k_1+k_2=-k} k_{1\alpha} k_{2\beta} A_\gamma(k_1) A_\delta(k_2) \\ & + \bar{a}_4(\ln \Lambda/\mu) \frac{1}{2} \int_{k_1, k_2} A_\nu(k_1) A_\nu(k_2) k_\mu A_\mu(-k-k_1-k_2). \end{aligned} \quad (9.59)$$

Compared with Φ for QED, Φ_R has an extra term proportional to the ϵ tensor. On the other hand, the asymptotic behavior of $S_{I,\Lambda}$ has the same form as for QED:

$$\begin{aligned} S_{I,\Lambda} & \xrightarrow{\Lambda \rightarrow \infty} \int d^4x \left[\Lambda^2 a_2(\ln \Lambda/\mu) \frac{1}{2} A_\mu^2 \right. \\ & + c_2(\ln \Lambda/\mu) \frac{1}{2} \partial_\mu A_\nu \partial_\mu A_\nu + d_2(\ln \Lambda/\mu) \frac{1}{2} (\partial_\mu A_\mu)^2 \\ & + a_4(\ln \Lambda/\mu) \frac{1}{4!} (A_\mu^2)^2 \\ & \left. + a_f(\ln \Lambda/\mu) \bar{\psi}_R \frac{1}{i} \not{\partial} \psi_R + a_3(\ln \Lambda/\mu) \bar{\psi}_R \not{A} \psi_R \right], \end{aligned} \quad (9.60)$$

which implies

$$\begin{aligned} k_\mu \frac{\delta S_{I,\Lambda}}{\delta A_\mu(k)} & \xrightarrow{\Lambda \rightarrow \infty} (\Lambda^2 a_2(\ln \Lambda/\mu) + k^2 (c_2 + d_2)(\ln \Lambda/\mu)) k_\mu A_\mu(-k) \\ & + a_3(\ln \Lambda/\mu) \int_p \bar{\psi}_R(-p-k) \not{k} \psi_R(p) \\ & + a_4(\ln \Lambda/\mu) \frac{1}{2} \int_{k_1, k_2} A_\nu(k_1) A_\nu(k_2) k_\mu A_\mu(-k-k_1-k_2). \end{aligned} \quad (9.61)$$

Hence, unless

$$b(0) = 0 \quad (9.62)$$

by chance, we cannot satisfy the Ward identity by adjusting the parameters in hands.

At 1-loop, the coefficient b can be calculated from Fig. 25 where B_α is replaced by A_α , and U is multiplied by the projection operator $\frac{1+\gamma_5}{2}$ from the left. We obtain

$$\begin{aligned} & \int_p \text{Tr} \frac{1+\gamma_5}{2} e U(-p-k, p) e \gamma_\alpha \frac{1-K((p-k_1)/\Lambda)}{\not{p}-\not{k}_1} e \gamma_\beta \\ & = -\frac{1}{2} T_{\alpha\beta}(k_1, k_2) \Big|_{e'=e} \xrightarrow{\Lambda \rightarrow \infty} -e^3 \frac{2}{3} \frac{1}{(4\pi)^2} \epsilon_{\alpha\beta\gamma\delta} k_{1\gamma} k_{2\delta}, \end{aligned} \quad (9.63)$$

where e' is replaced by e . There is another graph with $A_\alpha(k_1)$ and $A_\beta(k_2)$ interchanged. Hence, we obtain

$$b(0) = -e^3 \frac{2}{3} \frac{1}{(4\pi)^2}. \quad (9.64)$$

Therefore, we cannot satisfy the WT identity.

§10. Functional integral approach to the antifield formalism

In §§7 & 8 we have adapted the antifield formalism to the ERG framework. Though our discussion has been limited to perturbation theory, we have obtained an important conclusion: realization of symmetry is reduced to solving a classical algebraic problem. The purpose of the present section is to give a second look at the adaptation of the antifield formalism without relying on perturbative expansions.

The antifield formalism of Batalin and Vilkovisky¹⁶⁾⁴⁹⁾⁵¹⁾ can provide us with a systematic description of any symmetry present in a theory. In the following we apply the antifield formalism to the bare and Wilson actions. The quantum master equation (QME), satisfied by an action, signals the presence of symmetry, naturally extending the WT identity discussed in §5. As opposed to §§7 & 8, where we have constructed the continuum limit directly, we start from a bare action at a UV scale Λ_0 . Following the procedure of §2 we construct a Wilson action by integrating the fields over the momenta between Λ_0 and Λ ($< \Lambda_0$). With Λ_0 kept finite, the resulting ERG trajectory approaches the trajectory of the continuum limit $\bar{S}_\Lambda$ only as $\Lambda \rightarrow 0$. (See Fig. 16.) Nevertheless, by keeping Λ_0 finite, we can manage to write down more explicit expressions for the antifield dependence of the Wilson action,^{53),65),67)} compared with the perturbative construction in terms of the asymptotic behavior in §§7 & 8.

We discuss only a generic gauge theory in this section, and give concrete examples in the next section. A global symmetry can be treated in a similar manner,⁶⁴⁾⁵⁹⁾ though we do not discuss it here.^{*)} The organization of this section is as follows. In §10.1, we first review the classical antifield formalism, and then explain the modification necessary for quantized systems. For classical systems a symmetry is realized as a classical master equation (CME), and for quantum systems as a quantum master equation (QME). We then show, in §10.2, how to adapt the formalism in the presence of a UV cutoff, and in §10.3, how the QME relates to the WT identity. In §10.4 we derive explicit antifield dependence of the Wilson action, assuming a simple form for the BRST transformation of the bare action. In §10.5 we introduce the effective average action, and rewrite the QME for the Wilson action as the so-called “modified Slavnov-Taylor identity”⁴⁰⁾ for the average action. Finally, in §10.6 we discuss an implication of the nilpotency of the BRST transformation.

In this section, we adopt a notation that can handle bosonic and fermionic fields equally in order to respect the canonical structure of the antifield formalism. We use the symbol ϕ^A with a suffix A for fields, and distinguish their statistics by Grassmann parity $\epsilon_A \bmod 2$:

$$\epsilon(\phi^A) = \epsilon_A \equiv \begin{cases} 0 & \text{bosonic,} \\ 1 & \text{fermionic.} \end{cases} \quad (\bmod 2) \quad (10.1)$$

Though not mentioned explicitly in §§7 & 8, a field and its corresponding antifield form a canonical pair. In order to keep the canonical structure, we rescale an antifield

^{*)} The QME for chiral symmetry is an extension of the well-known Ginsparg-Wilson relation.⁴⁷⁾ In §6.5 we have briefly discussed the WT identity for the axial symmetry.

by the cutoff function $K(p/\Lambda)$. Hence, we must replace the antifield $\phi^*(p)$ of the previous sections by $K(p/\Lambda)\phi^*(p)$ to get the corresponding results in this section.

10.1. The Batalin-Vilkovisky antifield formalism

Let us consider a classical gauge fixed action $S_{cl}[\phi]$ that is invariant under the nilpotent BRST transformation $\delta\phi^{A,*}$

$$\delta S_{cl} \equiv \frac{\partial^r S_{cl}}{\partial \phi^A} \delta \phi^A = 0. \quad (10.2)$$

The fields ϕ^A represent collectively gauge and matter fields as well as (anti)ghosts and B-fields.***) The index A represents any index to distinguish different types of fields, such as momentum, the Lorentz index for a vector field, and the spinor index for a spinor field. Hence, the Einstein convention for repeated A includes integration over momenta.

By introducing an antifield ϕ_A^* for each field ϕ^A , we define an extended action

$$\bar{S}_{cl}[\phi, \phi^*] \equiv S_{cl}[\phi] + \phi_A^* \delta \phi^A. \quad (10.3)$$

The antifield ϕ_A^* has the opposite Grassmann parity to that of ϕ^A :

$$\epsilon(\phi_A^*) \equiv \epsilon(\phi^A) + 1 \pmod{2} \quad (10.4)$$

We also assign ghost numbers to ϕ^A and ϕ_A^* so that they add up to -1 .

In the space of ϕ^A and ϕ_A^* , we define a canonical structure by introducing an antibracket: for any field variables X and Y , we define

$$(X, Y) \equiv \frac{\partial^r X}{\partial \phi^A} \frac{\partial^l Y}{\partial \phi_A^*} - \frac{\partial^r X}{\partial \phi_A^*} \frac{\partial^l Y}{\partial \phi^A}. \quad (10.5)$$

Fields ϕ^A and their antifields ϕ_A^* are canonical conjugate pairs, satisfying

$$(\phi^A, \phi_B^*) = \delta_B^A, \quad (\phi^A, \phi^B) = (\phi_A^*, \phi_B^*) = 0. \quad (10.6)$$

We now define the BRST transformation of an arbitrary variable X by

$$\delta X \equiv (X, \bar{S}_{cl}). \quad (10.7)$$

This generalizes $\delta\phi^A$ for the fields ϕ^A ; for example, it gives the transformation of antifields by

$$\delta \phi_A^* = -\frac{\partial^l \bar{S}_{cl}}{\partial \phi^A}. \quad (10.8)$$

Following the definitions (10.3), (10.5), (10.7), we obtain

$$(\bar{S}_{cl}, \bar{S}_{cl}) = 2(\delta S_{cl} + \phi_A^* \delta^2 \phi^A), \quad (10.9)$$

*) The right (left) derivative is indicated by the superscript r (l).

**) The B field is an auxiliary field which is equal to the divergence of the gauge field $\partial_\mu A_\mu$ by the equation of motion. We introduce this to avoid the quadratic term $(\bar{c}^*)^2$ in $\bar{S}_{cl}$ (10.3).

where δS_{cl} is defined by (10·2). Hence, if $\delta S_{cl} = 0$ (10·2), and the original BRST transformation is nilpotent

$$\delta^2 \phi^A = 0, \quad (10\cdot10)$$

we obtain the **classical master equation (CME)**:

$$(\bar{S}_{cl}, \bar{S}_{cl}) = 0. \quad (10\cdot11)$$

Thus, CME is equivalent to the combination of the BRST invariance of S_{cl} (10·2) and the nilpotency (10·10). In fact the nilpotency of the BRST transformation (10·7) remains valid, even if δ acts on an arbitrary variable X , dependent on both ϕ^A and ϕ_A^* ; applying δ twice, we obtain

$$\delta^2 X = (\delta X, \bar{S}_{cl}) = ((X, \bar{S}_{cl}), \bar{S}_{cl}) = (X, (\bar{S}_{cl}, \bar{S}_{cl})). \quad (10\cdot12)$$

Hence, (10·11) implies the general nilpotency:

$$\delta^2 X = 0. \quad (10\cdot13)$$

We now apply the BV formalism to quantum systems with BRST invariance. Let $\bar{S}[\phi, \phi^*]$ be an action that defines a quantum system via functional integration over ϕ : $\bar{S}$ is either a bare action with a UV cutoff Λ_0 , or a Wilson action with a lower cutoff Λ . Under the BRST transformation of fields

$$\delta \phi^A = \frac{\partial^l \bar{S}}{\partial \phi_A^*}, \quad (10\cdot14)$$

the action changes by the **quantum master operator**:

$$\bar{\Sigma}[\phi, \phi^*] \equiv \frac{\partial^r \bar{S}}{\partial \phi^A} \frac{\partial^l \bar{S}}{\partial \phi_A^*} + \frac{\partial^r}{\partial \phi^A} \delta \phi^A = \frac{1}{2}(\bar{S}, \bar{S}) + \Delta \bar{S}, \quad (10\cdot15)$$

where we define

$$\Delta \equiv (-)^{\epsilon_A+1} \frac{\partial^r}{\partial \phi^A} \frac{\partial^r}{\partial \phi_A^*}. \quad (10\cdot16)$$

The first term is the classical change of the action under the BRST transformation (10·14). The second term is the contribution from the jacobian of the BRST transformation. The system is BRST invariant quantum mechanically if the two contributions cancel:

$$\bar{\Sigma}[\phi, \phi^*] = 0. \quad (10\cdot17)$$

We call this equation the **quantum master equation (QME)**. Note that the quantum master operator (10·15) appears somewhat different from (7·2); the cutoff dependent factor $K(p/\Lambda)$ is missing in (10·15). This is due to the adoption of a different normalization of the antifields, as mentioned in the last paragraph before §10.1.

In the antifield formalism, we define the quantum BRST transformation as

$$\delta_Q X \equiv (X, \bar{S}) + \Delta X \quad (10\cdot18)$$

for an arbitrary variable X . Without assuming QME, we obtain two important identities:

$$\delta_Q \bar{\Sigma}[\phi, \phi^*] = 0, \quad (10\cdot19)$$

$$\delta_Q^2 X = (X, \bar{\Sigma}[\phi, \phi^*]). \quad (10\cdot20)$$

These identities are algebraic, consequences of the definitions of the quantum master operator (10·15) and the quantum BRST transformation (10·18). The identity (10·19) is crucial for the perturbative construction of symmetric theories, as shown in §§7 & 8. Eq. (10·20) implies that the quantum BRST transformation (10·18) is nilpotent if and only if QME (10·17) holds. Note the importance of the second term in (10·18). If we defined the BRST transformation without it by

$$\delta X \equiv (X, \bar{S}), \quad (10\cdot21)$$

we would obtain

$$\delta^2 X = (X, (\bar{S}, \bar{S})), \quad (10\cdot22)$$

and hence nilpotency demands CME instead of QME. For a generic quantum system, however, what holds is QME, but not CME. Therefore, it is the quantum BRST transformation that can be nilpotent.

10.2. Scale change in the functional integral approach

The action $\bar{S}$ of the previous subsection is either a bare action $\bar{S}_B$ or the corresponding Wilson action $\bar{S}_A$.*) We recapitulate the important results of §2 using the notation adopted in the present section.

Let $\bar{S}_B$ be an action defined at the UV scale Λ_0 in the presence of antifields:

$$\bar{S}_B[\phi, \phi^*] \equiv S_B[\phi] + \phi^* \cdot \delta\phi, \quad (10\cdot23)$$

where S_B is the sum of the gaussian and interaction terms:

$$S_B[\phi] \equiv -\frac{1}{2}\phi \cdot K_0^{-1} D \cdot \phi + S_{I,B}[\phi]. \quad (10\cdot24)$$

Here, we have adopted the matrix notation for momentum integrals:

$$\phi \cdot K_0^{-1} D \cdot \phi \equiv \int_p \phi^A(-p) \frac{1}{K_0(p)} D_{AB}(p) \phi^B(p), \quad (10\cdot25)$$

$$\phi^* \cdot \delta\phi \equiv \int_p \phi_A^*(-p) \phi^A(p). \quad (10\cdot26)$$

We have assumed that the antifield dependence of $\bar{S}_B$ is strictly linear.

By introducing sources J_A , we define the generating functional

$$\bar{Z}_B[J, \phi^*] \equiv \int \mathcal{D}\phi \exp(\bar{S}_B[\phi, \phi^*] + K_0^{-1} J \cdot \phi). \quad (10\cdot27)$$

*) As in §2 we denote a bare action by S_B and a Wilson action by S_A . As in §§7 & 8 we also put a bar to denote the dependence on antifields.

In performing the above functional integral, we decompose the fields ϕ^A into the IR fields Φ^A and UV fields χ^A , where the propagators of each class of fields are given by

$$\begin{aligned} & K_0(p) (D_{AB}(p))^{-1} && \text{for } \phi^A, \\ & K(p) (D_{AB}(p))^{-1} && \text{for } \Phi^A, \\ & (K_0(p) - K(p)) (D_{AB}(p))^{-1} && \text{for } \chi^A. \end{aligned} \quad (10\cdot28)$$

Note that Φ^A carry the momenta below Λ , and χ^A those between Λ_0 and Λ .

We now introduce the Wilson action⁶⁵⁾⁶⁵⁾

$$\bar{S}_\Lambda[\Phi, \Phi^*] \equiv -\frac{1}{2} \Phi \cdot K^{-1} D \cdot \Phi + \bar{S}_{I,\Lambda}[\Phi, \Phi^*], \quad (10\cdot29)$$

where the interaction part is defined by the functional integral over the UV fields χ^A :

$$\exp(\bar{S}_{I,\Lambda}[\Phi, \Phi^*]) \equiv \int \mathcal{D}\chi \exp \left[-\frac{1}{2} \chi \cdot (K_0 - K)^{-1} D \cdot \chi + \bar{S}_{I,B}[\Phi + \chi, \phi^*] \right]. \quad (10\cdot30)$$

The two antifields ϕ^*, Φ^* are related simply as

$$K \Phi_A^* = K_0 \phi_A^*. \quad (10\cdot31)$$

We define the generating functional of the Wilson action by

$$\bar{Z}_\Lambda[J, \Phi^*] = \int \mathcal{D}\Phi \exp(\bar{S}_\Lambda[\Phi, \Phi^*] + K^{-1} J \cdot \Phi). \quad (10\cdot32)$$

The two generating functionals (10\cdot27) and (10\cdot32) are related by (2\cdot29), which is rewritten as

$$\bar{Z}_B[J, \phi^*] = N_J \bar{Z}_\Lambda[J, \Phi^*], \quad (10\cdot33)$$

where the normalization factor N_J is given by

$$\ln N_J = -\frac{(-)^{\epsilon_A}}{2} J_A K_0^{-1} K^{-1} (K_0 - K) (D^{-1})^{AB} J_B. \quad (10\cdot34)$$

Because of its importance, we wish to rederive it using the present notation. The main tool is the triviality of the gaussian integral:

$$\begin{aligned} & \int \mathcal{D}\theta \exp \left[-\frac{1}{2} \left(\theta - J (K_0 - K) D^{-1} \right) \cdot \frac{D}{K_0 K (K_0 - K)} \right. \\ & \quad \left. \cdot \left(\theta - (-)^{\epsilon(J)} D^{-1} (K_0 - K) J \right) \right] = \text{const}. \end{aligned} \quad (10\cdot35)$$

We substitute this into the partition function $\bar{Z}_B[J, \phi^*]$ defined by (10\cdot27), and perform a canonical change of variables from $\{\phi, \phi^*, \theta, \theta^*\}$ to $\{\Phi, \Phi^*, \chi, \chi^*\}$, where

$$\begin{aligned} \phi^A &= \Phi^A + \chi^A, & \theta^A &= (K_0 - K) \Phi^A - K \chi^A, \\ \phi_A^* &= K_0^{-1} [K \Phi_A^* + (K_0 - K) \chi_A^*], & \theta_A^* &= K_0^{-1} (\Phi_A^* - \chi_A^*). \end{aligned} \quad (10\cdot36)$$

Then, we obtain

$$\begin{aligned} \bar{Z}_B[J, \phi^*] = N_J \int \mathcal{D}\Phi \mathcal{D}\chi \exp \left[-\frac{1}{2} \Phi \cdot K^{-1} D \cdot \Phi + K^{-1} J \cdot \Phi \right. \\ \left. -\frac{1}{2} \chi \cdot (K_0 - K)^{-1} D \cdot \chi + \bar{S}_{I,B}[\Phi + \chi, \phi^*] \right], \end{aligned} \quad (10.37)$$

where N_J is given by (10.34). This is independent of the antifields χ^* , and we may adopt a gauge choice

$$\chi_A^* = 0. \quad (10.38)$$

This gives

$$K_0 \phi_A^* = K \Phi_A^*. \quad (10.39)$$

We will justify this relation shortly. With the above gauge choice, we obtain (10.33) & (10.34), where the Wilson action is defined by (10.29) & (10.30).

The partition function $\bar{Z}_B$ does not depend on the IR cutoff Λ so that

$$\Lambda \frac{\partial}{\partial \Lambda} \bar{Z}_B = 0. \quad (10.40)$$

This yields the Polchinski differential equation⁹⁴⁾

$$\begin{aligned} -\Lambda \frac{\partial}{\partial \Lambda} \bar{S}_\Lambda = \int_p (K^{-1} \Delta)(p) \left[\Phi^A(p) \frac{\partial^l \bar{S}_\Lambda}{\partial \Phi^A(p)} - \Phi_A^*(p) \frac{\partial^l \bar{S}_\Lambda}{\partial \Phi_A^*(p)} \right] \\ + \frac{1}{2} \int_p (-)^{\epsilon_A} (D^{-1} \Delta)^{AB}(p) \left[\frac{\partial^l \bar{S}_\Lambda}{\partial \Phi^B(-p)} \frac{\partial^r \bar{S}_\Lambda}{\partial \Phi^A(p)} + \frac{\partial^l \partial^r \bar{S}_\Lambda}{\partial \Phi^B(-p) \partial \Phi^A(p)} \right], \end{aligned} \quad (10.41)$$

where we define

$$\Delta(p/\Lambda) \equiv \Lambda \frac{\partial}{\partial \Lambda} K(p/\Lambda) \quad (10.42)$$

as in (2.47). $\bar{S}_\Lambda$ is completely determined by the Polchinski differential equation and the initial condition

$$\bar{S}_\Lambda[\Phi, \Phi^*] \Big|_{\Lambda=\Lambda_0} = \bar{S}_B[\Phi, \Phi^*]. \quad (10.43)$$

We can define a composite operator $\bar{\mathcal{O}}_\Lambda[\Phi, \Phi^*]$ as a functional that satisfies the differential equation (4.5). Using the present notation, we obtain

$$-\Lambda \frac{\partial}{\partial \Lambda} \bar{\mathcal{O}}_\Lambda = \bar{\mathcal{D}} \cdot \bar{\mathcal{O}}_\Lambda, \quad (10.44)$$

where

$$\bar{\mathcal{D}} \equiv -(K^{-1} \Delta) \Phi_A^* \frac{\partial^l}{\partial \Phi_A^*} + (-)^{\epsilon_A} (D^{-1} \Delta)^{AB} \left(\frac{\partial^l \bar{S}_{I,A}}{\partial \Phi^B} \frac{\partial^r}{\partial \Phi^A} + \frac{1}{2} \frac{\partial^l \partial^r}{\partial \Phi^B \partial \Phi^A} \right). \quad (10.45)$$

Note that the first term of (10·45), rescaling of the antifield, is a consequence of (10·39). Hence, the factor K^{-1} is necessary in order to make

$$K^{-1} \frac{\partial \bar{S}_\Lambda}{\partial \Phi_A^*} \quad (10·46)$$

composite operators.

For later convenience, we recall an alternative definition of composite operators. Given an arbitrary functional $\bar{\mathcal{O}}_B[\phi, \phi^*]$ at the UV scale, the corresponding IR composite operator $\bar{\mathcal{O}}_\Lambda[\Phi, \Phi^*]$ is constructed as

$$\begin{aligned} \bar{\mathcal{O}}_\Lambda[\Phi, \Phi^*] \exp(\bar{S}_{I,\Lambda}[\Phi, \Phi^*; \Lambda]) &\equiv \int \mathcal{D}\chi \bar{\mathcal{O}}_B[\Phi + \chi, \phi^*] \\ &\cdot \exp \left[-\frac{1}{2} \chi \cdot (K_0 - K)^{-1} D \cdot \chi + \bar{S}_{I,B}[\Phi + \chi, \phi^*] \right]. \end{aligned} \quad (10·47)$$

This implies

$$\langle \bar{\mathcal{O}}_\Lambda[\Phi, \Phi^*] \rangle_{\bar{S}_\Lambda, K^{-1}J} = N_J^{-1} \langle \bar{\mathcal{O}}_B[\phi, \phi^*] \rangle_{\bar{S}_B, K_0^{-1}J} \quad (10·48)$$

in the presence of arbitrary sources J coupled linearly to the fields.

10.3. WT identity and QME

In this subsection we consider the cutoff dependence of the quantum master operator. Our starting point is the BRST transformation of fields at the UV scale Λ_0 , given in terms of an anticommuting constant λ as

$$\phi^A \rightarrow \phi'^A = \phi^A + \delta_\lambda \phi^A, \quad \delta_\lambda \phi^A = \delta \phi^A \lambda = K_0 \mathcal{R}^A[\phi] \lambda. \quad (10·49)$$

In computing the generating functional (10·27), we consider changing integration variables from ϕ to ϕ' . If the jacobian is properly taken into account, the integral does not depend on the choice of integration variables. Hence, we obtain

$$\begin{aligned} \int \mathcal{D}\phi \left(K_0^{-1} J \cdot \delta \phi + \Sigma_B[\phi] + \frac{\partial^r (\phi^* \cdot \delta \phi)}{\partial \phi^A} \delta \phi^A \right) \\ \times \exp(\bar{S}_B[\phi, \phi^*] + K_0^{-1} J \cdot \phi) = 0, \end{aligned} \quad (10·50)$$

where $\Sigma_B[\phi]$ is the WT operator defined by

$$\Sigma_B[\phi] \equiv \frac{\partial^r S_B}{\partial \phi^A} \delta \phi^A + \frac{\partial^r}{\partial \phi^A} \delta \phi^A. \quad (10·51)$$

$\Sigma_B[\phi]$ is the sum of the change of the original gauge fixed action $S_B[\phi]$

$$\delta_\lambda S_B = \frac{\partial^r S_B}{\partial \phi^A} \delta_\lambda \phi^A, \quad (10·52)$$

and the contribution from the jacobian

$$\delta_\lambda \ln \mathcal{D}\phi = (-)^{\epsilon_A} \frac{\partial^r}{\partial \phi^A} \delta_\lambda \phi^A = \frac{\partial^r}{\partial \phi^A} \delta \phi^A \lambda. \quad (10·53)$$

The quantum master operator defined by^{*)}

$$\bar{\Sigma}_B = \frac{1}{2}(\bar{S}_B, \bar{S}_B) + \Delta \bar{S}_B \quad (10.54)$$

is indeed given as the sum of the WT operator and the third term of (10.50):

$$\bar{\Sigma}_B[\phi, \phi^*] = \frac{\partial^r \bar{S}_B}{\partial \phi^A} \delta \phi^A + \frac{\partial^r}{\partial \phi^A} \delta \phi^A = \Sigma_B[\phi] + \frac{\partial^r (\phi^* \cdot \delta \phi)}{\partial \phi^A} \delta \phi^A, \quad (10.55)$$

where we have used $\delta \phi^A = \partial^l \bar{S}_B / \partial \phi_A^*$. Hence, we obtain a simple relation between the QM and WT operators:

$$\bar{\Sigma}_B = \Sigma_B + \phi^* \cdot \delta^2 \phi. \quad (10.56)$$

As CME implies both the BRST invariance of the action and the nilpotency of the BRST transformation, QME also implies both. In contrast, the WT identity implies only the invariance of the action.

We now rewrite (10.50) as

$$\langle \bar{\Sigma}_B[\phi, \phi^*] \rangle_{\bar{S}_B, K_0^{-1}J} = -K_0^{-1}J \cdot \langle \delta \phi \rangle_{\bar{S}_B, K_0^{-1}J}. \quad (10.57)$$

Setting $\phi^* = 0$, we obtain an analogous relation for the WT operator:

$$\langle \Sigma_B[\phi] \rangle_{S_B, K_0^{-1}J} = -K_0^{-1}J \cdot \langle \delta \phi \rangle_{S_B, K_0^{-1}J}. \quad (10.58)$$

Using (10.49), we obtain

$$K_0^{-1} \langle \delta \phi \rangle_{\bar{S}_B, K_0^{-1}J} = \langle \mathcal{R}[\phi] \rangle_{\bar{S}_B, K_0^{-1}J} = \mathcal{R} \left[K_0 \partial_J^l \right] \bar{Z}_B[J, \phi^*], \quad (10.59)$$

where the differential operator $\mathcal{R}^A[K_0 \partial_J^l]$ is called the Slavnov operator. Setting $\phi^* = 0$ we obtain

$$K_0^{-1} \langle \delta \phi \rangle_{S_B, K_0^{-1}J} = \mathcal{R} \left[K_0 \partial_J^l \right] Z_B[J]. \quad (10.60)$$

Therefore, we can rewrite (10.57) and (10.58) as

$$\langle \bar{\Sigma}_B[\phi, \phi^*] \rangle_{\bar{S}_B, K_0^{-1}J} = -J \cdot \mathcal{R} \left[K_0 \partial_J^l \right] \bar{Z}_B[J, \phi^*], \quad (10.61)$$

$$\langle \Sigma_B[\phi] \rangle_{S_B, K_0^{-1}J} = -J \cdot \mathcal{R} \left[K_0 \partial_J^l \right] Z_B[J]. \quad (10.62)$$

We note that the above relations, being based upon the functional relation (10.50), are valid whether or not the bare action is invariant under the BRST transformation (10.49).

We now wish to transform (10.61) into an equivalent condition on the Wilson action $\bar{S}_A$. We first apply (10.47) to define the QM operator $\bar{\Sigma}_A[\Phi, \Phi^*]$ for the IR theory by

$$\begin{aligned} \bar{\Sigma}_A[\Phi, \Phi^*] \exp(\bar{S}_{I,A}[\Phi, \Phi^*]) &\equiv \int \mathcal{D}\chi \bar{\Sigma}_B[\Phi + \chi, \phi^*] \\ &\times \exp \left[-\frac{1}{2} \chi \cdot (K_0 - K)^{-1} D \cdot \chi + \bar{S}_{I,B}[\Phi + \chi, \phi^*] \right]. \end{aligned} \quad (10.63)$$

^{*)} Note that Δ here is (10.16), but not the log derivative of K .

Then, (10·48) implies

$$\langle \bar{\Sigma}_\Lambda[\Phi, \Phi^*] \rangle_{\bar{S}_\Lambda, K^{-1}J} = N_J^{-1} \langle \bar{\Sigma}_B[\phi, \phi^*] \rangle_{\bar{S}_B, K_0^{-1}J} . \quad (10\cdot64)$$

We also define the IR operator $\delta\Phi^A$ corresponding to $\delta\phi^A$ by

$$\begin{aligned} & K^{-1}\delta\Phi^A[\Phi, \Phi^*] \exp(\bar{S}_{I,\Lambda}[\Phi, \Phi^*]) \\ & \equiv \int \mathcal{D}\chi K_0^{-1}\delta\phi^A[\Phi + \chi] \exp\left[-\frac{1}{2}\chi \cdot (K_0 - K)^{-1}D \cdot \chi + \bar{S}_{I,B}[\Phi + \chi, \phi^*]\right] . \end{aligned} \quad (10\cdot65)$$

Eq. (10·48) implies

$$\langle K^{-1}\delta\Phi^A \rangle_{\bar{S}_\Lambda, K^{-1}J} = N_J^{-1} \langle K_0^{-1}\delta\phi^A \rangle_{\bar{S}_B, K_0^{-1}J} . \quad (10\cdot66)$$

Denoting

$$R^A[\Phi, \Phi^*] \equiv K^{-1}\delta\Phi^A[\Phi, \Phi^*] , \quad (10\cdot67)$$

we can rewrite (10·66) as

$$R^A[K\partial_J^l, \Phi^*] \bar{Z}_\Lambda[J, \Phi^*] = N_J^{-1} \mathcal{R}^A[K_0\partial_J^l] \bar{Z}_B[J, \phi^*] . \quad (10\cdot68)$$

Hence, using (10·64) and (10·68), we can transform (10·61) into

$$\langle \bar{\Sigma}_\Lambda[\Phi, \Phi^*] \rangle_{\bar{S}_\Lambda, K^{-1}J} = -J \cdot R[K\partial_J^l, \Phi^*] \bar{Z}_\Lambda[J, \Phi^*] , \quad (10\cdot69)$$

which is the desired relation.

The above relation implies

$$\bar{\Sigma}_\Lambda[\Phi, \Phi^*] = \frac{\partial^r \bar{S}_\Lambda[\Phi, \Phi^*]}{\partial \Phi^A} \delta\Phi^A + \frac{\partial^r}{\partial \Phi^A} \delta\Phi^A . \quad (10\cdot70)$$

This has the form expected for the quantum master operator; here the relevant action is the Wilson action $\bar{S}_\Lambda$ defined at an IR scale Λ , and the BRST transformation $\delta\Phi^A$ of the IR field is a composite operator if multiplied by K^{-1} .

Similarly, we can transform (10·62) into

$$\langle \Sigma_\Lambda[\Phi] \rangle_{S_\Lambda, K^{-1}J} = -J \cdot R[K\partial_J^l] Z_\Lambda[J] , \quad (10\cdot71)$$

where the WT operator and the BRST transformation are obtained by setting Φ^* to zero:

$$\Sigma_\Lambda[\Phi] \equiv \bar{\Sigma}_\Lambda[\Phi, \Phi^*] \big|_{\Phi^*=0} = \frac{\partial^r S_\Lambda[\Phi]}{\partial \Phi^A} \delta\Phi^A[\Phi] + \frac{\partial^r}{\partial \Phi^A} \delta\Phi^A[\Phi] , \quad (10\cdot72)$$

$$\delta\Phi^A[\Phi] \equiv \delta\Phi^A[\Phi, \Phi^*] \big|_{\Phi^*=0} = K R^A[\Phi] . \quad (10\cdot73)$$

So far our discussion has been kept very general; we have only assumed that the bare action $\bar{S}_B$ takes the form of (10·23) & (10·24). For symmetric theories the QME must hold: $\bar{\Sigma}_B$ and hence $\bar{\Sigma}_\Lambda$ must vanish. For renormalizable theories, the QME need not hold for any finite Λ_0 , but it must hold in the limit $\Lambda_0 \rightarrow \infty$. In general it is a difficult problem to construct a bare action that satisfies the QME exactly;

in §§7 & 8 we have discussed this problem directly in the limit $\Lambda_0 \rightarrow \infty$, but only perturbatively. For anomaly-free renormalizable theories, we can expect that the quantum master operator behaves as

$$\bar{\Sigma}_B[\phi, \phi^*] = O(1/\Lambda_0),$$

for a large but finite value of Λ_0 .^{*)} To be more precise, we expect

$$\langle \bar{\Sigma}_B[\phi, \phi^*] \rangle_{\bar{S}_B, K_0^{-1}J} = O(1/\Lambda_0), \quad (10.74)$$

or equivalently,

$$J \cdot \mathcal{R}[K_0 \partial_J^l] \bar{Z}_B[J, \phi^*] = O(1/\Lambda_0) \quad (10.75)$$

for an arbitrary source J .^{**)} This is the statement of the BRST invariance of the bare theory defined at the UV scale Λ_0 . As a consequence of (10.64), the QME for the Wilson action $\bar{S}_\Lambda$ is given either as

$$\bar{\Sigma}_\Lambda[\Phi, \Phi^*] = O(1/\Lambda_0), \quad (10.76)$$

or

$$J \cdot R[K \partial_J^l, \Phi^*] \bar{Z}_\Lambda[J, \Phi^*] = O(1/\Lambda_0). \quad (10.77)$$

Before concluding this subsection, we consider a particular class of BRST transformation

$$\delta\phi^A = K_0 \mathcal{R}^A[\phi] = K_0 \left(\mathcal{R}^{(1)A}_B(\Lambda_0) \phi^B + \frac{1}{2} \mathcal{R}^{(2)A}_{BC}(\Lambda_0) \phi^B \phi^C \right), \quad (10.78)$$

which are at most quadratic in fields. The BRST transformation for QED and YM theories belongs to this class. We wish to obtain an explicit expression for the corresponding IR composite operator $\delta\Phi^A$. In order to rewrite the expectation value

$$\langle \delta\phi^A \rangle_{K_0^{-1}J} = K_0 \mathcal{R}^A[K_0 \partial_J^l] \bar{Z}_B[J, \phi^*]$$

for the IR theory, we need to compute two things.

First, we compute the first order differential:

$$\begin{aligned} K_0 \frac{\partial^l}{\partial J_A} \bar{Z}_B[J, \phi^*] &= K_0 \frac{\partial^l}{\partial J_A} N_J \bar{Z}_\Lambda[J, \Phi^*] \\ &= N_J \left[(-)^{\epsilon_A+1} \left(\frac{(K_0 - K)}{K} D^{-1} \right)^{AB} J_B + K_0 \frac{\partial^l}{\partial J_A} \right] \bar{Z}_\Lambda[J, \Phi^*] \\ &= N_J \left\langle K_0 K^{-1} \Phi^A + (K_0 - K) (D^{-1})^{AB} \frac{\partial^l \bar{S}_\Lambda}{\partial \Phi^B} \right\rangle_{\bar{S}_\Lambda, K^{-1}J} \\ &= N_J \left\langle \Phi^A + (K_0 - K) (D^{-1})^{AB} \frac{\partial^l \bar{S}_{I,\Lambda}}{\partial \Phi^B} \right\rangle_{\bar{S}_\Lambda, K^{-1}J} \\ &= N_J \left\langle [\Phi^A]^* \right\rangle_{\bar{S}_\Lambda, K^{-1}J}, \end{aligned} \quad (10.79)$$

^{*)} We write $O(1/\Lambda_0)$ only to mean that it vanishes as $\Lambda_0 \rightarrow \infty$.

^{**)} Later in eqs. (11.4) and (11.15), we write $\bar{S}_B[\phi, \phi^*]$ explicitly for gauge theories. Thanks to renormalizability, the action has only a finite number of parameters. Our assumption is that we can tune the parameters so that eq. (10.74) holds.

where we define

$$\begin{aligned} [\Phi^A]^* &\equiv K_0 K^{-1} \Phi^A + (K_0 - K) (D^{-1})^{AB} \frac{\partial^l \bar{S}_\Lambda}{\partial \Phi^B} \\ &= \Phi^A + (K_0 - K) (D^{-1})^{AB} \frac{\partial^l \bar{S}_{I,\Lambda}}{\partial \Phi^B}. \end{aligned} \quad (10.80)$$

Second, we compute the second order differential:

$$\begin{aligned} K_0 \frac{\partial^l}{\partial J_A} K_0 \frac{\partial^l}{\partial J_B} \bar{Z}_B[J, \phi^*] &= K_0 \frac{\partial^l}{\partial J_A} K_0 \frac{\partial^l}{\partial J_B} N_J \bar{Z}_\Lambda[J, \Phi^*] \\ &\equiv N_J \left\langle [\Phi^A \Phi^B]^* \right\rangle_{\bar{S}_\Lambda, K^{-1}J}, \end{aligned} \quad (10.81)$$

where we define

$$\begin{aligned} [\Phi^A \Phi^B]^* &\equiv [\Phi^A]^* [\Phi^B]^* \\ &+ (K_0 - K) (D^{-1})^{AC} (K_0 - K) (D^{-1})^{BD} \frac{\partial^l \partial^l \bar{S}_{I,\Lambda}}{\partial \Phi^C \partial \Phi^D}. \end{aligned} \quad (10.82)$$

Note that nontrivial contributions arise from derivatives ∂_J acting on the normalization factor N_J .

Hence, using (10.68), we obtain

$$\langle \delta \Phi^A \rangle_{\bar{S}_\Lambda, K^{-1}J} = K \left\langle \mathcal{R}^{(1)A}_B(\Lambda_0) [\Phi^B]^* + \frac{1}{2} \mathcal{R}^{(2)A}_{BC}(\Lambda_0) [\Phi^B \Phi^C]^* \right\rangle_{\bar{S}_\Lambda, K^{-1}J}. \quad (10.83)$$

Since this is valid for arbitrary J , we obtain the operator equality

$$\delta \Phi^A = K \left(\mathcal{R}^{(1)A}_B(\Lambda_0) [\Phi^B]^* + \frac{1}{2} \mathcal{R}^{(2)A}_{BC}(\Lambda_0) [\Phi^B \Phi^C]^* \right). \quad (10.84)$$

It is important to mention the necessity of the cutoff function K_0 to make (10.84) UV finite.*)

10.4. Antifield dependence of the Wilson action

In this subsection we derive the general structure of the Wilson action $\bar{S}_\Lambda[\Phi, \Phi^*]$, in particular its dependence on the antifields. We first make some assumptions for the bare action that are general enough to accommodate the cases of QED and Yang-Mills theories. Then, we describe how the Wilson action depends on the antifields. In §11 we will give the Wilson actions for QED and Yang-Mills theories more explicitly.

As for QED, the Wilson action first obtained in ref. 53) has two different types of antifield dependence, one that can be written down explicitly and the other that appears as a linear shift of the field variables by antifields. These features are seen in Eqs. (11.7)-(11.9) in the next section.***) As for non-Abelian gauge theories, additional non-trivial antifield dependence exists. In this section, we will show how to

*) The potential UV divergence is hard to see in the matrix notation. It is hidden in the loop momentum integral contained in $\mathcal{R}^{(2)A}_{BC}[\Phi^B \Phi^C]$.

**) Eqs.(11.7)-(11.9) differ from those given in ref. 53), where no UV cutoff is introduced.

modify the method of 53) to treat the non-Abelian gauge symmetry. Based on this, we will give explicitly the Wilson action for Yang-Mills theories in (11.22).

Without losing generality, we assume that the bare action is linear in antifields and the BRST transformation quadratic in fields:

$$\bar{S}_B[\phi, \phi^*] = -\frac{1}{2}\phi^A K_0^{-1} D_{AB} \phi^B + \bar{S}_{I,B}[\phi, \phi^*], \quad (10.85)$$

$$\bar{S}_{I,B}[\phi, \phi^*] = S_{I,B}[\phi] + K_0 \phi_A^* \mathcal{R}^A[\phi], \quad (10.86)$$

$$\mathcal{R}^A[\phi] = \mathcal{R}^{(1)A}{}_B \phi^B + \frac{1}{2} \mathcal{R}^{(2)A}{}_{BC} \phi^B \phi^C. \quad (10.87)$$

The bare actions of QED and Yang-Mills theories can be given in this form.

The functional integral (10.30) over the UV fields χ defines the interaction part $\bar{S}_{I,A}$ of the Wilson action, $\bar{S}_A[\Phi, \Phi^*]$. We rewrite the integrand of (10.30) as

$$\begin{aligned} & -\frac{1}{2}\chi \cdot (K_0 - K)^{-1} D \cdot \chi + \bar{S}_{I,B}[\Phi + \chi, K_0^{-1} K \Phi^*] \\ & = -\frac{1}{2}\chi \cdot (K_0 - K)^{-1} D \cdot \chi + K \Phi_A^* \mathcal{R}^A[\Phi + \chi] + S_{I,B}[\Phi + \chi] \\ & = -\frac{1}{2}\chi \cdot (K_0 - K)^{-1} D \cdot \chi + K \Phi_A^* \mathcal{R}^A[\Phi] + \mathcal{J}_A \chi^A \\ & \quad + \frac{1}{2} K \Phi_A^* \chi^B \mathcal{R}^{(2)A}{}_{BC} \chi^C + S_{I,B}[\Phi + \chi], \end{aligned} \quad (10.88)$$

where the effective sources coupled to χ^A are given by

$$\mathcal{J}_A \equiv K \Phi_B^* \left(\mathcal{R}^{(1)B}{}_A + \mathcal{R}^{(2)B}{}_{CA} \Phi^C \right), \quad (10.89)$$

and

$$S_{I,B}[\Phi + \chi] \equiv \bar{S}_{I,B}[\Phi + \chi, K_0^{-1} K \Phi^*]|_{\Phi^*=0}. \quad (10.90)$$

For the terms quadratic in χ and linear in the antifield, we replace χ^A by the derivative $\partial/\partial \mathcal{J}^A$. The Wilson action then takes the form

$$\begin{aligned} \exp[\bar{S}_{I,A}[\Phi, \Phi^*]] &= \exp \left[K \Phi_A^* \left(\mathcal{R}^A[\Phi] + \frac{1}{2} \mathcal{R}^{(2)C}{}_{AB} \frac{\partial^l}{\partial \mathcal{J}_A} \frac{\partial^l}{\partial \mathcal{J}_B} \right) \right] \\ &\times \int \mathcal{D}\chi \exp \left(-\frac{1}{2} \chi \cdot (K_0 - K)^{-1} D \cdot \chi + \mathcal{J} \cdot \chi + S_{I,B}[\Phi + \chi] \right). \end{aligned} \quad (10.91)$$

In order to simplify the above further, we follow ref. 53) and introduce a change of variables. First, we complete the square with respect to χ in the first two terms of the integrand:

$$\begin{aligned} & -\frac{1}{2}\chi \cdot (K_0 - K)^{-1} D \cdot \chi + \mathcal{J} \cdot \chi \\ & = -\frac{1}{2}\chi' \cdot (K_0 - K)^{-1} D \cdot \chi' + \frac{1}{2}(-)^{\epsilon(\mathcal{J})} \mathcal{J} \cdot (K_0 - K) D^{-1} \mathcal{J}, \end{aligned} \quad (10.92)$$

where

$$\chi' \equiv \chi - \mathcal{J} \cdot (K_0 - K) D^{-1} = \chi - (-)^{\epsilon(\mathcal{J})} D^{-1} (K_0 - K) \cdot \mathcal{J}. \quad (10.93)$$

We then introduce new variables Φ'^A so that $\Phi^A + \chi^A = \Phi'^A + \chi'^A$:

$$\Phi'^A \equiv \Phi^A + \mathcal{J}_B(K_0 - K)(D^{-1})^{BA}. \quad (10.94)$$

Finally we obtain

$$\begin{aligned} \exp[\bar{S}_{I,\Lambda}[\Phi, \Phi^*]] &= \exp\left[K\Phi_A^*\mathcal{R}^A[\Phi] + \frac{K}{2}\Phi_A^*\mathcal{R}^{(2)C}_{AB}\frac{\partial^l}{\partial\mathcal{J}_A}\frac{\partial^l}{\partial\mathcal{J}_B}\right] \\ &\times \exp\left[\frac{1}{2}(-)^{\epsilon(\mathcal{J})}\mathcal{J} \cdot (K_0 - K)D^{-1} \cdot \mathcal{J}\right] \\ &\times \int \mathcal{D}\chi' \exp\left[-\frac{1}{2}\chi' \cdot (K_0 - K)^{-1}D \cdot \chi' + S_{I,B}[\Phi' + \chi']\right]. \end{aligned} \quad (10.95)$$

Using the definition of $S_{I,\Lambda}$ in the absence of antifields

$$\exp[S_{I,\Lambda}[\Phi']] \equiv \int \mathcal{D}\chi' \exp\left(-\frac{1}{2}\chi' \cdot (K_0 - K)^{-1}D \cdot \chi' + S_{I,B}[\Phi' + \chi']\right), \quad (10.96)$$

we obtain our final expression for the Wilson action:

$$\begin{aligned} \bar{S}_\Lambda[\Phi, \Phi^*] &= -\frac{1}{2}\Phi^A K^{-1}D_{AB}\Phi^B + K\Phi_A^*\mathcal{R}^A[\Phi] \\ &+ \ln\left[\exp\left\{\frac{K}{2}\Phi_A^*\mathcal{R}^{(2)A}_{BC}\frac{\partial^l}{\partial\mathcal{J}_B}\frac{\partial^l}{\partial\mathcal{J}_C}\right\}\right. \\ &\quad \left.\times \exp\left\{\frac{1}{2}(-)^{\epsilon(\mathcal{J})}\mathcal{J} \cdot (K_0 - K)D^{-1} \cdot \mathcal{J} + S_{I,\Lambda}[\Phi']\right\}\right]. \end{aligned} \quad (10.97)$$

With (10.97) we can construct the action $\bar{S}_{I,\Lambda}[\Phi, \Phi^*]$ from $S_{I,\Lambda}[\Phi]$. The antifield dependence comes partly from $\mathcal{J}$ (10.89), and partly from the shifted variables Φ' (10.94). The remaining antifield dependence is solely generated by the exponentiated differential operator $(K/2)\Phi^*R^{(2)}\partial_{\mathcal{J}}\partial_{\mathcal{J}}$ in the logarithm. This complicates $\bar{S}_\Lambda$ and the BRST transformation of the fields Φ^A . Once the antifields are removed, however, we have simple expressions:⁶⁶⁾

$$\begin{aligned} \delta\Phi^A|_{\Phi^*=0} &= \left[\frac{\partial^l \bar{S}_{I,\Lambda}}{\partial\Phi_A^*}\right]_{\Phi^*=0} \\ &= K\left(\mathcal{R}^{(1)A}_B[\Phi^A] + \frac{1}{2}\mathcal{R}^{(2)A}_{BC}[\Phi^A \Phi^B]\right), \end{aligned} \quad (10.98)$$

where

$$\begin{aligned} [\Phi^A] &\equiv \Phi^A + (K_0 - K)(D^{-1})^{AB}\frac{\partial^l S_{I,\Lambda}}{\partial\Phi^B}, \\ [\Phi^A \Phi^B] &\equiv [\Phi^A][\Phi^B] \end{aligned} \quad (10.99)$$

$$+ (K_0 - K)(D^{-1})^{AC}(K_0 - K)(D^{-1})^{BD}\frac{\partial^l \partial^l S_{I,\Lambda}}{\partial\Phi^C \partial\Phi^D}. \quad (10.100)$$

We can obtain the same results from (10.84) simply by setting the antifields to zero.

Before closing this subsection, we make a comment on the ghost equation of motion: we wish to point out that the actions for gauge theories depend on the antighost and the antifield through a particular linear combination.

Let us consider the classical action (10·3) either for QED or for Yang-Mills theories. The gauge-fixing and the ghost parts of $S_{cl}[\phi]$ are written as⁷³⁾

$$\delta\left(\bar{c}^a F^a(\phi)\right), \quad (10\cdot101)$$

where $F^a(\phi)$ is a gauge-fixing condition. If we choose the covariant gauge fixing

$$F^a(\phi) = -i\left(\partial_\mu a_\mu^a + \frac{\xi}{2}b^a\right), \quad (10\cdot102)$$

where a_μ^a is a gauge field, and b an auxiliary field, we obtain

$$\begin{aligned} \delta\left(\bar{c}^a F^a(\phi)\right) + \phi_A^* \delta\phi^A &= ib^a F^a(\phi) + \left(\bar{c}^a \frac{\partial^r F^a}{\partial\phi^A} + \phi_A^*\right) \delta\phi^A \\ &= \left(-i\bar{c}^a \partial_\mu + a_\mu^{a*}\right) \delta a_\mu^a + \dots \end{aligned} \quad (10\cdot103)$$

Therefore, in $\bar{S}_{cl}[\phi, \phi^*]$, the antifield $a_\mu^{a*}(-p)$ appears as a linear combination with the antighost, $a_\mu^{a*}(-p) + \bar{c}(-p)p_\mu$. Hence, the action $\bar{S}_{cl}[\phi, \phi^*]$ satisfies

$$\frac{\partial^l \bar{S}_{cl}}{\partial \bar{c}(-p)} - p_\mu \frac{\partial^l \bar{S}_{cl}}{\partial a_\mu^{a*}(-p)} = 0. \quad (10\cdot104)$$

We now assume that the bare action satisfies the same equation:

$$K_0 \frac{\partial^l \bar{S}_B}{\partial \bar{c}(-p)} - p_\mu K_0^{-1} \frac{\partial^l \bar{S}_B}{\partial a_\mu^{a*}(-p)} = 0. \quad (10\cdot105)$$

This implies the same relation for the Wilson action

$$K \frac{\partial^l \bar{S}_\Lambda}{\partial \bar{C}(-p)} - p_\mu K^{-1} \frac{\partial^l \bar{S}_\Lambda}{\partial A_\mu^{a*}(-p)} = 0, \quad (10\cdot106)$$

because $K \partial \bar{S}_\Lambda / \partial \bar{\Phi}^A$ and $K^{-1} \partial \bar{S}_\Lambda / \partial \Phi_A^*$ are both composite operators. We call Eqs. (10·104), (10·105), (10·106) the ghost equations of motion. We encountered these before, first as (6·51) and then as (8·23). In fact in §8.2, (8·23) plays an important role in simplifying the perturbative construction of the theory. We will assume (10·105) and hence (10·106) when we discuss examples in the next section.

10.5. QME and the modified Slavnov-Taylor identity

In §2.4 we have discussed the relation between a Wilson action S_Λ and the corresponding effective average action $\Gamma_{B,\Lambda}$. The two are related by a Legendre transformation. A theory can be constructed equally well in terms of S_Λ or $\Gamma_{B,\Lambda}$. As for realization of symmetry, however, we have so far considered only the Wilson action. In this subsection we make a brief detour to write down the identity for

$\bar{\Gamma}_{B,\Lambda}$, corresponding to the QME for $\bar{S}_\Lambda$. This is called the **modified Slavnov-Taylor identity**.⁴⁰⁾ We also give a brief note on the same subject as Appendix D. Compared with Appendix D, we employ a more functional method here.

As in §2.4 we introduce an action with both UV and IR cutoffs:

$$\bar{S}_{B,\Lambda}[\phi, \phi^*] \equiv -\frac{1}{2}\phi \cdot (K_0 - K)^{-1} D \cdot \phi + S_{I,B}[\phi] + K_0 \phi^* \cdot \mathcal{R}[\phi] . \quad (10\cdot107)$$

The gaussian term suppresses the momentum modes $p^2 > \Lambda_0^2$ or $p^2 < \Lambda^2$. The generating functional is defined by

$$\exp [\bar{W}_{B,\Lambda}[J, \phi^*]] \equiv \int \mathcal{D}\phi \exp \left(\bar{S}_{B,\Lambda}[\phi, \phi^*] + K_0^{-1} J \cdot \phi \right) . \quad (10\cdot108)$$

The Legendre transform of $\bar{W}_{B,\Lambda}$ is defined by

$$\bar{\Gamma}_{B,\Lambda}[\varphi, \phi^*] \equiv \bar{W}_{B,\Lambda}[J, \phi^*] - K_0^{-1} J \cdot \varphi , \quad (10\cdot109)$$

where φ is defined in terms of J as

$$K_0^{-1} \varphi^A(p) \equiv \frac{\partial^l \bar{W}_{B,\Lambda}[J, \phi^*]}{\partial J^A(-p)} . \quad (10\cdot110)$$

In the following we wish to consider the functional integral over the QM operator for the bare action:

$$\bar{\Sigma}_{B,\Lambda}^{1PI} \equiv \exp [-\bar{W}_{B,\Lambda}[J, \phi^*]] \int \mathcal{D}\phi \bar{\Sigma}_B[\phi, \phi^*] \exp (\bar{S}_{B,\Lambda}[\phi, \phi^*] + K_0^{-1} J \cdot \phi) , \quad (10\cdot111)$$

and express this in terms of the effective average action $\bar{\Gamma}_{B,\Lambda}$.

As a preparation, we first compute the derivatives of $\bar{\Gamma}_{B,\Lambda}[\varphi, \phi^*]$:

$$\frac{\partial^l \bar{\Gamma}_{B,\Lambda}}{\partial \phi_A^*} = \frac{\partial^l \bar{W}_{B,\Lambda}}{\partial \phi_A^*} + \frac{\partial^l \bar{J}_C}{\partial \phi_A^*} \left(\frac{\partial^l \bar{W}_{B,\Lambda}}{\partial J^C} - K_0^{-1} \varphi^C \right) = \frac{\partial^l \bar{W}_{B,\Lambda}}{\partial \phi_A^*} , \quad (10\cdot112)$$

$$\frac{\partial^r \bar{\Gamma}_{B,\Lambda}}{\partial \varphi^A} = \left(\frac{\partial^r \bar{W}_{B,\Lambda}}{\partial J^C} - (-)^{\epsilon_B} K_0^{-1} \varphi^C \right) \frac{\partial^r J^B}{\partial \varphi^A} - K_0^{-1} J^A = -K_0^{-1} J^A . \quad (10\cdot113)$$

Hence, we obtain

$$\begin{aligned} \frac{\partial^r \bar{\Gamma}_{B,\Lambda}}{\partial \varphi^A} \frac{\partial^l \bar{\Gamma}_{B,\Lambda}}{\partial \phi_A^*} &= -K_0^{-1} J^A \frac{\partial^l \bar{W}_{B,\Lambda}}{\partial \phi_A^*} = -e^{-\bar{W}_{B,\Lambda}} \left(K_0^{-1} J^A \frac{\partial^l}{\partial \phi_A^*} \right) e^{\bar{W}_{B,\Lambda}} \\ &= -e^{-\bar{W}_{B,\Lambda}} \int \mathcal{D}\phi \frac{\partial^r}{\partial \phi^A} e^{K_0^{-1} J \cdot \phi} \cdot \frac{\partial^l \bar{S}_{B,\Lambda}}{\partial \phi_A^*} e^{\bar{S}_{B,\Lambda}[\phi, \phi^*]} \\ &= e^{-\bar{W}_{B,\Lambda}} \int \mathcal{D}\phi \bar{\Sigma}_{B,\Lambda} \exp [\bar{S}_{B,\Lambda}[\phi, \phi^*] + K_0^{-1} J \cdot \phi] , \end{aligned} \quad (10\cdot114)$$

where $\bar{\Sigma}_{B,\Lambda}$ is the QM operator for the action $\bar{S}_{B,\Lambda}$:

$$\bar{\Sigma}_{B,\Lambda} \equiv \frac{1}{2}(\bar{S}_{B,\Lambda}, \bar{S}_{B,\Lambda}) + \Delta \bar{S}_{B,\Lambda} . \quad (10\cdot115)$$

To find the difference between $\bar{\Sigma}_{B,\Lambda}$ and $\bar{\Sigma}_B$, we note

$$\bar{S}_B[\phi, \phi^*] = \bar{S}_{B,\Lambda}[\phi, \phi^*] + \frac{1}{2}\phi \cdot R_\Lambda \cdot \phi, \quad (10.116)$$

where we define

$$[R_\Lambda(p)]_{AB} \equiv D_{AB}(p) \left(\frac{1}{K_0 - K} - \frac{1}{K_0} \right). \quad (10.117)$$

Hence, we obtain

$$\bar{\Sigma}_B = \bar{\Sigma}_{B,\Lambda} + [R_\Lambda]_{BA} \phi^B \frac{\partial^l \bar{S}_{B,\Lambda}}{\partial \phi_A^*}. \quad (10.118)$$

The functional integral over the difference gives

$$\begin{aligned} & e^{-\bar{W}_{B,\Lambda}} \int \mathcal{D}\phi [R_\Lambda]_{BA} \phi^B \frac{\partial^l \bar{S}_{B,\Lambda}}{\partial \phi_A^*} \exp [\bar{S}_{B,\Lambda}[\phi, \phi^*] + K_0^{-1} J \cdot \phi] \\ &= e^{-\bar{W}_{B,\Lambda}} [R_\Lambda]_{BA} K_0 \frac{\partial^l}{\partial J_B} \frac{\partial^l}{\partial \phi_A^*} e^{\bar{W}_{B,\Lambda}} \\ &= [R_\Lambda]_{BA} K_0 \left(\frac{\partial^l \varphi^C}{\partial J_B} \frac{\partial^l}{\partial \varphi^C} \frac{\partial^l \bar{\Gamma}_{B,\Lambda}}{\partial \phi_A^*} + K_0^{-1} \varphi^B \frac{\partial^l \bar{\Gamma}_{B,\Lambda}}{\partial \phi_A^*} \right), \end{aligned} \quad (10.119)$$

where we have used (10.112), (10.113). To summarize so far, the functional integral over (10.118) gives

$$\begin{aligned} \bar{\Sigma}_{B,\Lambda}^{1PI} &= \frac{\partial^r \bar{\Gamma}_{B,\Lambda}}{\partial \varphi^A} \frac{\partial^l \bar{\Gamma}_{B,\Lambda}}{\partial \phi_A^*} \\ &+ [R_\Lambda]_{BA} \left(K_0 \frac{\partial^l \varphi^C}{\partial J_B} \frac{\partial^l}{\partial \varphi^C} \frac{\partial^l \bar{\Gamma}_{B,\Lambda}}{\partial \phi_A^*} + \varphi^B \frac{\partial^l \bar{\Gamma}_{B,\Lambda}}{\partial \phi_A^*} \right). \end{aligned} \quad (10.120)$$

To further rewrite the second term in the above, we introduce the notation:

$$(\bar{\Gamma}_{B,\Lambda}^{(2)})_{CA} \equiv \frac{\partial^l}{\partial \varphi^C} \frac{\partial^r \bar{\Gamma}_{B,\Lambda}}{\partial \varphi^A} = -K_0^{-1} \frac{\partial^l J^A}{\partial \varphi^C}, \quad (10.121)$$

where (10.113) is used. Then, the inverse is written as

$$\frac{\partial^l \varphi^A}{\partial J_C} = -K_0^{-1} (\bar{\Gamma}_{B,\Lambda}^{(2)})_{CA}^{-1}. \quad (10.122)$$

Hence, we can rewrite (10.120) as

$$\begin{aligned} \bar{\Sigma}_{B,\Lambda}^{1PI} &= \frac{\partial^r \bar{\Gamma}_{B,\Lambda}}{\partial \varphi^A} \frac{\partial^l \bar{\Gamma}_{B,\Lambda}}{\partial \phi_A^*} \\ &+ [R_\Lambda]_{BA} \left(-(\bar{\Gamma}_{B,\Lambda}^{(2)})_{BC}^{-1} \frac{\partial^l}{\partial \varphi^C} \frac{\partial^l \bar{\Gamma}_{B,\Lambda}}{\partial \phi_A^*} + \varphi^B \frac{\partial^l \bar{\Gamma}_{B,\Lambda}}{\partial \phi_A^*} \right). \end{aligned} \quad (10.123)$$

Imposing this to vanish, we obtain the modified Slavnov-Taylor identity:⁴⁰⁾

$$\bar{\Sigma}_{B,\Lambda}^{1PI} = 0. \quad (10.124)$$

In the limit of $\Lambda \rightarrow 0$, R_Λ goes to zero and $\bar{\Gamma}_{B,\Lambda}$ becomes $\bar{\Gamma}_B$. Therefore, the modified ST identity reduces to the Zinn-Justin equation.

10.6. On the nilpotency without antifields

As shown toward the end of §10.1, the quantum master equation (10·17) implies the nilpotency of the quantum BRST transformation δ_Q , defined by (10·18). Even if δ_Q is nilpotent, the reduced BRST transformation

$$\delta \equiv \delta_Q|_{\Phi^*=0} \quad (10\cdot125)$$

that appears in the WT operator (10·72) is not nilpotent in general. In this subsection we compute δ^2 , and study how it differs from zero, assuming the nilpotency $\delta_Q^2 = 0$ of the full BRST transformation.

Take a functional $\mathcal{O}_\Lambda[\Phi]$ of the field Φ defined at an IR scale Λ , and consider its BRST transformation:

$$\delta_Q \mathcal{O}_\Lambda[\Phi] = \frac{\partial^r \mathcal{O}_\Lambda[\Phi]}{\partial \Phi^A} \frac{\partial^l \bar{S}_\Lambda}{\partial \Phi_A^*}. \quad (10\cdot126)$$

Expanding the Wilson action in powers of antifields

$$\bar{S}_\Lambda[\Phi, \Phi^*] = S_\Lambda[\Phi] + \Phi_A^* f^A[\Phi] + \frac{1}{2} \Phi_A^* \Phi_B^* f^{BA}[\Phi] + \mathcal{O}\left((\Phi^*)^3\right), \quad (10\cdot127)$$

we obtain an expansion

$$\delta_Q \Phi^A \equiv \frac{\partial^l \bar{S}_\Lambda}{\partial \Phi_A^*} = f^A[\Phi] + \Phi_B^* f^{BA}[\Phi] + \mathcal{O}\left((\Phi^*)^2\right). \quad (10\cdot128)$$

Setting $\Phi^* = 0$, we obtain

$$\delta \Phi^A = f^A[\Phi]. \quad (10\cdot129)$$

By the definition (10·18), the nilpotency of δ_Q implies

$$0 = \left(\frac{\partial^r}{\partial \Phi^A} \delta_Q \mathcal{O}_\Lambda \right) \frac{\partial^l \bar{S}_\Lambda}{\partial \Phi_A^*} - \left(\frac{\partial^r}{\partial \Phi_A^*} \delta_Q \mathcal{O}_\Lambda \right) \frac{\partial^l \bar{S}_\Lambda}{\partial \Phi^A} - (-)^{\epsilon_A} \frac{\partial^r}{\partial \Phi^A} \frac{\partial^r}{\partial \Phi_A^*} \left(\delta_Q \mathcal{O}_\Lambda \right). \quad (10\cdot130)$$

Setting Φ^* to zero, the first term on the right gives $\delta^2 \mathcal{O}_\Lambda$. Therefore, we obtain

$$\delta^2 \mathcal{O}_\Lambda = \left[\frac{\partial^r}{\partial \Phi_A^*} \delta_Q \mathcal{O}_\Lambda \right]_{\Phi^*=0} \frac{\partial^l S_\Lambda}{\partial \Phi^A} + (-)^{\epsilon_A} \frac{\partial^r}{\partial \Phi^A} \left[\frac{\partial^r}{\partial \Phi_A^*} \delta_Q \mathcal{O}_\Lambda \right]_{\Phi^*=0}. \quad (10\cdot131)$$

Using the expansions (10·127) & (10·128), we obtain

$$\begin{aligned} \delta^2 \mathcal{O}_\Lambda[\Phi] &= - \left(\frac{\partial^r \mathcal{O}_\Lambda}{\partial \Phi^A} f^{AB} \right) \frac{\partial^r S_\Lambda}{\partial \Phi^B} - \frac{\partial^r}{\partial \Phi^B} \left(\frac{\partial^r \mathcal{O}_\Lambda}{\partial \Phi^A} f^{AB} \right) \\ &= -e^{-S_\Lambda} \frac{\partial^r}{\partial \Phi^B} \left(\frac{\partial^r \mathcal{O}_\Lambda}{\partial \Phi^A} f^{AB} e^{S_\Lambda} \right). \end{aligned} \quad (10\cdot132)$$

In particular, we obtain

$$\delta^2 \Phi^A = -f^{AB} \frac{\partial^r S_\Lambda}{\partial \Phi^B} - \frac{\partial^r}{\partial \Phi^B} f^{AB} = -e^{-S_\Lambda} \frac{\partial^r}{\partial \Phi^B} (f^{AB} e^{S_\Lambda}). \quad (10\cdot133)$$

Thus, we have found that δ fails to be nilpotent in a particular way that $e^{S_\Lambda} \delta^2 \mathcal{O}_\Lambda$ is a total derivative, or using the terminology introduced in §4.2, $\delta^2 \mathcal{O}_\Lambda$ is a generalized equation of motion.*)

*) Note that $f^{AB}[\Phi]$ is not necessarily a composite operator. Hence, we are abusing the terminology introduced for genuine composite operators.

§11. Examples: the Wilson actions for QED and Yang-Mills theories

The purpose of this section is to apply the general result (10.97) on the antifield dependence of the Wilson action to QED and the Yang-Mills theories. As for QED this has already been done in §8.1, from a consideration of the ERG differential equation. This method works only for QED, but not for YM theories. For the latter we have adopted a perturbative approach in §8.2, and have shown how to construct the continuum limit of the Wilson action by loop expansions. In §8.2, we have discussed the antifield dependence of only the asymptotic behavior of the Wilson action, since that is all we need for proving the theory's existence. In contrast, we have adopted a unifying approach in §10, and have determined the antifield dependence of the Wilson action explicitly by starting from a bare action that has simple antifield dependence.*)

11.1. QED

We first consider QED and reproduce the results from 65) and 53). Our derivation follows closely that of 53) except that we pay more careful attention to the necessity of a UV cutoff for the bare theory.

Denoting the UV fields by

$$\phi^A = \{a_\mu, b, c, \bar{c}, \varphi, \bar{\varphi}\}, \quad (11.1)$$

and the UV antifields by

$$\phi_A^* = \{a_\mu^*, b^*, c^*, \bar{c}^*, \varphi^*, \bar{\varphi}^*\}, \quad (11.2)$$

the bare action of QED is written as follows**):

$$\bar{S}_B[\phi, \phi^*] = -\frac{1}{2}\phi \cdot K_0^{-1}D \cdot \phi + K_0\phi_A^*\mathcal{R}^A[\phi] + S_{I,B}[\phi], \quad (11.3)$$

where

$$\begin{aligned} \frac{1}{2}\phi \cdot K_0^{-1}D \cdot \phi &\equiv \int_p K_0^{-1}(p) \left[\frac{1}{2}a_\mu(-p)(p^2\delta_{\mu\nu} - p_\mu p_\nu)a_\nu(p) + i\bar{c}(-p)p^2c(p) \right. \\ &\quad \left. - b(-p)\left\{ ip_\mu a_\mu(p) + \frac{\xi}{2}b(p) \right\} + \bar{\varphi}(-p)(\not{p} + im)\varphi(p) \right], \\ K_0\phi_A^*\mathcal{R}^A[\phi] &\equiv \int_p K_0(p) \left[a_\mu^*(-p)(-i)p_\mu c(p) + \bar{c}^*(-p) \cdot ib(p) \right. \\ &\quad \left. + ie \int_q \varphi^*(-p)c(q)\varphi(p-q) + ie \int_q \bar{\varphi}(-p-q)c(q)\bar{\varphi}^*(p) \right], \end{aligned}$$

) The antifield dependence of the bare action is linear in ϕ^ and at most quadratic in ϕ as in (10.87).

**) We have introduced the auxiliary field b to avoid the quadratic dependence of $\bar{S}_B$ on $\bar{c}^*$. The result (8.12) is obtained from (11.11) by integrating out B .

and

$$\begin{aligned}
S_{I,B}[\phi] = & \int_p \left[\frac{a_2}{2} A_0^2 a_\mu(-p) a_\mu(p) + \frac{z_2}{2} a_\mu(-p) p^2 a_\mu(p) + \frac{z'_2}{2} a_\nu(-p) p_\nu p_\mu a_\mu(p) \right. \\
& + \frac{z_4}{8} \int_{q,k,r} \delta^4(p+q+k+r) a_\nu(p) a_\nu(q) a_\mu(k) a_\mu(r) \\
& \left. + \bar{\varphi}(-p) (z_f \not{p} + z_m im) \varphi(p) + z_3 \int_q \bar{\varphi}(-p) \not{q} \varphi(p-q) \right]. \quad (11.4)
\end{aligned}$$

The coefficients

$$a_2, z_2, z'_2, z_3, z_4, z_f, z_m \quad (11.5)$$

are given such dependence on $\ln \Lambda_0/\mu$ that we obtain a finite continuum limit.

Denoting the IR fields and antifields by

$$\Phi^A = \{A_\mu, B, C, \bar{C}, \psi, \bar{\psi}\}, \quad \Phi_A^* = \{A_\mu^*, B^*, C^*, \bar{C}^*, \psi^*, \bar{\psi}^*\}, \quad (11.6)$$

the effective sources defined by (10.89) are given by

$$\begin{aligned}
\mathcal{J}_c(-p) = & -iK(p) p_\mu A_\mu^*(-p) \\
& -ie \int_q K(q) (\psi^*(-q) \psi(q-p) - \bar{\psi}(-q-p) \bar{\psi}^*(q)) , \quad (11.7a)
\end{aligned}$$

$$\mathcal{J}_B(-p) = iK(p) \bar{C}^*(-p), \quad (11.7b)$$

$$\mathcal{J}_\psi(-p) = ie \int_q K(q) \psi^*(-q) C(q-p), \quad (11.7c)$$

$$\mathcal{J}_{\bar{\psi}}(-p) = -ie \int_q K(q) C(q-p) \bar{\psi}^*(-q). \quad (11.7d)$$

Since the ghost and antighost remain free, and are absent in the interaction action $S_{I,B}$, the source $\mathcal{J}_c$ does not appear in the IR action. This simplifies the application of the general result (10.97). The differential operators, $(K\Phi^*/2)R^{(2)}\partial_{\mathcal{J}}\partial_{\mathcal{J}}$ in (10.97), generate no contribution. The Wilson action is thus given as

$$\begin{aligned}
\bar{S}_\Lambda[\Phi, \Phi^*] = & -\frac{1}{2} \Phi^A K^{-1} D_{AB} \Phi^B + K \Phi_A^* \mathcal{R}^A[\Phi] \\
& + \frac{1}{2} (-)^{\epsilon(\mathcal{J})} \mathcal{J} \cdot (K_0 - K) D^{-1} \mathcal{J} + S_{I,\Lambda}[\Phi'], \quad (11.8)
\end{aligned}$$

where the shifted variables are defined by

$$A'_\mu(p) = A_\mu(p) + \frac{ip_\mu}{p^2} (K_0(p) - K(p)) \mathcal{J}_B(p), \quad (11.9a)$$

$$\psi'(p) = \psi(p) - \frac{(K_0(p) - K(p))}{\not{p} + im} \mathcal{J}_{\bar{\psi}}(p), \quad (11.9b)$$

$$\bar{\psi}'(-p) = \bar{\psi}(-p) + \mathcal{J}_\psi(-p) \frac{(K_0(p) - K(p))}{\not{p} + im}. \quad (11.9c)$$

The term quadratic in the effective sources is given by

$$\frac{1}{2}(-)^{\epsilon(\mathcal{J})} \mathcal{J} \cdot (K_0 - K) D^{-1} \mathcal{J} = -\mathcal{J}_\psi(-p) \frac{(K_0(p) - K(p))}{\not{p} + im} \mathcal{J}_{\bar{\psi}}(p). \quad (11.10)$$

Hence, the total Wilson action is

$$\begin{aligned} \bar{S}_A[\Phi, \Phi^*] = \int_p \left[-K(p)^{-1} \left\{ \frac{1}{2} A_\mu(-p) (p^2 \delta_{\mu\nu} - p_\mu p_\nu) A_\nu(p) + \bar{C}(-p) i p^2 C(p) \right. \right. \\ \left. - B(-p) \left(i p_\mu A_\mu(p) + \frac{\xi}{2} B(p) \right) + \bar{\psi}(-p) (\not{p} + im) \psi(p) \right\} \\ + K(p) \left\{ \bar{C}^*(-p) \cdot i B(p) + A_\mu^*(-p) (-i) p_\mu C(p) \right. \\ \left. + i e \int_q (\psi^*(-p) C(q) \psi(p-q) + \bar{\psi}(-p-q) C(q) \bar{\psi}^*(p)) \right\} \\ \left. + e^2 \int_{q,k} K(p+q) K(p-k) \psi^*(-p-q) \frac{(K_0(p) - K(p))}{\not{p} + im} \right. \\ \left. \times \bar{\psi}^*(p-k) C(q) C(k) \right] + S_{I,A}[A'_\mu, \psi', \bar{\psi}']. \quad (11.11) \end{aligned}$$

Since QED is renormalizable, we can tune the $\ln \Lambda_0/\mu$ dependence of the bare parameters (11.5) to obtain the continuum limit of $S_{I,A}$. The remaining dependence of $\bar{S}_A$ on Λ_0 comes from K_0 in the $\psi^* \bar{\psi}^*$ term and the shifted variables $A'_\mu, \psi', \bar{\psi}'$. In the limit $\Lambda_0 \rightarrow \infty$, K_0 is simply replaced by 1, and we obtain the continuum limit of $\bar{S}_A$ as in 53). A canonical transformation that eliminates the quadratic terms in the antifield gives the action obtained in ref. 65). Perturbative construction of $S_{I,A}$ that satisfies the WT identity, $\Sigma_A = 0$, has already been discussed in §6.1.

Finally, we observe that $\bar{S}_A$ given by (11.11) satisfies the ghost equation of motion (10.106). This is a consequence of (10.105) satisfied by the bare action (11.3) as we have pointed out at the end of §10.4.

11.2. Yang-Mills theories

We consider a Yang-Mills theory without matter described by the UV action^{*)}

$$\bar{S}_B[\phi, \phi^*] = -\frac{1}{2} \phi^A K_0^{-1} D_{AB} \phi^B + \bar{S}_{I,B}[\phi, \phi^*], \quad (11.12)$$

where fields and antifields are collectively denoted as

$$\phi^A = \{a_\mu, b, c, \bar{c}\}, \quad \phi_A^* = \{a_\mu^*, b^*, c^*, \bar{c}^*\}. \quad (11.13)$$

The bare action consists of the kinetic part

$$-\frac{1}{2} \phi^A K_0^{-1} D_{AB} \phi^B \equiv - \int_p K_0^{-1}(p) \left[\frac{1}{2} a_\mu(-p) \cdot (p^2 \delta_{\mu\nu} - p_\mu p_\nu) a_\nu(p) \right.$$

^{*)} We suppress the group index and use the notations: $A \cdot B \equiv A^a B^a$, $A^2 \equiv A^a A^a$ and $A \cdot (B \times C) \equiv f^{abc} A^a B^b C^c$.

$$-b(-p) \cdot \{ip_\mu a_\mu(p) + \frac{\xi}{2}b(p)\} + i\bar{c}(-p) \cdot p^2 c(p) \Big] \quad (11.14)$$

and the interaction part

$$\begin{aligned} \bar{S}_{I,B}[\phi, \phi^*] &\equiv S_{I,B}[\phi] + K_0 \phi_A^* \left(\mathcal{R}^{(1)A}{}_B \phi^B + \frac{1}{2} \Phi^B \mathcal{R}^{(2)A}{}_{BC} \phi^C \right) \\ &= \int_p \left[\frac{a_2}{2} \Lambda_0^2 a_\mu(-p) a_\mu(p) + \frac{z_1}{2} p^2 a_\mu(p) a_\mu(-p) + \frac{z_2}{2} p_\mu p_\nu a_\mu(-p) a_\nu(p) \right] \\ &\quad + z_3 \int_{p,q} p_\nu a_\mu(-p) \cdot [a_\mu(q) \times a_\nu(p-q)] \\ &\quad + \int_{p_1, \dots, p_4} \delta(\Sigma p_i) \left[\frac{z_4}{8} a_\mu(p_1) \cdot a_\mu(p_2) a_\nu(p_3) \cdot a_\nu(p_4) + \frac{z_5}{8} a_\mu(p_1) \cdot a_\nu(p_2) a_\mu(p_3) \cdot a_\nu(p_4) \right] \\ &\quad + \int_p \left[K_0(p) \left\{ \bar{c}^*(-p) \cdot ib(p) - ia_\mu^*(-p) \cdot p_\mu c(p) + \frac{z_8}{2} c^*(-p) \cdot \int_q c(q) \times c(p-q) \right\} \right. \\ &\quad \left. + (K_0^2(p) a_\mu^*(-p) + p_\mu \bar{c}(-p)) \cdot \left\{ -iz_6 p_\mu c(p) + z_7 \int_q a_\mu(p-q) \times c(q) \right\} \right]. \quad (11.15) \end{aligned}$$

Here, we have assumed the gauge group SU(2) to write down $S_{I,B}[\phi]$; more terms are available and need to be introduced for higher symmetry. The coefficients $a_2, z_1, \dots, z_8$ depend on $\ln \Lambda_0/\mu$.

The Wilson action with the IR fields & antifields

$$\Phi^A = \{A_\mu^a, B^a, C^a, \bar{C}^a\}, \quad \Phi_A^* = \{A_\mu^*, B^*, C^*, \bar{C}^*\} \quad (11.16)$$

is given by the general formula (10.97). Let us get a concrete expression. The effective sources read

$$\mathcal{J}_\mu(-p) = -z_7 \int_q K(q) K_0(q) A_\mu^*(q) \times C(-p-q), \quad (11.17a)$$

$$\mathcal{J}_B(-p) = iK(p) \bar{C}^*(-p), \quad (11.17b)$$

$$\begin{aligned} \mathcal{J}_c(-p) &= -i(1 + z_6 K_0(p)) K(p) p_\mu A_\mu^*(-p) + z_7 \int_q K(q) K_0(q) A_\mu^*(q) \times A_\mu(-p-q) \\ &\quad + z_8 \int_q K(q) C^*(q) \times C(-p-q), \quad (11.17c) \end{aligned}$$

and the differential operator w.r.t. $\mathcal{J}$ is given by

$$\begin{aligned} \frac{K}{2} \Phi_C^* \frac{\partial^l}{\partial \mathcal{J}_A} R^{(2)C}{}_{AB} \frac{\partial^l}{\partial \mathcal{J}_B} &= z_7 \int_{p,q} K(p) K_0(p) A_\mu^*(-p) \cdot \left(\frac{\partial^l}{\partial \mathcal{J}_\mu(-p+q)} \times \frac{\partial^l}{\partial \mathcal{J}_c(-q)} \right) \\ &\quad + \frac{z_8}{2} \int_{p,q} K(p) C^*(-p) \cdot \left(\frac{\partial^l}{\partial \mathcal{J}_c(-p+q)} \times \frac{\partial^l}{\partial \mathcal{J}_c(-q)} \right). \quad (11.18) \end{aligned}$$

The shifted variables defined by (10.94) are given by

$$A'_\mu(p) = A_\mu(p) + \frac{K_0(p) - K(p)}{p^2} \left[\delta_{\mu\nu} - (1 - \xi) \frac{p_\mu p_\nu}{p^2} \right] \mathcal{J}_\nu(p)$$

$$\begin{aligned}
& + \frac{ip_\mu}{p^2} (K_0(p) - K(p)) \mathcal{J}_B(p) , \\
\bar{C}'(p) &= \bar{C}(p) - \frac{i}{p^2} (K_0(p) - K(p)) \mathcal{J}_c(p) .
\end{aligned} \tag{11.19}$$

It is easy to see that $S_{I,A}[\Phi']$ does not depend on B :

$$S_{I,A}[\Phi'] = S_{I,A}[A'_\mu, \bar{C}', C] . \tag{11.20}$$

The quadratic term in $\mathcal{J}$ that appears inside the exponentials of (10.97) takes the following concrete form:

$$\begin{aligned}
\frac{1}{2}(-)^{\epsilon(\mathcal{J})} \mathcal{J} \cdot (K_0 - K) D^{-1} \mathcal{J} &= i \frac{(K_0(p) - K(p))}{p^2} p_\mu \mathcal{J}_\mu(-p) \mathcal{J}_B(p) \\
&+ \frac{1}{2} \mathcal{J}_\mu(-p) \frac{(K_0(p) - K(p))}{p^2} \left[\delta_{\mu\nu} - (1 - \xi) \frac{p_\mu p_\nu}{p^2} \right] \mathcal{J}_\nu(p) .
\end{aligned} \tag{11.21}$$

Putting together the above results, we obtain the Wilson action:

$$\begin{aligned}
\bar{S}_A[\Phi, \Phi^*] &= \int_p \left[-K^{-1}(p) \left\{ \frac{1}{2} A_\mu(-p) \cdot (p^2 \delta_{\mu\nu} - p_\mu p_\nu) A_\nu(p) + \bar{C}(-p) \cdot ip^2 C(p) \right. \right. \\
&\quad \left. \left. - B(-p) \cdot \left(ip_\mu A_\mu(p) + \frac{\xi}{2} B(p) \right) \right\} \right. \\
&\quad + K(p) \bar{C}^*(-p) \cdot iB(p) \\
&\quad + K(p) A_\mu^*(-p) \cdot \left\{ (-i)(1 + z_6 K_0(p)) p_\mu C(p) \right. \\
&\quad \left. + z_7 K_0(p) \int_q A_\mu(p - q) \times C(q) \right\} \\
&\quad \left. + K(p) C^*(-p) \cdot z_8 \frac{1}{2} \int_q C(p - q) \times C(q) \right] \\
&+ \log \left[\exp \left\{ \frac{K}{2} \Phi_A^* \frac{\partial^l}{\partial \mathcal{J}_B} R^{(2)A}_{BC} \frac{\partial^l}{\partial \mathcal{J}_C} \right\} \right. \\
&\quad \left. \times \exp \left\{ \frac{1}{2} (-)^{\epsilon(\mathcal{J})} \mathcal{J} \cdot (K_0 - K) D^{-1} \cdot \mathcal{J} + S_{I,A}[\Phi'] \right\} \right] . \tag{11.22}
\end{aligned}$$

In writing down the bare action (11.12) we have assumed the ghost equation of motion (10.105) to hold. Hence, the Wilson action (11.22) must satisfy (10.106), i.e., the Wilson action must depend on the antighost and the antifield for the gauge field only through the linear combination, $p_\mu \bar{C}^a(-p) + K^2(p) A_\mu^{a*}(-p)$. The way we have written (11.22) for $\bar{S}_A$, this is not manifest.

§12. Concluding remarks

We hope we have succeeded in achieving the two goals that we have set in §1. Formulating a field theory with a momentum cutoff is the most natural, and it accords with Wilson's non-perturbative definition of a field theory. It is still surprising that we can describe the continuum limit using a finite momentum cutoff, but it does not come for free: the cutoff function needs to be smooth, and the Wilson action contains an infinite number of terms. The formalism of ERG is simple enough that we only need several pages to summarize the essence of all done in this review. We have given a quick summary as Appendix F.

We have introduced two ways of realizing continuous symmetry: one with the WT identity, and another with the QME. Among the examples we have discussed in §§6 & 8, only YM theories need the QME for proving the possibility of construction. Let us try to understand this. The WT identity is the invariance of the Wilson action under infinitesimal transformations, and it knows nothing about the algebra formed by the transformations. We need the antifield formalism to incorporate an algebraic structure. Take the example of the two dimensional $O(N)$ non-linear sigma model discussed in §6.4. Commuting two generators that move the N -th axis, we get an element of $O(N-1)$, under which the action is manifestly invariant. Take the Wess-Zumino model. Commuting two supersymmetry generators, we get a translation, under which the action is manifestly invariant. Hence, either in the non-linear sigma model or the Wess-Zumino model, the algebra of infinitesimal transformations does not constrain the theory. Hence, the WT identity suffices. This is not the case with YM theories; we need the antifield formalism to incorporate the full algebra of BRST transformations.*)

As for perturbative applications, our list of examples is not exhaustive. Among those missing, supersymmetric YM theories are particularly important. The ERG formalism has been already applied,^{6),23),35)} but our understanding is still incomplete regarding the non-renormalization properties.¹⁰⁴⁾¹¹⁹⁾ We also think it important to prove the non-renormalization theorem for chiral anomalies using the ERG formalism; to our knowledge it has not been done.

As for non-perturbative applications, the order-by-order solution of the WT identity or QME is inappropriate. Even approximate solutions will do as long as they are given in closed form. Progress along this line is hoped for.

In this review we have been primarily concerned with formulating various symmetric theories. The formalism has matured enough that we are ready to address more physical questions.

Acknowledgment

The authors would like to thank Prof. K. Higashijima for suggestions and encouragement.

*) The antifield formalism is necessary only to prove the possibility of perturbative construction. The WT identity suffices for the order-by-order construction.

Appendix A

— Generalized diffusion equations —

In §2 we have derived a particular ERG differential equation (2·51) for the interaction part of the Wilson action, $S_{I,A}$, defined by (2·23). The purpose of this appendix is twofold:

1. to rewrite (2·23) as a formula that gives the entire Wilson action,
2. to generalize (2·23) to incorporate the original Wilson's differential equation¹²⁴⁾ and Polchinski's⁹⁴⁾ under the same footing.

This is done by (A·2), where different choices of A_t, Z_t correspond to Wilson's and Polchinski's ERG differential equations.

In the following we adopt a notation slightly different from that in the main text. We write S_t for S_A , and S_0 for S_B , where t is defined by

$$t = \ln \frac{\Lambda_0}{\Lambda} \geq 0. \quad (\text{A} \cdot 1)$$

The logarithmic parameter t grows, as the momentum cutoff Λ decreases.

A.1. Derivation of the generalized diffusion equations

We consider a one-parameter family of actions $S_t[\phi]$ ($t \geq 0$) for a real scalar field ϕ . We generate the t -dependence by a gaussian integral transformation of the following type:^{121), 124)}

$$\exp [S_t[\phi]] = \int [d\phi'] \exp \left[-\frac{1}{2} \int_p A_t(p)^2 (\phi(p) - Z_t(p)\phi'(p)) (\phi(-p) - Z_t(p)\phi'(-p)) + S_0[\phi'] \right]. \quad (\text{A} \cdot 2)$$

We impose the two properties

$$\begin{aligned} \text{(i)} \quad & 1/A_0(p) = 0, \\ \text{(ii)} \quad & Z_0(p) = 1, \end{aligned} \quad (\text{A} \cdot 3)$$

in order to assure the limit

$$\lim_{t \rightarrow 0+} S_t = S_0. \quad (\text{A} \cdot 4)$$

The two factors Z_t and $A_t Z_t$ have the following physical meanings:


1. blocking factor $Z_t(p)$ — $Z_t(p)\phi(p)$ is the Fourier transform of a block spin. Denoting the Fourier inverse of $\phi(p)$ and $Z_t(p)$ by the same symbols, we write

$$\phi(x) = \int_p e^{ipx} \phi(p), \quad Z_t(x) = \int_p e^{ipx} Z_t(p). \quad (\text{A} \cdot 5)$$

Then, the inverse Fourier transform of $Z_t(p)\phi(p)$ is the block spin or average field:

$$\phi_t(x) \equiv \int_p e^{ipx} Z_t(p)\phi(p) = \int d^D y Z_t(x-y)\phi(y) \quad (\text{A} \cdot 6)$$

The size of the domain of $Z_t(x)$ should grow exponentially as e^t .

Fig. 27. More integration for higher p^2 and higher t .

2. $A_t(p)Z_t(p)$ — This has to do with incomplete functional integration. The gaussian factor in (A.2) implies that $\phi(p)$ is equal to the block spin $Z_t(p)\phi'(p)$ up to order $1/A_t(p)$. Equivalently $\phi'(p)$ is integrated around $\phi(p)/Z_t(p)$ up to order $1/(A_t(p)Z_t(p))$. We take the squared width of integration, $1/(A_t Z_t(p))^2$, a growing function of both t and p^2 . (Fig. 27)

We will shortly give examples of Z_t and $A_t Z_t$.

We now wish to derive the differential equation for $S_t[\phi]$, analogous to the diffusion equation. We first consider the t -derivative:

$$\begin{aligned}
 & \partial_t \exp[S_t[\phi]] \\
 &= \int [d\phi'] \int_p \left\{ -\partial_t A_t(p) \cdot A_t(p) (\phi(p) - Z_t(p)\phi'(p)) \cdot (\phi(-p) - Z_t(p)\phi'(-p)) \right. \\
 & \quad \left. + A_t^2(p) \partial_t Z_t(p) \phi'(p) (\phi(-p) - Z_t(p)\phi'(-p)) \right\} \\
 & \quad \times \exp \left[-\frac{1}{2} \int_q A_t(q)^2 (\phi(q) - Z_t(q)\phi'(q)) (\phi(-q) - Z_t(q)\phi'(-q)) + S_0[\phi'] \right] \\
 &= \int [d\phi'] \int_p \left\{ -(\partial_t A_t(p) \cdot A_t(p) + A_t^2(p) \partial_t \ln Z_t(p)) \right. \\
 & \quad \times (\phi(p) - Z_t(p)\phi'(p)) (\phi(-p) - Z_t(p)\phi'(-p)) \\
 & \quad \left. + A_t^2(p) \partial_t \ln Z_t(p) \cdot \phi(p) (\phi(-p) - Z_t(p)\phi'(-p)) \right\} \exp[\dots], \quad (\text{A.7})
 \end{aligned}$$

where we have suppressed the argument of the last exponential. We compare the above with the first and second order functional derivatives of $\exp[S_t]$:

$$-\frac{\delta}{\delta\phi(p)} \exp[S_t[\phi]] = \int [d\phi'] A_t(p)^2 (\phi(-p) - Z_t(p)\phi'(-p)) \exp[\dots], \quad (\text{A.8})$$

$$\begin{aligned}
 \frac{\delta^2}{\delta\phi(p)\delta\phi(-p)} \exp[S_t[\phi]] &= \int [d\phi] A_t(p)^2 (\phi(-p) - Z_t(p)\phi'(-p)) \\
 & \quad \times A_t(p)^2 (\phi(-p) - Z_t(p)\phi'(-p)) \exp[\dots], \quad (\text{A.9})
 \end{aligned}$$

where we have ignored an additive constant proportional to the space volume.

Thus, we obtain the generalized diffusion equation

$$\begin{aligned} \partial_t \exp [S_t[\phi]] = \int_p \left[-\frac{1}{A_t^2(p)} \partial_t \ln (A_t Z_t)(p) \frac{\delta^2}{\delta \phi(p) \delta \phi(-p)} \right. \\ \left. - \partial_t \ln Z_t(p) \cdot \phi(p) \frac{\delta}{\delta \phi(p)} \right] \exp [S_t[\phi]] . \end{aligned} \quad (\text{A}\cdot 10)$$

This can be rewritten for S_t as

$$\begin{aligned} \partial_t S_t = \int_p \left[F_t(p) \cdot \phi(p) \frac{\delta S_t}{\delta \phi(p)} \right. \\ \left. + G_t(p) \cdot \frac{1}{2} \left\{ \frac{\delta S_t}{\delta \phi(p)} \frac{\delta S_t}{\delta \phi(-p)} + \frac{\delta^2 S_t}{\delta \phi(p) \delta \phi(-p)} \right\} \right] , \end{aligned} \quad (\text{A}\cdot 11)$$

where

$$\begin{cases} F_t(p) & \equiv -\partial_t \ln Z_t(p) , \\ G_t(p) & \equiv -\frac{2}{A_t(p)^2} \partial_t \ln (A_t Z_t)(p) . \end{cases} \quad (\text{A}\cdot 12)$$

The equation for S_t has the same form as the ERG differential equation (2.55) for the full action.

We now define the generating functional of the connected correlation functions:

$$\exp [W_t[J]] \equiv \int [d\phi] \exp \left[S_t[\phi] + \int_p J(-p) \phi(p) \right] . \quad (\text{A}\cdot 13)$$

Using the technique familiar from §2.2, we can compute $W_t[J]$ in terms of $W_0[J]$ as follows:

$$\begin{aligned} \exp [W_t[J]] &= \int [d\phi] e^{\int_p J(p) \phi(-p)} \int [d\phi'] \exp \left[\right. \\ &\quad \left. -\frac{1}{2} \int_p A_t(p)^2 (\phi(p) - Z_t(p) \phi'(p)) (\phi(-p) - Z_t(p) \phi'(-p)) + S_0[\phi'] \right] \\ &= \int [d\phi] \int [d\phi'] \exp \left[\int_p J(p) (\phi(-p) - Z_t(p) \phi'(-p)) \right. \\ &\quad \left. -\frac{1}{2} \int_p A_t(p)^2 (\phi(p) - Z_t(p) \phi'(p)) (\phi(-p) - Z_t(p) \phi'(-p)) \right. \\ &\quad \left. + S_0[\phi'] + \int_p J(p) Z_t(p) \phi'(-p) \right] . \end{aligned} \quad (\text{A}\cdot 14)$$

Now, shifting ϕ by $Z_t \phi'$, we obtain

$$\begin{aligned} \exp [W_t[J]] &= \int [d\phi] \exp \left[-\frac{1}{2} \int_p A_t(p)^2 \phi(p) \phi(-p) + \int_p J(-p) \phi(p) \right] \\ &\quad \times \int [d\phi'] \exp \left[S_0[\phi'] + \int_p J(p) Z_t(p) \phi'(-p) \right] \end{aligned}$$

$$= \exp \left[-\frac{1}{2} \int_p \frac{1}{A_t(p)^2} J(p) J(-p) + W_0 [Z_t(p) J(p)] \right]. \quad (\text{A}\cdot 15)$$

Hence, we obtain

$$W_t[J] = W_0[Z_t J] - \frac{1}{2} \int_p \frac{1}{A_t(p)^2} J(p) J(-p). \quad (\text{A}\cdot 16)$$

This means that nothing is lost in going from S_0 to S_t ; we can construct W_0 from W_t , and vice versa. Differentiating the above equation with respect to J , we obtain the following equation for the connected correlation functions:

$$\begin{cases} \langle \phi(p) \phi(-p) \rangle_{S_t} &= \frac{1}{A_t(p)^2} + Z_t(p)^2 \langle \phi(p) \phi(-p) \rangle_{S_0}, \\ \langle \phi(p_1) \cdots \phi(p_n) \rangle_{S_t} &= \prod_{i=1}^n Z_t(p_i) \cdot \langle \phi(p_1) \cdots \phi(p_n) \rangle_{S_0}. \end{cases} \quad (n > 1) \quad (\text{A}\cdot 17)$$

This is a generalization of (2.43). Hence, $\phi(p)$ of S_t has the same correlations as the block spin $Z_t(p)\phi(p)$ of the original S_0 . The smearing factor A_t affects only the two-point correlation function. This is a well known general characteristic of diffusion processes.*)

We now choose a particular blocking function, given in terms of a cutoff function K :

$$Z_t(p) = \frac{K(pe^t/\Lambda_0)}{K(p/\Lambda_0)} \exp \left[\frac{1}{2} \int_0^t dt' \eta(t') \right], \quad (\text{A}\cdot 18)$$

where η is an anomalous dimension, which affects only the overall normalization of the field. For $\eta = 0$, we obtain

$$Z_t(p) \begin{cases} = 1 & (p^2 < e^{-2t} \Lambda_0^2), \\ \sim 0 & (p^2 > e^{-2t} \Lambda_0^2), \end{cases} \quad (\text{A}\cdot 19)$$

if we assume $K(p/\Lambda) = 1$ for $p^2 < \Lambda^2$. Hence, the support of the Fourier transform $Z_t(x)$ has size e^t/Λ_0 in space, growing exponentially as t increases. This choice of Z_t corresponds to

$$F_t(p) \equiv -\partial_t \ln Z_t(p) = \frac{\Delta(pe^t/\Lambda_0)}{K(pe^t/\Lambda_0)} - \frac{\eta(t)}{2}. \quad (\text{A}\cdot 20)$$

With the above choice for $F_t(p)$, we obtain

$$\begin{aligned} G_t(p) &= -2 Z_t(p)^2 \partial_t \ln A_t Z_t(p) \\ &= -2 e^{\int_0^t \eta} \frac{K(pe^t/\Lambda_0)^2}{K(p/\Lambda_0)^2} \frac{\partial_t \ln A_t Z_t(p)}{(A_t Z_t(p))^2}. \end{aligned} \quad (\text{A}\cdot 21)$$

Hence,

$$\partial_t \frac{1}{(A_t Z_t(p))^2} = e^{-\int_0^t \eta} \frac{K(p/\Lambda_0)^2}{K(pe^t/\Lambda_0)^2} G_t(p), \quad (\text{A}\cdot 22)$$

*) The diffusion equation $\partial_t P(x, t) = D \partial_x^2 P(x, t)$ preserves the connected part of the correlation $\langle x^n \rangle = \int dx x^n P(x, t)$ for $n > 2$.

where we impose the initial condition

$$\lim_{t \rightarrow 0^+} \frac{1}{(A_t Z_t(p))^2} = 0. \quad (\text{A}\cdot 23)$$

Let us now look at two examples of $G_t(p)$ and calculate the corresponding $A_t Z_t(p)$.

Example 1 (Wilson)

We choose

$$G_t(p) = 2 \frac{e^{2t}}{\Lambda_0^2} \left(\frac{\Delta(pe^t/\Lambda_0)}{K(pe^t/\Lambda_0)} + 1 - \frac{1}{2}\eta(t) \right). \quad (\text{A}\cdot 24)$$

This is the original choice of Wilson discussed in sect. 11 of [124].^{*)} This gives

$$\begin{aligned} \partial_t \frac{\Lambda_0^2}{(A_t Z_t(p))^2} &= 2e^{2t-\int_0^t \eta} \frac{K(p/\Lambda_0)^2}{K(pe^t/\Lambda_0)^2} \left(\frac{\Delta(pe^t/\Lambda_0)}{K(pe^t/\Lambda_0)} + 1 - \frac{\eta}{2} \right) \\ &= \partial_t \left(e^{2t-\int_0^t \eta} \frac{K(p/\Lambda_0)^2}{K(pe^t/\Lambda_0)^2} \right). \end{aligned} \quad (\text{A}\cdot 25)$$

Hence, we obtain

$$\frac{\Lambda_0^2}{(A_t Z_t(p))^2} = e^{2t-\int_0^t \eta} \frac{K(p/\Lambda_0)^2}{K(pe^t/\Lambda_0)^2} - 1, \quad (\text{A}\cdot 26)$$

using the initial condition (A·23). Thus, using (A·18), we obtain

$$\frac{1}{A_t(p)^2} = \frac{1}{\Lambda_0^2} \left(1 - \frac{K(pe^t/\Lambda_0)^2}{K(p/\Lambda_0)^2} e^{\int_0^t \eta} \right). \quad (\text{A}\cdot 27)$$

This implies

$$\begin{cases} \langle \phi(p)\phi(-p) \rangle_{S_t} - \frac{e^{2t}}{\Lambda_0^2} &= e^{\int_0^t \eta} \frac{K(pe^t/\Lambda_0)^2}{K(p/\Lambda_0)^2} \left(\langle \phi(p)\phi(-p) \rangle_{S_0} - \frac{1}{\Lambda_0^2} \right), \\ \langle \phi(p_1) \cdots \phi(p_{2n}) \rangle_{S_t} &= e^{n \int_0^t \eta} \prod_{i=1}^{2n} \frac{K(p_i e^t/\Lambda_0)}{K(p_i/\Lambda_0)} \cdot \langle \phi(p_1) \cdots \phi(p_{2n}) \rangle_{S_0}. \end{cases} \quad (\text{A}\cdot 28)$$

Wilson's choice is characterized by the following behavior of the two-point function as $t \rightarrow \infty$:

$$\langle \phi(p)\phi(-p) \rangle_{S_t} \xrightarrow{t \rightarrow \infty} \frac{e^{2t}}{\Lambda_0^2}. \quad (\text{A}\cdot 29)$$

Example 2 (Polchinski)

The Polchinski equation (2·55) is obtained if we choose $\eta = 0$, and

$$G_t(p) = \frac{\Delta(pe^t/\Lambda_0)}{p^2 + m^2}. \quad (\text{A}\cdot 30)$$

^{*)} We have not introduced rescaling of space. This is discussed briefly in Appendix C.

Here, we make a slightly generalized choice without assuming $\eta = 0$:

$$G_t(p) = \frac{1}{p^2 + m^2(t)} \left\{ \Delta(pe^t/\Lambda_0) - \left(\eta(t) + \frac{b_m(t)}{p^2 + m^2(t)} \right) K(pe^t/\Lambda_0) (1 - K(pe^t/\Lambda_0)) \right\}, \quad (\text{A}\cdot 31)$$

where $m^2(t)$ is a t -dependent squared mass, and

$$b_m(t) \equiv \frac{dm^2(t)}{dt}. \quad (\text{A}\cdot 32)$$

Let us find the corresponding $A_t Z_t(p)$; we find

$$\begin{aligned} \partial_t \frac{1}{(A_t Z_t(p))^2} &= e^{-\int_0^t \eta} \frac{K(p/\Lambda_0)^2}{K(pe^t/\Lambda_0)^2} \frac{1}{p^2 + m^2(t)} \\ &\times \left\{ \Delta(pe^t/\Lambda_0) - \left(\eta(t) + \frac{b_m(t)}{p^2 + m^2(t)} \right) K(pe^t/\Lambda_0) (1 - K(pe^t/\Lambda_0)) \right\} \\ &= K(p/\Lambda_0)^2 \partial_t \left\{ \frac{e^{-\int_0^t \eta}}{p^2 + m^2(t)} \left(\frac{1}{K(pe^t/\Lambda_0)} - 1 \right) \right\}. \end{aligned} \quad (\text{A}\cdot 33)$$

Hence,

$$\begin{aligned} \frac{1}{(A_t Z_t(p))^2} &= K(p/\Lambda_0)^2 \\ &\cdot \left\{ \frac{e^{-\int_0^t \eta}}{p^2 + m^2(t)} \left(\frac{1}{K(pe^t/\Lambda_0)} - 1 \right) - \frac{1}{p^2 + m^2(0)} \left(\frac{1}{K(p/\Lambda_0)} - 1 \right) \right\}. \end{aligned} \quad (\text{A}\cdot 34)$$

Thus, we obtain

$$\begin{aligned} \frac{1}{A_t(p)^2} &= K(pe^t/\Lambda_0)^2 \\ &\cdot \left\{ \frac{1}{p^2 + m^2(t)} \left(\frac{1}{K(pe^t/\Lambda_0)} - 1 \right) - \frac{e^{\int_0^t \eta}}{p^2 + m^2(0)} \left(\frac{1}{K(p/\Lambda_0)} - 1 \right) \right\}. \end{aligned} \quad (\text{A}\cdot 35)$$

Hence, we obtain the following t dependence of the correlation functions:

$$\begin{aligned} \langle \phi(p) \phi(-p) \rangle_{S_t} &= \frac{K(pe^t/\Lambda_0) (1 - K(pe^t/\Lambda_0))}{p^2 + m^2(t)} \\ &= e^{\int_0^t \eta} \frac{K(pe^t/\Lambda_0)^2}{K(p/\Lambda_0)^2} \left(\langle \phi(p) \phi(-p) \rangle_{S_0} - \frac{K(p/\Lambda_0) (1 - K(p/\Lambda_0))}{p^2 + m^2(0)} \right), \end{aligned} \quad (\text{A}\cdot 36)$$

and

$$\langle \phi(p_1) \cdots \phi(p_{2n}) \rangle_{S_t} = e^{n \int_0^t \eta} \prod_{i=1}^{2n} \frac{K(p_i e^t/\Lambda_0)}{K(p_i/\Lambda_0)} \cdot \langle \phi(p_1) \cdots \phi(p_{2n}) \rangle_{S_0}. \quad (\text{A}\cdot 37)$$

These reduce to (2.43) if we take $\eta = 0$ and t -independent m^2 .

A.2. Extention to fermions

In extending the generalized diffusion equations to fermions, we only consider the spin $\frac{1}{2}$ fields in $D = 4$. We generate the following one-parameter family of actions:

$$\begin{aligned} \exp [S_t[\psi, \bar{\psi}]] &\equiv \int [d\psi' d\bar{\psi}'] \exp [S_0[\psi', \bar{\psi}'] \\ &- \int_p \{ \bar{\psi}(-p) - Z_t(p) \bar{\psi}'(-p) \} \bar{A}_t(-p) A_t(p) \{ \psi(p) - Z_t(p) \psi'(p) \}] , \end{aligned} \quad (\text{A}\cdot 38)$$

where Z_t is a scalar, and A_t & $\bar{A}_t$ are 4-by-4 matrices. We choose

$$Z_0 = 1, \quad \frac{1}{\bar{A}_0 A_0} = 0, \quad (\text{A}\cdot 39)$$

so that

$$\lim_{t \rightarrow 0+} S_t = S_0. \quad (\text{A}\cdot 40)$$

Let us define the generating functional by

$$\exp [W_t[\bar{\xi}, \eta]] \equiv \int [d\psi d\bar{\psi}] \exp \left[S_t[\psi, \bar{\psi}] + \int (\bar{\xi} \psi + \bar{\psi} \eta) \right]. \quad (\text{A}\cdot 41)$$

We then obtain the familiar relation

$$W_t[\bar{\xi}, \eta] = W_0[Z_t \bar{\xi}, Z_t \eta] + \int \bar{\xi}(-p) \frac{1}{\bar{A}_t(-p) A_t(p)} \eta(p), \quad (\text{A}\cdot 42)$$

where $1/(\bar{A}_t A_t)$ denotes the inverse of the matrix $\bar{A}_t A_t$.

To derive the differential equation for S_t , it is convenient to rewrite

$$\begin{aligned} \exp [S_t[\psi, \bar{\psi}]] &= \int [d\psi' d\bar{\psi}'] \exp \left[- \int_p \bar{\psi}'(-p) \psi'(p) \right. \\ &\quad \left. + S_0 \left[\frac{1}{Z_t} \left(\frac{1}{\bar{A}_t} \psi' + \psi \right), \frac{1}{Z_t} \left(\bar{\psi}' \frac{1}{\bar{A}_t} + \bar{\psi} \right) \right] \right]. \end{aligned} \quad (\text{A}\cdot 43)$$

From this, we obtain the following ERG differential equation:

$$\begin{aligned} \partial_t S_t &= \int_p \left[F_t(p) \left\{ S_t \frac{\overleftarrow{\delta}}{\delta \psi(p)} \psi(p) + \bar{\psi}(-p) \frac{\overrightarrow{\delta}}{\delta \bar{\psi}(-p)} S_t \right\} \right. \\ &\quad \left. - \text{Tr } G_t(p) \left\{ \frac{\overrightarrow{\delta}}{\delta \bar{\psi}(-p)} S_t \cdot S_t \frac{\overleftarrow{\delta}}{\delta \psi(p)} + \frac{\overrightarrow{\delta}}{\delta \bar{\psi}(-p)} S_t \frac{\overleftarrow{\delta}}{\delta \psi(p)} \right\} \right], \end{aligned} \quad (\text{A}\cdot 44)$$

where

$$\begin{cases} F_t(p) &\equiv -\partial_t \ln Z_t(p), \\ G_t(p) &\equiv Z_t(p)^2 \partial_t \{ 1 / (Z_t(p)^2 \bar{A}_t(-p) A_t(p)) \}. \end{cases} \quad (\text{A}\cdot 45)$$

Example 1 (Polchinski)

We take

$$Z_t(p) \equiv \frac{K (pe^t/\Lambda_0)}{K (p/\Lambda_0)} = \frac{K}{K_0}, \quad (\text{A.46a})$$

$$\frac{1}{\bar{A}_t(-p)A_t(p)} \equiv \frac{1}{\not{p} + im} K \left(1 - \frac{K}{K_0}\right). \quad (\text{A.46b})$$

Then we obtain

$$\begin{cases} F_t(p) & \equiv \Delta(pe^t/\Lambda_0)/K (pe^t/\Lambda_0), \\ G_t(p) & \equiv \Delta(pe^t/\Lambda_0)/(\not{p} + im). \end{cases} \quad (\text{A.47})$$

Example 2 (chirality breaking diffusion)

We keep the same blocking factor Z_t , but take the smearing factor

$$\frac{1}{\bar{A}_t A_t} = \frac{1}{im} K \left(1 - \frac{K}{K_0}\right) \quad (\text{A.48})$$

so that

$$G_t(p) = \frac{\Delta(pe^t/\Lambda_0)}{im}. \quad (\text{A.49})$$

The smearing factor breaks chirality explicitly. Even if we start from the chiral invariant action

$$S_0[\psi, \bar{\psi}] = - \int_p \frac{1}{K_0(p)} \bar{\psi}(-p) \not{p} \psi(p), \quad (\text{A.50})$$

the diffusion process breaks chiral symmetry explicitly, and $S_{t>0}$ is not manifestly chiral invariant. But it does inherit the chiral invariance of the original action S_0 , as will be shown at the end of the next subsection. It is straightforward to find

$$S_t[\psi, \bar{\psi}] = - \int_p \bar{\psi}(-p) D_t(p) \psi(p), \quad (\text{A.51})$$

where

$$D_t(p) \equiv im \frac{K_0}{K} \frac{\not{p}}{(K_0 - K)\not{p} + imK}. \quad (\text{A.52})$$

This satisfies the initial condition:

$$D_0(p) = \frac{\not{p}}{K_0}. \quad (\text{A.53})$$

Now, we take $\Lambda_0 \rightarrow \infty$, while we keep $\Lambda = \Lambda_0 e^{-t}$ finite. We obtain the continuum limit:

$$D_\Lambda(p) = \frac{im}{K(p/\Lambda)} \frac{\not{p}}{(1 - K(p/\Lambda))\not{p} + imK(p/\Lambda)}. \quad (\text{A.54})$$

This has an interesting property:

$$D_\Lambda(p) \longrightarrow \begin{cases} \frac{im}{K(p/\Lambda)} & (p \gg \Lambda), \\ \not{p} & (p < \Lambda). \end{cases} \quad (\text{A.55})$$

For $p^2 < \Lambda^2$, chiral invariance becomes manifest.

A.3. WT operator for chiral invariance

As an application, let us formulate the WT identity for chiral invariance. At scale Λ_0 , we define the WT operator

$$\Sigma_0[\psi, \bar{\psi}] \equiv \int_p \left(S_0 \frac{\overleftarrow{\delta}}{\delta\psi(p)} \gamma_5 \psi(p) + \bar{\psi}(-p) \gamma_5 \frac{\overrightarrow{\delta}}{\delta\bar{\psi}(-p)} S_0 \right) \quad (\text{A.56})$$

for chiral invariance (axial invariance to be more precise). $\Sigma_0 = 0$ implies that the action S_0 is invariant under the chiral transformation:

$$\psi(p) \longrightarrow \gamma_5 \psi(p), \quad \bar{\psi}(p) \longrightarrow \bar{\psi}(p) \gamma_5. \quad (\text{A.57})$$

We wish to study how the chiral symmetry is realized by the Wilson action S_t with $t > 0$.

A familiar calculation gives

$$\begin{aligned} & \int [d\psi d\bar{\psi}] \Sigma_0[\psi, \bar{\psi}] \exp \left[S_0[\psi, \bar{\psi}] + \int (\bar{\xi} \psi + \bar{\psi} \eta) \right] \\ &= \int [d\psi d\bar{\psi}] \Sigma_t[\psi, \bar{\psi}] \exp \left[S_t[\psi, \bar{\psi}] + \int \frac{1}{Z_t} (\bar{\xi} \psi + \bar{\psi} \eta) - \int \frac{1}{Z_t^2} \bar{\xi} \frac{1}{\bar{A}_t A_t} \eta \right], \end{aligned} \quad (\text{A.58})$$

where

$$\begin{aligned} \Sigma_t \equiv & - \int_p \text{Tr} \gamma_5 \left[\psi(p) \cdot S_t \frac{\overleftarrow{\delta}}{\delta\psi(p)} + \frac{\overrightarrow{\delta}}{\delta\bar{\psi}(-p)} S_t \cdot \bar{\psi}(-p) \right] \\ & - \int_p \text{Tr} \left\{ \gamma_5, \frac{1}{\bar{A}_t A_t} \right\} \left[\frac{\overrightarrow{\delta}}{\delta\bar{\psi}(-p)} S_t \cdot S_t \frac{\overleftarrow{\delta}}{\delta\psi(p)} + \frac{\overrightarrow{\delta}}{\delta\bar{\psi}(-p)} S_t \frac{\overleftarrow{\delta}}{\delta\psi(p)} \right]. \end{aligned} \quad (\text{A.59})$$

Hence, the chiral invariance of S_t is given by

$$\Sigma_t = 0. \quad (\text{A.60})$$

For a free theory^{*)}

$$S_t = - \int_p \bar{\psi}(-p) D_t(p) \psi(p), \quad (\text{A.61})$$

the relation $\Sigma_t = 0$ reduces to the Ginsparg-Wilson relation⁴⁷⁾

$$\{\gamma_5, D_t(p)\} - D_t(p) \left\{ \gamma_5, \frac{1}{\bar{A}_t A_t} \right\} D_t(p) = 0. \quad (\text{A.62})$$

ERG gives the following t -dependence:

$$D_t(p) = \bar{A}_t A_t - \bar{A}_t A_t Z_t \frac{1}{D_0 + Z_t^2 \bar{A}_t A_t} Z_t \bar{A}_t A_t. \quad (\text{A.63})$$

^{*)} The quantum master equation and its solutions have been studied for interacting fermions in Refs. 64), 68), 69). See also Ref. 58)

Substituting this into the Ginsparg-Wilson relation above, we obtain what is expected:

$$\{\gamma_5, D_0\} = 0, \quad (\text{A}\cdot 64)$$

which is equivalent to

$$\Sigma_0 = 0. \quad (\text{A}\cdot 65)$$

Let us reexamine the second example of the previous subsection, where we have chosen the blocking factor (A·46a) and the chirality breaking smearing factor (A·48):

$$Z_t(p) = \frac{K}{K_0}, \quad \frac{1}{A_t A_t} = \frac{1}{im} K \left(1 - \frac{K}{K_0} \right). \quad (\text{A}\cdot 66)$$

(A·59) gives

$$\begin{aligned} \Sigma_t \equiv & - \int_p \text{Tr} \gamma_5 \left[\psi(p) \cdot S_t \frac{\overleftarrow{\delta}}{\delta \psi(p)} + \frac{\overrightarrow{\delta}}{\delta \bar{\psi}(-p)} S_t \cdot \bar{\psi}(-p) \right. \\ & \left. + \frac{2K_0}{imK} (K_0 - K) \left\{ \frac{\overrightarrow{\delta}}{\delta \bar{\psi}(-p)} S_t \cdot S_t \frac{\overleftarrow{\delta}}{\delta \psi(p)} + \frac{\overrightarrow{\delta}}{\delta \bar{\psi}(-p)} S_t \frac{\overleftarrow{\delta}}{\delta \psi(p)} \right\} \right]. \end{aligned} \quad (\text{A}\cdot 67)$$

The free action

$$S_t[\psi, \bar{\psi}] = - \int_p \bar{\psi}(-p) im \frac{K_0}{K} \frac{\not{p}}{(K_0 - K)\not{p} + imK} \psi(p), \quad (\text{A}\cdot 68)$$

given by (A·51) and (A·52), indeed satisfies the WT identity:

$$\Sigma_t = 0. \quad (\text{A}\cdot 69)$$

Appendix B

— Applications of composite operators —

In the main text we have shown how to realize symmetry in terms of either a WT composite operator Σ_A or a QM composite operator $\bar{\Sigma}_A$. In this appendix we describe two further applications of composite operators: one to beta functions, and the other to universality.

B.1. Beta functions

We recall that the Wilson action of the ϕ^4 theory in $D = 4$ is defined by the Polchinski equation

$$-\Lambda \frac{\partial}{\partial \Lambda} S_{I,\Lambda}[\phi] = \int_p \frac{\Delta(p/\Lambda)}{p^2 + m^2} \frac{1}{2} \left\{ \frac{\delta S_{I,\Lambda}[\phi]}{\delta \phi(-p)} \frac{\delta S_{I,\Lambda}[\phi]}{\delta \phi(p)} + \frac{\delta^2 S_{I,\Lambda}[\phi]}{\delta \phi(-p) \delta \phi(p)} \right\}, \quad (\text{B}\cdot 1)$$

and the asymptotic behavior

$$S_{I,\Lambda} \xrightarrow{\Lambda \rightarrow \infty} \int d^4x \left[(\Lambda^2 a_2(\ln \Lambda/\mu) + m^2 b_2(\ln \Lambda/\mu)) \frac{\phi^2}{2} + c_2(\ln \Lambda/\mu) \frac{1}{2} (\partial_\mu \phi)^2 + a_4(\ln \Lambda/\mu) \frac{\phi^4}{4!} \right]. \quad (\text{B}\cdot 2)$$

The Wilson action is parametrized not only by m^2 which appears in the differential equation, but also by the three parameters $b_2(0), c_2(0), a_4(0)$. Here, for the sake of simplicity, we make a simple choice:¹⁰⁶⁾

$$b_2(0) = c_2(0) = 0, \quad a_4(0) = -\lambda. \quad (\text{B}\cdot 3)$$

We call this the MS scheme for its similarity to the minimal subtraction scheme for dimensional regularization.^{111), 112)} We immediately notice, though, that the MS scheme depends on the choice of μ . The MS scheme (B·3) for a different μ corresponds to $b_2(0), c_2(0), a_4(0)$ that do not satisfy (B·3) for the original μ . It is the purpose of this appendix to derive the μ dependence of the Wilson action in the MS scheme (B·3).¹⁰⁸⁾ (See also 31), 57) for derivations of beta functions from the ERG. Ref. 91) derives the same mass independent RG equation as (B·22).)

Since μ can be regarded as a constant external field, the derivative of the Wilson action

$$-\mu \frac{\partial S_\Lambda}{\partial \mu} \quad (\text{B}\cdot 4)$$

is a composite operator. (Here, m^2 and λ are fixed.) To be more precise, it is a dimension 4 composite operator with zero momentum. Hence, it must be a linear combination of the three linearly independent composite operators

$$m^2 \left[\frac{\phi^2}{2} \right]_A(0), \quad \left[\frac{\phi^4}{4!} \right]_A(0), \quad \left[\frac{1}{2} (\partial_\mu \phi)^2 \right]_A(0), \quad (\text{B}\cdot 5)$$

which are defined at the end of §4.3. The asymptotic behavior (B·2) implies the following asymptotic behavior:

$$-\mu \frac{\partial S_\Lambda}{\partial \mu} \xrightarrow{\Lambda \rightarrow \infty} \int d^4x \left[\left(\Lambda^2 \dot{a}_2(\ln \Lambda/\mu) + m^2 \dot{b}_2(\ln \Lambda/\mu) \right) \frac{\phi^2}{2} + \dot{c}_2(\ln \Lambda/\mu) \frac{1}{2} (\partial_\mu \phi)^2 + \dot{a}_4(\ln \Lambda/\mu) \frac{\phi^4}{4!} \right], \quad (\text{B} \cdot 6)$$

where

$$\dot{a}_2(\ln \Lambda/\mu) \equiv \Lambda \frac{\partial}{\partial \Lambda} a_2(\ln \Lambda/\mu), \text{ etc.} \quad (\text{B} \cdot 7)$$

Hence, we obtain

$$-\mu \frac{\partial S_\Lambda}{\partial \mu} = \dot{b}_2(0) m^2 \left[\frac{\phi^2}{2} \right]_\Lambda (0) + \dot{c}_2(0) \left[\frac{1}{2} (\partial_\mu \phi)^2 \right]_\Lambda (0) + \dot{a}_4(0) \left[\frac{\phi^4}{4!} \right]_\Lambda (0), \quad (\text{B} \cdot 8)$$

where the coefficients depend only on λ .

We now wish to rewrite the above equation as a physically meaningful equation. For this purpose, we construct an alternative basis of dimension 4 composite operators with zero momentum.

We first consider

$$\mathcal{O}_\lambda \equiv -\partial_\lambda S_\Lambda = -\partial_\lambda S_{I,\Lambda}. \quad (\text{B} \cdot 9)$$

This is a composite operator of the type (4·34), since λ can be regarded as a constant external field. This satisfies

$$\begin{aligned} \langle \mathcal{O}_\lambda \phi(p_1) \cdots \phi(p_n) \rangle^\infty &\equiv \prod_{i=1}^n \frac{1}{K(p_i/\Lambda)} \cdot \langle \mathcal{O}_\lambda \phi(p_1) \cdots \phi(p_n) \rangle_{S_\Lambda} \\ &= -\partial_\lambda \langle \phi(p_1) \cdots \phi(p_n) \rangle^\infty. \end{aligned} \quad (\text{B} \cdot 10)$$

Similarly, we wish to construct a composite operator $\mathcal{O}_m$ that gives

$$\langle \mathcal{O}_m \phi(p_1) \cdots \phi(p_n) \rangle^\infty = -\partial_{m^2} \langle \phi(p_1) \cdots \phi(p_n) \rangle^\infty \quad (\text{B} \cdot 11)$$

for any n . For $n \neq 2$, the m^2 derivative of S_Λ will do:

$$-\partial_{m^2} \langle \phi(p_1) \cdots \phi(p_n) \rangle^\infty = \prod_{i=1}^n \frac{1}{K(p_i/\Lambda)} \cdot \langle (-\partial_{m^2} S_\Lambda) \phi(p_1) \cdots \phi(p_n) \rangle_{S_\Lambda}. \quad (\text{B} \cdot 12)$$

But, for $n = 2$, we obtain

$$\begin{aligned} &-\partial_{m^2} \langle \phi(p) \phi(-p) \rangle^\infty \\ &= -\partial_{m^2} \left(\frac{1}{K^2} \langle \phi(p) \phi(-p) \rangle_{S_\Lambda} + \frac{1 - 1/K}{p^2 + m^2} \right) \\ &= \frac{1}{K^2} \left(\langle (-\partial_{m^2} S_\Lambda) \phi(p) \phi(-p) \rangle_{S_\Lambda} + \frac{K(K-1)}{(p^2 + m^2)^2} \right). \end{aligned} \quad (\text{B} \cdot 13)$$

Hence, we need $\mathcal{O}'$ that satisfies

$$\begin{cases} \langle \mathcal{O}' \phi(p) \phi(-p) \rangle_{S_\Lambda} = \frac{K(K-1)}{(p^2+m^2)^2}, \\ \langle \mathcal{O}' \phi(p_1) \cdots \phi(p_n) \rangle_{S_\Lambda} = 0 \quad (n \neq 2). \end{cases} \quad (\text{B}\cdot 14)$$

This is easily constructed as

$$\mathcal{O}' = \int_p \frac{K(K-1)}{(p^2+m^2)^2} \frac{1}{2} \left\{ \frac{\delta S_\Lambda}{\delta \phi(p)} \frac{\delta S_\Lambda}{\delta \phi(-p)} + \frac{\delta^2 S_\Lambda}{\delta \phi(p) \delta \phi(-p)} \right\}. \quad (\text{B}\cdot 15)$$

Hence, we obtain

$$\begin{aligned} \mathcal{O}_m &= -\partial_{m^2} S_\Lambda + \mathcal{O}' \\ &= -\partial_{m^2} S_\Lambda - \int_p \frac{K(1-K)}{(p^2+m^2)^2} \frac{1}{2} \left\{ \frac{\delta S_\Lambda}{\delta \phi(p)} \frac{\delta S_\Lambda}{\delta \phi(-p)} + \frac{\delta^2 S_\Lambda}{\delta \phi(p) \delta \phi(-p)} \right\}. \end{aligned} \quad (\text{B}\cdot 16)$$

It is straightforward to check that this $\mathcal{O}_m$ satisfies the ERG differential equation (4.3).

The third and last composite operator is defined by

$$\mathcal{N} \equiv - \int_p K(p/\Lambda) \left\{ [\phi]_\Lambda(p) \frac{\delta S_\Lambda}{\delta \phi(p)} + \frac{\delta}{\delta \phi(p)} [\phi]_\Lambda(p) \right\}, \quad (\text{B}\cdot 17)$$

where

$$[\phi]_\Lambda(p) \equiv \frac{1}{K} \phi(p) + \frac{1-K}{p^2+m^2} \frac{\delta S_\Lambda}{\delta \phi(-p)} \quad (\text{B}\cdot 18)$$

is the composite operator corresponding to $\phi(p)$. $\mathcal{N}$ is a composite operator of the type (4.32), and it has a simple correlation function

$$\langle \mathcal{N} \phi(p_1) \cdots \phi(p_n) \rangle^\infty = n \langle \phi(p_1) \cdots \phi(p_n) \rangle^\infty \quad (\text{B}\cdot 19)$$

for any n .

Thus, we have obtained an alternative basis of dimension 4 scalar composite operators with zero momentum, consisting of

$$\mathcal{O}_\lambda, \quad m^2 \mathcal{O}_m, \quad \mathcal{N}. \quad (\text{B}\cdot 20)$$

We can use this basis, instead of the basis consisting of (B.5), to expand $-\mu \frac{\partial}{\partial \mu} S_\Lambda$:

$$-\mu \frac{\partial S_\Lambda}{\partial \mu} = \beta_m(\lambda) m^2 \mathcal{O}_m + \beta(\lambda) \mathcal{O}_\lambda + \gamma(\lambda) \mathcal{N}, \quad (\text{B}\cdot 21)$$

where the coefficients are all functions of λ alone, since they cannot depend on $\ln \Lambda/\mu$. The physical meaning of this equation is clear. It implies, for the correlation functions,

$$\left(-\mu \frac{\partial}{\partial \mu} + \beta_m m^2 \partial_{m^2} + \beta \partial_\lambda - n \gamma \right) \langle \phi(p_1) \cdots \phi(p_n) \rangle^\infty = 0. \quad (\text{B}\cdot 22)$$

This is a mass independent renormalization group equation.^{(111),(118)}

B.2. Universality

Universality is an important concept in renormalization theory. To construct a continuum limit, there are always more than one way. The independence of the continuum limit on the particular method of construction is called universality. For example, if we use a lattice to construct a continuum theory, the limit should not depend on what kind of lattice, whether square or cubic, we use.

In the following we examine universality in two restricted senses. First, we wish to show that the continuum limit (i.e., the correlation functions with the suffix ∞) does not depend on the particular asymptotic conditions we use to select a solution of the ERG differential equation. Second, we wish to show that the continuum limit does not depend on the choice of a cutoff function K .

Scheme dependence

Let us compare two solutions of the same Polchinski differential equation (B.1). One is S_A satisfying the MS condition (B.3), and the other is $S_A + \delta S_A$ satisfying

$$\begin{cases} b_2(0) = \delta b_2, \\ c_2(0) = \delta c_2, \\ a_4(0) = -\lambda + \delta a_4, \end{cases} \quad (\text{B.23})$$

where $\delta b_2, \delta c_2, \delta a_4$ are all infinitesimal constants. Then, δS_A is a composite operator given by

$$\delta S_A = \delta b_2 m^2 \left[\frac{\phi^2}{2} \right]_A(0) + \delta c_2 \left[\frac{1}{2} (\partial_\mu \phi)^2 \right]_A(0) + \delta a_4 \left[\frac{\phi^4}{4!} \right]_A(0). \quad (\text{B.24})$$

Using the alternative basis, we can rewrite this in the form

$$\delta m^2 \mathcal{O}_m + \frac{\delta z}{2} \mathcal{N} + \delta \lambda \mathcal{O}_\lambda. \quad (\text{B.25})$$

This implies that $S_A + \delta S_A$, which does not satisfy the MS condition (B.3), gives the same continuum limit as the Wilson action satisfying (B.3) with squared mass $m^2 + \delta m^2$ and coupling $\lambda + \delta \lambda$. The only difference is in the normalization of fields by a factor $\frac{1}{2}\delta z$. Extending this, we can show that the action with any choice of $b_2(0), c_2(0), a_4(0)$ is equivalent to an action satisfying the MS condition (B.3) up to field normalization.

Dependence on the choice of K

In our discussions so far, we have always kept a choice of the cutoff function K . Physics should not depend on the choice of K , and we will show this in the following.

Let S_A be a solution of the ERG equation with the cutoff function $K + \delta K$, infinitesimally different from K . We would like to construct an equivalent action S'_A that has K as the cutoff function. We first recall that the correlation functions in the continuum limit are given by

$$\begin{cases} \langle \phi(p)\phi(-p) \rangle^\infty = \frac{1}{(K+\delta K)^2} \langle \phi(p)\phi(-p) \rangle_{S_A} + \frac{1-1/(K+\delta K)}{p^2+m^2}, \\ \langle \phi(p_1) \cdots \phi(p_n) \rangle^\infty = \prod_{i=1}^n \frac{1}{(K+\delta K)(p_i)} \cdot \langle \phi(p_1) \cdots \phi(p_n) \rangle_{S_A}. \end{cases} \quad (\text{B.26})$$

This can be rewritten as

$$\langle \phi(p)\phi(-p) \rangle^\infty = \frac{1}{K^2} \left\{ \langle (1 - \delta K/K) \phi(p) (1 - \delta K/K) \phi(-p) \rangle_{S_\Lambda} + \frac{\delta K}{p^2 + m^2} \right\} + \frac{1 - 1/K}{p^2 + m^2}, \quad (\text{B}\cdot 27)$$

$$\langle \phi(p_1) \cdots \phi(p_n) \rangle^\infty = \prod_{i=1}^n \frac{1}{K(p_i)} \times \langle (1 - \delta K/K) \phi(p_1) \cdots (1 - \delta K/K) \phi(p_n) \rangle_{S_\Lambda}. \quad (\text{B}\cdot 28)$$

We now construct a new Wilson action by

$$S'_\Lambda \equiv S_\Lambda + \int_p \frac{\delta K}{K} \phi(p) \frac{\delta S_\Lambda}{\delta \phi(p)} + \int_p \frac{\delta K}{p^2 + m^2} \frac{1}{2} \left\{ \frac{\delta S_\Lambda}{\delta \phi(p)} \frac{\delta S_\Lambda}{\delta \phi(-p)} + \frac{\delta^2 S_\Lambda}{\delta \phi(p) \delta \phi(-p)} \right\}. \quad (\text{B}\cdot 29)$$

This has the correlation functions:

$$\begin{cases} \langle \phi(p)\phi(-p) \rangle_{S'_\Lambda} &= \langle (1 - \delta K/K) \phi(p) (1 - \delta K/K) \phi(-p) \rangle_{S_\Lambda} + \frac{\delta K}{p^2 + m^2}, \\ \langle \phi(p_1) \cdots \phi(p_n) \rangle_{S'_\Lambda} &= \langle (1 - \delta K/K) \phi(p_1) \cdots (1 - \delta K/K) \phi(p_n) \rangle_{S_\Lambda}, \end{cases} \quad (\text{B}\cdot 30)$$

so that

$$\langle \phi(p)\phi(-p) \rangle^\infty = \frac{1}{K^2} \langle \phi(p)\phi(-p) \rangle_{S'_\Lambda} + \frac{1 - 1/K}{p^2 + m^2}, \quad (\text{B}\cdot 31)$$

$$\langle \phi(p_1) \cdots \phi(p_n) \rangle^\infty = \prod_{i=1}^n \frac{1}{K(p_i)} \cdot \langle \phi(p_1) \cdots \phi(p_n) \rangle_{S'_\Lambda}. \quad (\text{B}\cdot 32)$$

This implies

1. that S'_Λ satisfies the Polchinski equation with the cutoff function K ,
2. that S'_Λ gives the same continuum limit as S_Λ .

Hence, the continuum limit does not depend on the choice of a cutoff function. Note that even if S_Λ obeys the MS condition (B·3), S'_Λ does not necessarily obey it.

Appendix C

— ERG differential equations for fixed points —

In the main text we use the Polchinski differential equation, (2·51) for $S_{I,\Lambda}$ and (2·55) for S_Λ , to construct the continuum limit of renormalizable theories. This equation is **NOT** what was originally proposed by K. G. Wilson for his non-perturbative studies of field theory.¹²⁴⁾ As we lower the cutoff Λ , the cutoff function $K(p/\Lambda)$ keeps changing its momentum dependence, and hence the Polchinski equation has no fixed point. In this appendix we explain how to modify the Polchinski equation to obtain an ERG differential equation that can have non-trivial fixed points.

To have a fixed point, we must do two things:

1. make everything dimensionless by multiplying an appropriate power of the cutoff Λ ,
2. rescale the field to normalize the kinetic term.

Let us explain the above one by one:

1. For the real scalar theory in D dimensions, the Fourier transform $\phi(p)$ has mass dimension $-\frac{D+2}{2}$. Using the dimensionless ratio

$$\bar{p} \equiv \frac{p}{\Lambda}, \quad (\text{C} \cdot 1)$$

we define the dimensionless field

$$\bar{\phi}(\bar{p}) \equiv \Lambda^{\frac{D+2}{2}} \phi(p). \quad (\text{C} \cdot 2)$$

We then rewrite the Wilson action

$$\begin{aligned} S_\Lambda[\phi] = & -\frac{1}{2} \int_p \frac{p^2}{K(p/\Lambda)} \phi(p) \phi(-p) + \sum_{n=1}^{\infty} \frac{1}{(2n)!} \int_{p_1, \dots, p_{2n}} \phi(p_1) \cdots \phi(p_{2n}) \\ & \times (2\pi)^D \delta^{(D)}(p_1 + \cdots p_{2n}) \mathcal{V}_{2n}(\Lambda; p_1, \dots, p_{2n}) \end{aligned} \quad (\text{C} \cdot 3)$$

as

$$\begin{aligned} \bar{S}_t[\bar{\phi}] \equiv & -\frac{1}{2} \int_{\bar{p}} \frac{\bar{p}^2}{K(\bar{p})} \bar{\phi}(-\bar{p}) \bar{\phi}(\bar{p}) + \sum_{n=1}^{\infty} \frac{1}{(2n)!} \int_{\bar{p}_1, \dots, \bar{p}_{2n}} \bar{\phi}(\bar{p}_1) \cdots \bar{\phi}(\bar{p}_{2n}) \\ & \times (2\pi)^D \delta^{(D)}(\bar{p}_1 + \cdots \bar{p}_{2n}) \bar{\mathcal{V}}_{2n}(t; \bar{p}_1, \dots, \bar{p}_{2n}), \end{aligned} \quad (\text{C} \cdot 4)$$

where we define

$$\bar{\mathcal{V}}_{2n}(t; \bar{p}_1, \dots, \bar{p}_{2n}) \equiv \Lambda^{(n-1)D-2n} \mathcal{V}_{2n}(\Lambda; p_1, \dots, p_{2n}) \quad (\text{C} \cdot 5)$$

so that

$$S_\Lambda[\phi] = \bar{S}_t[\bar{\phi}]. \quad (\text{C} \cdot 6)$$

The parameter t is a dimensionless parameter such that

$$\Lambda \propto e^{-t}. \quad (\text{C} \cdot 7)$$

If $S_A[\phi]$ satisfies the Polchinski equation, then $\bar{S}_t$ satisfies

$$\begin{aligned} \partial_t \bar{S}_t = & \int_p \left\{ p_\mu \frac{\partial \bar{\phi}(p)}{\partial p_\mu} + \left(\frac{D+2}{2} + \frac{\Delta(p)}{K(p)} \right) \phi(p) \right\} \frac{\delta \bar{S}_t}{\delta \bar{\phi}(p)} \\ & + \int_p \frac{\Delta(p)}{p^2} \frac{1}{2} \left(\frac{\delta \bar{S}_t}{\delta \bar{\phi}(p)} \frac{\delta \bar{S}_t}{\delta \bar{\phi}(-p)} + \frac{\delta^2 \bar{S}_t}{\delta \bar{\phi}(p) \delta \bar{\phi}(-p)} \right), \end{aligned} \quad (\text{C}\cdot 8)$$

where we have omitted the bar above the integration variable $\bar{p}$.

2. To obtain a fixed point, we must normalize the field to satisfy

$$\left. \frac{\partial}{\partial p^2} \bar{\mathcal{V}}_2(t; p, -p) \right|_{p^2=0} = 0 \quad (\text{C}\cdot 9)$$

This further modifies the differential equation to

$$\begin{aligned} \partial_t \bar{S}_t = & \int_p \left\{ p_\mu \frac{\partial \bar{\phi}(p)}{\partial p_\mu} + \left(\frac{D+2}{2} - \frac{\eta_t}{2} + \frac{\Delta(p)}{K(p)} \right) \bar{\phi}(p) \right\} \frac{\delta \bar{S}_t}{\delta \bar{\phi}(p)} \\ & + \int_p \frac{1}{p^2} \{ \Delta(p) - \eta_t K(p)(1 - K(p)) \} \frac{1}{2} \left(\frac{\delta \bar{S}_t}{\delta \bar{\phi}(p)} \frac{\delta \bar{S}_t}{\delta \bar{\phi}(-p)} + \frac{\delta^2 \bar{S}_t}{\delta \bar{\phi}(p) \delta \bar{\phi}(-p)} \right), \end{aligned} \quad (\text{C}\cdot 10)$$

where the constant η_t , **anomalous dimension**, is determined as

$$\eta_t = \frac{-\frac{\partial}{\partial p^2} \int_q \frac{\Delta(q)}{q^2} \bar{\mathcal{V}}_4(t; q, -q, p, -p) \Big|_{p^2=0}}{1 - \frac{\partial}{\partial p^2} \int_q \frac{K(q)(1-K(q))}{q^2} \bar{\mathcal{V}}_4(t; q, -q, p, -p) \Big|_{p^2=0}}. \quad (\text{C}\cdot 11)$$

The last ERG differential equation (C·10) has not only the gaussian fixed point

$$\bar{S}_G[\bar{\phi}] \equiv -\frac{1}{2} \int_p \bar{\phi}(-p) \bar{\phi}(p) \frac{p^2}{K(p)}, \quad (\text{C}\cdot 12)$$

but also a non-trivial fixed point, called the Wilson-Fisher fixed point, if $D < 4$.

In the above we have modified the ERG differential equation not only to have a fixed point, but also to give simple t -dependence to the correlation functions, similar to (2·43). We find the following results:

$$\begin{aligned} & \frac{1}{K(pe^t)^2} \langle \bar{\phi}(pe^t) \bar{\phi}(-pe^t) \rangle_{\bar{S}_t} + \frac{1 - 1/K(pe^t)}{p^2 e^{2t}} \\ & = \exp \left[-2t + \int^t dt' \eta_{t'} \right] \cdot (t\text{-independent}), \end{aligned} \quad (\text{C}\cdot 13a)$$

$$\begin{aligned} & \prod_{i=1}^n \frac{1}{K(p + ie^t)} \cdot \langle \bar{\phi}(p_1 e^t) \cdots \bar{\phi}(p_n e^t) \rangle_{\bar{S}_t} \\ & = \exp \left[t \left(D - n \frac{D+2}{2} \right) + \frac{n}{2} \int^t dt' \eta_{t'} \right] \cdot (t\text{-independent}). \end{aligned} \quad (\text{C}\cdot 13b)$$

The original differential equation, given in 124), is based upon Wilson's ERG differential equation (Example 1 of §A.1), and is somewhat different from the above:

$$\begin{aligned} \partial_t \bar{S}_t = & \int_{\bar{p}} \left\{ p_\mu \frac{\partial \bar{\phi}(p)}{\partial p_\mu} + \left(\frac{D+2}{2} - \frac{\eta_t}{2} + \frac{\Delta(p)}{K(p)} \right) \phi(p) \right\} \frac{\delta \bar{S}_t}{\delta \bar{\phi}(p)} \\ & + \int_p \left\{ \frac{\Delta(p)}{K(p)} + 1 - \frac{\eta_t}{2} \right\} \frac{1}{2} \left(\frac{\delta \bar{S}_t}{\delta \bar{\phi}(p)} \frac{\delta \bar{S}_t}{\delta \bar{\phi}(-p)} + \frac{\delta^2 \bar{S}_t}{\delta \bar{\phi}(p) \delta \bar{\phi}(-p)} \right). \end{aligned} \quad (\text{C}\cdot 14)$$

The difference in the second line comes from the choice of G_t (A·24). In 124), the function $\Delta(p)/K(p)$ is denoted as

$$\rho(p). \quad (\text{C}\cdot 15)$$

Expanding $\bar{S}_t$ in powers of fields

$$\bar{S}_t = - \sum_{n=1}^{\infty} \frac{1}{(2n)!} \int \bar{\phi}(p_1) \cdots \bar{\phi}(p_{2n}) (2\pi)^D \delta^{(D)}(p_1 + \cdots + p_{2n}) u_{2n}(t; p_1, \dots, p_{2n}), \quad (\text{C}\cdot 16)$$

the anomalous dimension η_t is determined by the normalization condition

$$\left. \frac{\partial}{\partial p^2} u_2(t; p, -p) \right|_{p^2=0} = 1 \quad (\text{C}\cdot 17)$$

as

$$\frac{\eta_t}{2} = \frac{-2u_2(t; 0, 0) + \frac{\partial}{\partial p^2} \int_q \left(\frac{\Delta(q)}{K(q)} + 1 \right) \frac{1}{2} u_4(t; q, -q, p, -p) \Big|_{p^2=0}}{1 - 2u_2(t; 0, 0) + \frac{\partial}{\partial p^2} \frac{1}{2} \int_q u_4(t; q, -q, p, -p) \Big|_{p^2=0}} \quad (\text{C}\cdot 18)$$

(C·14) gives slightly different t -dependence to the two-point correlation function:

$$\begin{aligned} & \frac{1}{K(pe^t)^2} \left(\langle \bar{\phi}(pe^t) \bar{\phi}(-pe^t) \rangle_{\bar{S}_t} - 1 \right) \\ & = \exp \left[-2t + \int^t dt' \eta_{t'} \right] \cdot (t\text{-independent}), \end{aligned} \quad (\text{C}\cdot 19a)$$

$$\begin{aligned} & \prod_{i=1}^n \frac{1}{K(p_i e^t)} \cdot \langle \bar{\phi}(p_1 e^t) \cdots \bar{\phi}(p_n e^t) \rangle_{\bar{S}_t} \\ & = \exp \left[t \left(D - n \frac{D+2}{2} \right) + \frac{n}{2} \int^t dt' \eta_{t'} \right] \cdot (t\text{-independent}). \end{aligned} \quad (\text{C}\cdot 19b)$$

Appendix D

— Symmetry of the effective average action —

The purpose of this appendix is to rewrite the WT identity and quantum master equation of the Wilson action as those of the corresponding effective average action. We have briefly discussed this in §10.5, where a more functional method is used than the method here. Our main tool is the relation between a composite operator and its 1PI sibling, given by (4.45) & (4.46).

We first consider a generic WT composite operator

$$\Sigma_\Lambda[\phi] \equiv \int_p K(p/\Lambda) \left[\mathcal{O}_\Lambda[\phi](p) \frac{\delta S_\Lambda}{\delta \phi(p)} + \frac{\delta}{\delta \phi(p)} \mathcal{O}_\Lambda[\phi](p) \right], \quad (\text{D}\cdot 1)$$

where $\mathcal{O}_\Lambda[\phi](p)$ is a composite operator of momentum p giving an infinitesimal transformation of $\phi(p)$. The corresponding 1PI composite operator is defined by

$$\Sigma_\Lambda^{1PI}[\Phi] \equiv \Sigma_\Lambda[\phi], \quad (\text{D}\cdot 2)$$

where

$$\Phi(p) \equiv \frac{1}{K(p/\Lambda)} \phi(p) + \frac{1 - K(p/\Lambda)}{p^2 + m^2} \frac{\delta S_\Lambda}{\delta \phi(-p)}. \quad (\text{D}\cdot 3)$$

(In this appendix, we consider only the continuum limit $\Lambda_0 \rightarrow \infty$.) Since Φ and ϕ coincide asymptotically, $\Sigma_\Lambda[\phi]$ and $\Sigma_\Lambda^{1PI}[\Phi]$ share the same asymptotic behavior.

In order to express $\Sigma_\Lambda[\phi]$ in terms of Φ , we consider the two composite operators $\mathcal{O}_\Lambda[\phi]$ and $K(p) \frac{\delta S_\Lambda}{\delta \phi(p)}$ one by one.

(i) $\mathcal{O}_\Lambda[\phi]$ — The corresponding 1PI composite operator is given by

$$\mathcal{O}_\Lambda^{1PI}[\Phi] = \mathcal{O}_\Lambda[\phi]. \quad (\text{D}\cdot 4)$$

(ii) $K(p) \frac{\delta S_\Lambda}{\delta \phi(p)}$ — Writing Γ_Λ for $\Gamma_{B,\Lambda}$ for short, (2.85) and (2.86) give

$$\Phi(p) = -\frac{1}{K} \frac{1 - K}{p^2 + m^2} \frac{\delta \Gamma_\Lambda[\Phi]}{\delta \Phi(-p)} + \frac{1 - K}{p^2 + m^2} \frac{\delta S_\Lambda[\phi]}{\delta \phi(-p)}. \quad (\text{D}\cdot 5)$$

Hence,

$$K(p/\Lambda) \frac{\delta S_\Lambda}{\delta \phi(-p)} = -(p^2 + m^2) \Phi(p) + \frac{\delta \Gamma_{I,\Lambda}[\Phi]}{\delta \Phi(-p)}, \quad (\text{D}\cdot 6)$$

where the interaction part $\Gamma_{I,\Lambda}[\Phi]$ is defined so that

$$\Gamma_\Lambda[\Phi] = -\frac{1}{2} \int_p \frac{p^2 + m^2}{1 - K(p/\Lambda)} \Phi(-p) \Phi(p) + \Gamma_{I,\Lambda}[\Phi]. \quad (\text{D}\cdot 7)$$

Thus, we obtain

$$\begin{aligned} \Sigma_\Lambda^{1PI}[\Phi] &= \int_p \left[\left\{ -(p^2 + m^2) \Phi(-p) + \frac{\delta \Gamma_{I,\Lambda}[\Phi]}{\delta \Phi(p)} \right\} \mathcal{O}_\Lambda^{1PI}[\Phi](p) \right. \\ &\quad \left. + K(p/\Lambda) \int_q \frac{\delta \Phi(q)}{\delta \phi(p)} \frac{\delta \mathcal{O}_\Lambda^{1PI}[\Phi](p)}{\delta \Phi(q)} \right] \end{aligned}$$

$$\begin{aligned}
&= \int_p \left[\left\{ -(p^2 + m^2)\Phi(-p) + \frac{\delta\Gamma_{I,\Lambda}[\Phi]}{\delta\Phi(p)} \right\} \mathcal{O}_\Lambda^{1PI}[\Phi](p) \right. \\
&\quad \left. - \int_q \left(\Gamma_\Lambda^{(2)} \right)^{-1}(q, -p) R_\Lambda(p) \frac{\delta\mathcal{O}_\Lambda^{1PI}[\Phi](p)}{\delta\Phi(q)} \right], \quad (\text{D}\cdot 8)
\end{aligned}$$

where we have used (2.97)

$$\frac{\delta\Phi(q)}{\delta\phi(p)} = - \left(\Gamma_\Lambda^{(2)} \right)^{-1}(q, -p) \frac{p^2 + m^2}{1 - K(p/\Lambda)}, \quad (\text{D}\cdot 9)$$

and

$$R_\Lambda(p) \equiv (p^2 + m^2) \frac{K(p/\Lambda)}{1 - K(p/\Lambda)}. \quad (\text{D}\cdot 10)$$

For example, the WT identity for QED

$$\begin{aligned}
k_\mu \frac{\delta S_{I,\Lambda}}{\delta A_\mu(k)} &= e \int_p K(p/\Lambda) \left(-S_\Lambda \frac{\overleftarrow{\delta}}{\delta\psi(p)} [\psi](p-k) + \text{Tr} [\psi](p-k) \frac{\overleftarrow{\delta}}{\delta\psi(p)} \right. \\
&\quad \left. + [\bar{\psi}](-p-k) \frac{\overrightarrow{\delta}}{\delta\bar{\psi}(-p)} S_\Lambda - \text{Tr} \frac{\overrightarrow{\delta}}{\delta\bar{\psi}(-p)} [\bar{\psi}](-p-k) \right) \quad (\text{D}\cdot 11)
\end{aligned}$$

is rewritten as

$$\begin{aligned}
k_\mu \frac{\delta\Gamma_{I,\Lambda}}{\delta\mathcal{A}_\mu(k)} &= e \int_p \left[\bar{\Psi}(-p-k) \not{k} \Psi(p) \right. \\
&\quad - \Gamma_{I,\Lambda} \frac{\overleftarrow{\delta}}{\delta\Psi(p+k)} \Psi(p) + \bar{\Psi}(-p-k) \frac{\overrightarrow{\delta}}{\delta\Psi(-p)} \Gamma_{I,\Lambda} \\
&\quad \left. + K((p+k)/\Lambda) \text{Tr} \Psi(p) \frac{\overleftarrow{\delta}}{\delta\psi(p+k)} - K(p/\Lambda) \text{Tr} \frac{\overrightarrow{\delta}}{\delta\bar{\psi}(-p)} \bar{\Psi}(-p-k) \right], \quad (\text{D}\cdot 12)
\end{aligned}$$

where

$$\begin{cases} \mathcal{A}_\mu(k) &= [A_\mu]_\Lambda(k), \\ \Psi(p) &= [\psi]_\Lambda(p), \\ \bar{\Psi}(-p) &= [\bar{\psi}]_\Lambda(-p). \end{cases} \quad (\text{D}\cdot 13)$$

We next consider the quantum master operator

$$\bar{\Sigma}_\Lambda[\phi, \phi^*] \equiv \int_p K(p/\Lambda) \left[\frac{\delta\bar{S}_\Lambda}{\delta\phi(p)} \frac{\overrightarrow{\delta}}{\delta\phi^*(-p)} \bar{S}_\Lambda + \frac{\overrightarrow{\delta}}{\delta\phi^*(-p)} \frac{\delta\bar{S}_\Lambda}{\delta\phi(p)} \right]. \quad (\text{D}\cdot 14)$$

The antifield ϕ^* is a classical external field, and it does not affect the Legendre transformation between $\bar{S}_\Lambda[\phi, \phi^*]$ and $\bar{\Gamma}_\Lambda[\Phi, \phi^*]$. The Legendre transformation is given by

$$\bar{\Gamma}_\Lambda[\Phi, \phi^*] = \bar{S}_\Lambda[\phi, \phi^*] + \int_p \frac{p^2 + m^2}{1 - K} \left(\frac{1}{2K} \phi(p) \phi(-p) - \Phi(p) \phi(-p) \right), \quad (\text{D}\cdot 15)$$

where

$$\Phi(p) = \frac{1}{K}\phi(p) + \frac{1-K}{p^2+m^2} \frac{\delta \bar{S}_\Lambda[\phi, \phi^*]}{\delta \phi(-p)}. \quad (\text{D}\cdot 16)$$

From the general relation (2.73), we obtain

$$\frac{\vec{\delta}}{\delta \phi^*(-p)} \bar{S}_\Lambda[\phi, \phi^*] = \frac{\vec{\delta}}{\delta \phi^*(-p)} \bar{\Gamma}_\Lambda[\Phi, \phi^*]. \quad (\text{D}\cdot 17)$$

Hence, the 1PI quantum master operator is given by

$$\begin{aligned} \bar{\Sigma}_\Lambda^{1PI}[\Phi, \phi^*] = \int_p \left[\left\{ -(p^2+m^2)\Phi(-p) + \frac{\delta \bar{\Gamma}_{I,\Lambda}[\Phi, \phi^*]}{\delta \Phi(p)} \right\} \frac{\vec{\delta}}{\delta \phi^*(-p)} \bar{\Gamma}_\Lambda[\Phi, \phi^*] \right. \\ \left. - \int_q \left(\bar{\Gamma}_\Lambda^{(2)} \right)^{-1}(q, -p) R_\Lambda(p) \frac{\vec{\delta}}{\delta \phi^*(-p)} \frac{\delta \bar{\Gamma}_\Lambda[\Phi, \phi^*]}{\delta \Phi(q)} \right]. \end{aligned} \quad (\text{D}\cdot 18)$$

Since

$$\bar{\Sigma}_\Lambda[\phi, \phi^*] = \bar{\Sigma}_\Lambda^{1PI}[\Phi, \phi^*] \quad (\text{D}\cdot 19)$$

by construction, $\bar{\Sigma}_\Lambda$ and $\bar{\Sigma}_\Lambda^{1PI}$ share the same asymptotic behavior. For YM theories,

$$\bar{\Sigma}_\Lambda^{1PI}[\Phi, \phi^*] = 0 \quad (\text{D}\cdot 20)$$

is called the **modified Slavnov-Taylor (ST) identity**, first obtained by Ellwanger.⁴⁰⁾

To summarize, for the effective average action, the WT identity is given by

$$\Sigma_\Lambda^{1PI}[\Phi] = 0, \quad (\text{D}\cdot 21)$$

where $\Sigma_\Lambda^{1PI}[\Phi]$ is given by (D-8), and the quantum master equation is given by the modified ST identity

$$\bar{\Sigma}_\Lambda^{1PI}[\Phi, \phi^*] = 0, \quad (\text{D}\cdot 22)$$

where $\bar{\Sigma}_\Lambda^{1PI}[\Phi, \phi^*]$ is given by (D-18).

In the limit $\Lambda \rightarrow 0+$, the effective average action becomes the effective action (the generating functional of the 1PI correlation functions):

$$\begin{cases} \lim_{\Lambda \rightarrow 0+} \Gamma_\Lambda[\Phi] &= \Gamma[\Phi], \\ \lim_{\Lambda \rightarrow 0+} \bar{\Gamma}_\Lambda[\Phi, \phi^*] &= \bar{\Gamma}[\Phi, \phi^*]. \end{cases} \quad (\text{D}\cdot 23)$$

Since $K(p/\Lambda)$ vanishes in this limit, the identities (D-21) & (D-22) reduce to

$$\begin{cases} \int_p \frac{\delta \Gamma[\Phi]}{\delta \Phi(p)} \mathcal{O}_{\Lambda=0}^{1PI}[\Phi](p) = 0, \\ \int_p \frac{\delta \bar{\Gamma}[\Phi, \phi^*]}{\delta \Phi(p)} \frac{\vec{\delta}}{\delta \phi^*(-p)} \bar{\Gamma}[\Phi, \phi^*] = 0. \end{cases} \quad (\text{D}\cdot 24)$$

The latter is the Zinn-Justin equation,¹²⁵⁾ familiar from YM theories.

Appendix E

—— $O(N)$ linear sigma model ——

In this subsection we apply the antifield formalism to the $O(N)$ linear sigma model. The model is a typical example of theories with their symmetry realized linearly. In such cases, it should not be surprising that the WT identity $\Sigma_A = 0$ or the QME $\bar{\Sigma}_A = 0$ for the Wilson action boils down to the invariance of the action under the naïve linear transformation.^{*)} Though we do not discuss it here, the Wess-Zumino model with auxiliary fields has its supersymmetry linearly realized, and the supersymmetry of the Wilson action is invariant under the naïve linear supersymmetry transformation.^{(35), (93), (102)}

We consider a theory of N real scalar fields ϕ_i invariant under the following linear $O(N)$ transformation:

$$\delta\phi_i(p) = \xi^a T_{ij}^a \phi_j(p), \quad (i = 1, \dots, N) \quad (\text{E.1})$$

where ξ^a ($a = 1, \dots, N(N-1)/2$) is an infinitesimal constant. The N -by- N matrices T^a are real and antisymmetric, and they satisfy the commutation relation

$$[T^a, T^b] = -f^{abc} T^c, \quad (\text{E.2})$$

where the structure constants are real and completely antisymmetric, and satisfy the Jacobi identity:

$$f^{abd} f^{dce} + f^{bcd} f^{dae} + f^{cad} f^{dbe} = 0. \quad (\text{E.3})$$

The Wilson action is given as

$$S_A = S_{F,\Lambda} + S_{I,\Lambda}, \quad (\text{E.4})$$

where the free part is

$$S_{F,\Lambda} \equiv \frac{1}{2} \int_p \phi_i(p) \phi_i(-p) \frac{p^2 + m^2}{K(p/\Lambda)}. \quad (\text{E.5})$$

Using the composite operator corresponding to the elementary field

$$[\phi_i]_\Lambda(p) \equiv \phi_i(p) + \frac{1 - K(p/\Lambda)}{p^2 + m^2} \frac{\delta S_{I,\Lambda}}{\delta \phi_i(-p)}, \quad (\text{E.6})$$

the WT composite operator is given by

$$\Sigma_A[\phi] \equiv \int_p K(p/\Lambda) \left[\frac{\delta S_A}{\delta \phi_i(p)} \xi^a T_{ij}^a [\phi_j]_\Lambda(p) + \frac{\delta}{\delta \phi_i(p)} (\xi^a T_{ij}^a [\phi_j]_\Lambda(p)) \right]. \quad (\text{E.7})$$

The WT identity

$$\Sigma_A = 0 \quad (\text{E.8})$$

^{*)} In preparing this appendix, we have benefited from the discussions with Drs. K. Ülker and L. Akant.

is equivalent to the cutoff independent identities

$$\sum_{j=1}^n \left\langle \phi_{i_1}(p_1) \cdots \xi^a T_{ij}^a \phi_k(p_j) \cdots \phi_{i_n}(p_n) \right\rangle^\infty = 0 \quad (\text{E}\cdot 9)$$

for any n .

We now wish to show that the WT identity (E·8) is equivalent to the naïve identity:

$$\int_p \frac{\delta S_{I,\Lambda}}{\delta \phi_i(p)} \xi^a T_{ij}^a \phi_j(p) = 0. \quad (\text{E}\cdot 10)$$

This is done in two steps. First we show the vanishing of the jacobian:

$$\begin{aligned} & \int_p K(p/\Lambda) \frac{\delta}{\delta \phi_i(p)} (\xi^a T_{ij}^a [\phi_j]_\Lambda(p)) \\ &= \int_p K(p/\Lambda) \xi^a T_{ij}^a \left(\delta_{ij} (2\pi)^4 \delta(0) + \frac{1 - K(p/\Lambda)}{p^2 + m^2} \frac{\delta^2 S_{I,\Lambda}}{\delta \phi_i(p) \delta \phi_j(-p)} \right) \\ &= 0. \end{aligned} \quad (\text{E}\cdot 11)$$

This vanishes because of the antisymmetry of T^a . Second we consider

$$\begin{aligned} & \int_p K(p/\Lambda) \frac{\delta S_\Lambda}{\delta \phi_i(p)} \xi^a T_{ij}^a [\phi_j]_\Lambda(p) \\ &= \int_p K(p/\Lambda) \left[-(p^2 + m^2) \frac{1}{K(p/\Lambda)} \phi_i(-p) + \frac{\delta S_{I,\Lambda}}{\delta \phi_i(p)} \right] \\ & \quad \times \xi^a T_{ij}^a \left[\phi_j(p) + \frac{1 - K(p/\Lambda)}{p^2 + m^2} \frac{\delta S_{I,\Lambda}}{\delta \phi_j(-p)} \right] \\ &= \int_p K(p/\Lambda) \left[-\frac{1 - K(p/\Lambda)}{K(p/\Lambda)} \phi_i(-p) \frac{\delta S_{I,\Lambda}}{\delta \phi_j(-p)} + \frac{\delta S_{I,\Lambda}}{\delta \phi_i(p)} \phi_j(p) \right] \xi^a T_{ij}^a \\ &= - \int_p \phi_i(-p) \frac{\delta S_{I,\Lambda}}{\delta \phi_j(-p)} \xi^a T_{ij}^a, \end{aligned} \quad (\text{E}\cdot 12)$$

where we have used the antisymmetry of T^a again. Hence, the WT identity is equivalent to (E·10).

We now promote ξ^a to anticommuting ghost fields, constant in space. We then introduce the fermionic antifields $\phi_i^*(p)$ for $\phi_i(p)$ and bosonic constant antifields ξ^{*a} for ξ^a . We define the action by

$$\begin{aligned} \bar{S}_\Lambda \equiv S_{F,\Lambda} &+ \int_p \phi_i^*(-p) \xi^a T_{ij}^a \phi_j(p) + \frac{1}{2} \int_p \phi_i^*(p) \xi^a T_{ij}^a \phi_k^* \xi^b T_{kj}^b \frac{1 - K(p/\Lambda)}{p^2 + m^2} \\ &+ S_{I,\Lambda} [\phi_i^{sh}(p)] - \frac{1}{2} \xi^{a*} f^{abc} \xi^b \xi^c, \end{aligned} \quad (\text{E}\cdot 13)$$

where the interaction action is the same as before, except that the fields are shifted by antifields:

$$\phi_i^{sh}(p) \equiv \phi_i(p) + \phi_j^*(p) \xi^a T_{ji}^a \frac{1 - K(p/\Lambda)}{p^2 + m^2}. \quad (\text{E}\cdot 14)$$

$\bar{S}_\Lambda$ is constructed so that it satisfies the same Polchinski differential equation as S_Λ . The last Λ -independent term generates the O(N) transformation of the ghosts:

$$\delta \xi^a = \frac{\partial \bar{S}_\Lambda}{\partial \xi^{a*}} = -\frac{1}{2} f^{abc} \xi^b \xi^c. \quad (\text{E}\cdot 15)$$

The QM operator is defined by

$$\begin{aligned} \bar{\Sigma}_\Lambda[\phi, \phi^*; \xi, \xi^*] \equiv & \int_p K(p/\Lambda) \left\{ \frac{\delta \bar{S}_\Lambda}{\delta \phi_i(p)} \cdot \frac{\vec{\delta}}{\delta \phi_i^*(-p)} \bar{S}_\Lambda + \frac{\delta}{\delta \phi_i(p)} \frac{\vec{\delta}}{\delta \phi_i^*(-p)} \bar{S}_\Lambda \right\} \\ & + \frac{\partial \bar{S}_\Lambda}{\partial \xi^{c*}} \frac{\vec{\partial}}{\partial \xi^c} \bar{S}_\Lambda. \end{aligned} \quad (\text{E}\cdot 16)$$

We wish to verify that the QME

$$\bar{\Sigma}_\Lambda[\phi, \phi^*; \xi, \xi^*] = 0 \quad (\text{E}\cdot 17)$$

is equivalent to the WT identity (E·8), and hence equivalent to the naïve identity (E·10). Verification is straightforward but takes several steps.

1. We first compute

$$\begin{aligned} \frac{\vec{\delta}}{\delta \phi_i^*(-p)} \bar{S}_\Lambda &= \xi^a T_{ij}^a \left[\phi_j(p) + \phi_k^*(p) \xi^b T_{kj}^b \frac{1 - K(p/\Lambda)}{p^2 + m^2} \right. \\ &\quad \left. + \frac{1 - K(p/\Lambda)}{p^2 + m^2} \frac{\delta S_{I,\Lambda}[\phi^{sh}]}{\delta \phi_j^{sh}(-p)} \right] \\ &= \xi^a T_{ij}^a [\phi_j]_{\Lambda}(p) \Big|_{\phi \rightarrow \phi^{sh}}. \end{aligned} \quad (\text{E}\cdot 18)$$

Hence,

$$\frac{\vec{\delta}}{\delta \phi_i^*(-p)} \frac{\delta \bar{S}_\Lambda}{\delta \phi_i(p)} = \xi^a T_{ij}^a \frac{1 - K(p/\Lambda)}{p^2 + m^2} \frac{\delta^2 S_{I,\Lambda}[\phi^{sh}]}{\delta \phi_j^{sh}(-p) \delta \phi_i^{sh}(p)}, \quad (\text{E}\cdot 19)$$

and therefore

$$\int_p K(p/\Lambda) \frac{\vec{\delta}}{\delta \phi_i^*(-p)} \frac{\delta \bar{S}_\Lambda}{\delta \phi_i(p)} = 0 \quad (\text{E}\cdot 20)$$

by the antisymmetry of T^a .

2. We then compute

$$\begin{aligned} \frac{\delta \bar{S}_\Lambda}{\delta \phi_i(p)} &= -\frac{1}{K(p/\Lambda)} (p^2 + m^2) \phi_i(-p) + \phi_j^*(-p) \xi^a T_{ji}^a + \frac{\delta S_{I,\Lambda}[\phi^{sh}]}{\delta \phi_i^{sh}(p)} \\ &= -\frac{1}{K(p/\Lambda)} (p^2 + m^2) \left(\phi_i(-p) + \phi_j^*(-p) \xi^a T_{ji}^a \frac{1 - K(p/\Lambda)}{p^2 + m^2} \right) \\ &\quad + \frac{\delta S_{I,\Lambda}[\phi^{sh}]}{\delta \phi_i^{sh}(p)} + \frac{1}{K(p/\Lambda)} \phi_j^*(-p) \xi^a T_{ji}^a \\ &= \frac{\delta S_\Lambda[\phi^{sh}]}{\delta \phi_i^{sh}(p)} + \frac{1}{K(p/\Lambda)} \phi_j^*(-p) \xi^a T_{ji}^a. \end{aligned} \quad (\text{E}\cdot 21)$$

3. We therefore obtain

$$\begin{aligned}
& \int_p K(p/\Lambda) \frac{\delta \bar{S}_\Lambda}{\delta \phi_i(p)} \cdot \frac{\vec{\delta}}{\delta \phi_i^*(-p)} \bar{S}_\Lambda \\
&= \int_p K(p/\Lambda) \left[\frac{\delta S_\Lambda[\phi^{sh}]}{\delta \phi_i^{sh}(p)} + \frac{1}{K(p/\Lambda)} \phi_j^*(-p) \xi^a T_{ji}^a \right] \xi^b T_{ik}^b [\phi_k]_\Lambda(p) \Big|_{\phi \rightarrow \phi^{sh}} \\
&= \Sigma_\Lambda \Big|_{\phi \rightarrow \phi^{sh}} + \int_p \phi_j^*(-p) \xi^a T_{ji}^a \xi^b T_{jk}^b [\phi_k]_\Lambda(p) \Big|_{\phi \rightarrow \phi^{sh}}. \tag{E.22}
\end{aligned}$$

Hence,

$$\bar{\Sigma}_\Lambda = \Sigma_\Lambda \Big|_{\phi \rightarrow \phi^{sh}} + \int_p \phi_j^*(-p) \xi^a T_{ji}^a \xi^b T_{jk}^b [\phi_k]_\Lambda(p) \Big|_{\phi \rightarrow \phi^{sh}} + \frac{\partial \bar{S}_\Lambda}{\partial \xi^{c*}} \frac{\vec{\partial}}{\partial \xi^c} \bar{S}_\Lambda. \tag{E.23}$$

4. Using

$$\xi^a T_{ji}^a \xi^b T_{jk}^b = -\frac{1}{2} f^{abc} \xi^a \xi^b T_{jk}^c = \frac{\partial \bar{S}_\Lambda}{\partial \xi^{c*}} T_{jk}^c, \tag{E.24}$$

we obtain

$$\begin{aligned}
& \int_p \phi_j^*(-p) \xi^a T_{ji}^a \xi^b T_{ik}^b [\phi_k]_\Lambda(p) \Big|_{\phi \rightarrow \phi^{sh}} \\
&= \frac{\partial \bar{S}_\Lambda}{\partial \xi^{c*}} \int_p \phi_i^*(-p) T_{ij}^c \left[\phi_j(p) + \phi_k^*(p) \xi^d T_{kj}^d \frac{1 - K(p/\Lambda)}{p^2 + m^2} \right. \\
&\quad \left. + \frac{1 - K(p/\Lambda)}{p^2 + m^2} \frac{\delta S_{I,\Lambda}[\phi^{sh}]}{\delta \phi_j^{sh}(-p)} \right]. \tag{E.25}
\end{aligned}$$

5. We compute

$$\begin{aligned}
\frac{\vec{\partial}}{\partial \xi^c} \bar{S}_\Lambda &= - \int_p \phi_i^*(-p) T_{ij}^c \left[\phi_j(p) + \phi_k^*(p) \xi^d T_{kj}^d \frac{1 - K(p/\Lambda)}{p^2 + m^2} \right. \\
&\quad \left. + \frac{1 - K(p/\Lambda)}{p^2 + m^2} \frac{\delta S_{I,\Lambda}[\phi^{sh}]}{\delta \phi_j^{sh}(-p)} \right] + f^{abc} \xi^{a*} \xi^b. \tag{E.26}
\end{aligned}$$

Since

$$\frac{\partial S_\Lambda}{\partial \xi^{c*}} f^{abc} \xi^{a*} \xi^b = -\frac{1}{2} f^{cde} \xi^d \xi^e f^{abc} \xi^{a*} \xi^b = -\frac{1}{2} f^{cde} f^{abc} \xi^d \xi^e \xi^b \xi^{a*} = 0 \tag{E.27}$$

by the Jacobi identity, we obtain

$$\int_p \phi_j^*(-p) \xi^a T_{ji}^a \xi^b T_{jk}^b [\phi_k]_\Lambda(p) \Big|_{\phi \rightarrow \phi^{sh}} + \frac{\partial \bar{S}_\Lambda}{\partial \xi^{c*}} \frac{\vec{\partial}}{\partial \xi^c} \bar{S}_\Lambda = 0. \tag{E.28}$$

6. We finally obtain the desired relation

$$\bar{\Sigma}_\Lambda[\phi, \phi^*; \xi, \xi^*] = \Sigma_\Lambda[\phi] \Big|_{\phi \rightarrow \phi^{sh}}. \tag{E.29}$$

The last relation is already familiar from QED, for which we have derived (8·15), even though the abelian gauge symmetry of QED is not strictly linear in the sense used here.

To conclude, we have found that for the O(N) global symmetry, the following three equations are equivalent:

$$\delta S_\Lambda \equiv \int_p \frac{\delta S_\Lambda}{\delta \phi_i(p)} \xi^a T_{ij}^a \phi_j(p) = 0, \quad (\text{E} \cdot 30\text{a})$$

$$\Sigma_\Lambda[\phi] \equiv \int_p K(p/\Lambda) \frac{\delta S_\Lambda}{\delta \phi_i(p)} \xi^a T_{ij}^a [\phi_j]_\Lambda(p) = 0, \quad (\text{E} \cdot 30\text{b})$$

$$\bar{\Sigma}_\Lambda[\phi, \phi^*; \xi, \xi^*] \equiv \int_p K(p/\Lambda) \frac{\delta \bar{S}_\Lambda}{\delta \phi_i(p)} \frac{\vec{\delta}}{\delta \phi_i^*(-p)} \bar{S}_\Lambda + \frac{\partial \bar{S}_\Lambda}{\partial \xi^{c*}} \frac{\vec{\partial}}{\partial \xi^c} \bar{S}_\Lambda = 0. \quad (\text{E} \cdot 30\text{c})$$

The jacobian of the linear transformation is 1:

$$\int_p K(p/\Lambda) \xi^a T_{ij}^a \frac{\delta [\phi_j]_\Lambda(p)}{\delta \phi_i(p)} = \int_p K(p/\Lambda) \frac{\delta}{\delta \phi_i(p)} \frac{\vec{\delta}}{\delta \phi_i^*(-p)} \bar{S}_\Lambda = 0. \quad (\text{E} \cdot 31)$$

Appendix F

— Quick summary —

ERG

1. A cutoff function $K(p)$ and its log derivative $\Delta(p) \equiv -2p^2 \frac{d}{dp^2} K(p)$:

$$K(0) = 1, \quad K(p) \xrightarrow{p^2 \rightarrow \infty} 0. \quad (\text{F}\cdot 1)$$

We usually (not always) choose

$$K(p) = 1 \quad (p^2 < 1). \quad (\text{F}\cdot 2)$$

2. The interaction part $S_{I,\Lambda}$ of the Wilson action satisfies the Polchinski equation

$$-\Lambda \frac{\partial}{\partial \Lambda} S_{I,\Lambda} = \frac{1}{2} \int_p \frac{\Delta(p/\Lambda)}{p^2 + m^2} \left\{ \frac{\delta S_{I,\Lambda}}{\delta \phi(p)} \frac{\delta S_{I,\Lambda}}{\delta \phi(-p)} + \frac{\delta^2 S_{I,\Lambda}}{\delta \phi(p) \delta \phi(-p)} \right\}. \quad (\text{F}\cdot 3)$$

3. The full Wilson action

$$S_\Lambda \equiv -\frac{1}{2} \int_p \frac{p^2 + m^2}{K(p/\Lambda)} \phi(p) \phi(-p) + S_{I,\Lambda} \quad (\text{F}\cdot 4)$$

satisfies the ERG differential equation

$$\begin{aligned} -\Lambda \frac{\partial}{\partial \Lambda} S_\Lambda = \int_p \frac{\Delta(p/\Lambda)}{p^2 + m^2} & \left[\frac{p^2 + m^2}{K(p/\Lambda)} \phi(p) \frac{\delta S_\Lambda}{\delta \phi(p)} \right. \\ & \left. + \frac{1}{2} \left\{ \frac{\delta S_\Lambda}{\delta \phi(p)} \frac{\delta S_\Lambda}{\delta \phi(-p)} + \frac{\delta^2 S_\Lambda}{\delta \phi(p) \delta \phi(-p)} \right\} \right]. \end{aligned} \quad (\text{F}\cdot 5)$$

4. Under the initial condition $S_{I,\Lambda_0} = S_{I,B}$, the integral solution of the Polchinski equation is given by

$$\begin{aligned} \exp[S_{I,\Lambda}[\phi]] = \int [d\phi'] \exp \Big[& \\ -\frac{1}{2} \int_p \frac{p^2 + m^2}{K(p/\Lambda_0) - K(p/\Lambda)} & \phi'(-p) \phi'(p) + S_{I,B}[\phi + \phi'] \Big]. \end{aligned} \quad (\text{F}\cdot 6)$$

5. For renormalizable theories, $S_{I,\Lambda}$ can be determined by the Polchinski equation and its asymptotic behavior:

$$S_{I,\Lambda} \xrightarrow{\Lambda \rightarrow \infty} S_{I,\Lambda}^{asympt}. \quad (\text{F}\cdot 7)$$

For the ϕ^4 theory in $D = 4$, we choose

$$\begin{aligned} S_{I,\Lambda}^{asympt} = \int d^4x \Big[& (\Lambda^2 a_2 (\ln \Lambda/\mu) + m^2 b_2 (\ln \Lambda/\mu)) \frac{1}{2} \phi^2 \\ & + c_2 (\ln \Lambda/\mu) \frac{1}{2} \partial_\mu \phi \partial_\mu \phi + a_4 (\ln \Lambda/\mu) \frac{1}{4!} \phi^4 \Big], \end{aligned}$$

where the coefficients a_2, b_2, c_2, a_4 are determined by the Polchinski equation up to

$$b_2(0), \quad c_2(0), \quad a_4(0), \quad (\text{F}\cdot 8)$$

which constitute the renormalized parameters of the theory. A particularly simple choice is

$$b_2(0) = c_2(0) = 0, \quad a_4(0) = -\lambda. \quad (\text{F}\cdot 9)$$

6. No loss of information along the ERG flow: the correlation functions

$$\begin{cases} \langle \phi(p)\phi(-p) \rangle^\infty & \equiv \frac{1}{K(p/\Lambda)^2} \langle \phi(p)\phi(-p) \rangle_{S_\Lambda} + \frac{1-1/K(p/\Lambda)}{p^2+m^2}, \\ \langle \phi(p_1)\cdots\phi(p_n) \rangle^\infty & \equiv \prod_{i=1}^n \frac{1}{K(p_i/\Lambda)} \cdot \langle \phi(p_1)\cdots\phi(p_n) \rangle_{S_\Lambda}, \quad (n > 2) \end{cases} \quad (\text{F}\cdot 10)$$

are independent of Λ .

7. $\mathcal{O}_\Lambda$ is a composite operator if $S_\Lambda + \epsilon \mathcal{O}_\Lambda$ satisfies the ERG differential equation to first order in ϵ , or equivalently if $\mathcal{O}_\Lambda$ satisfies the differential equation

$$\begin{aligned} -\Lambda \frac{\partial}{\partial \Lambda} \mathcal{O} &= \int_p \frac{\Delta(p/\Lambda)}{p^2+m^2} \left[\left\{ \frac{p^2+m^2}{K(p/\Lambda)} \phi(p) + \frac{\delta S_\Lambda}{\delta \phi(-p)} \right\} \frac{\delta}{\delta \phi(p)} \right. \\ &\quad \left. + \frac{1}{2} \frac{\delta^2}{\delta \phi(p) \delta \phi(-p)} \right] \mathcal{O}_\Lambda \\ &= \int_p \frac{\Delta(p/\Lambda)}{p^2+m^2} \left[\frac{\delta S_{I,\Lambda}}{\delta \phi(-p)} \frac{\delta}{\delta \phi(p)} + \frac{1}{2} \frac{\delta^2}{\delta \phi(p) \delta \phi(-p)} \right] \mathcal{O}_\Lambda. \end{aligned} \quad (\text{F}\cdot 11)$$

Given an initial condition $\mathcal{O}_{\Lambda_0} = \mathcal{O}_B$, the integral formula for $\mathcal{O}_\Lambda$ is

$$\begin{aligned} \mathcal{O}_\Lambda[\phi] \exp[S_{I,\Lambda}[\phi]] &= \int [d\phi'] \mathcal{O}_B[\phi + \phi'] \\ &\times \exp \left[-\frac{1}{2} \int_p \frac{p^2+m^2}{K(p/\Lambda_0) - K(p/\Lambda)} \phi'(-p) \phi'(p) + S_{I,B}[\phi + \phi'] \right]. \end{aligned} \quad (\text{F}\cdot 12)$$

8. The Λ independent correlation functions of a composite operator:

$$\langle \mathcal{O}_\Lambda \phi(p_1) \cdots \phi(p_n) \rangle^\infty = \prod_{i=1}^n \frac{1}{K(p_i/\Lambda)} \cdot \langle \mathcal{O} \phi(p_1) \cdots \phi(p_n) \rangle_{S_\Lambda}. \quad (\text{F}\cdot 13)$$

9. Examples of composite operators

(a) $\int_p K(p/\Lambda) \frac{\delta S_\Lambda}{\delta \phi(-p)}$ (Equation of motion)

$$\int_p K(p/\Lambda) \left\langle \frac{\delta S_\Lambda}{\delta \phi(-p)} \phi(p_1) \cdots \phi(p_n) \right\rangle^\infty = - \sum_{i=1}^n \langle \cdots \phi(p_i + p) \cdots \rangle^\infty. \quad (\text{F}\cdot 14)$$

(b) $\frac{\delta S_\Lambda}{\delta J(-p)}$, where J is an external source.

$$\left\langle \frac{\delta S_\Lambda}{\delta J(-p)} \phi(p_1) \cdots \phi(p_n) \right\rangle^\infty = \frac{\delta}{\delta J(-p)} \langle \phi(p_1) \cdots \phi(p_n) \rangle^\infty. \quad (\text{F}\cdot 15)$$

(c) $[\phi]_\Lambda(p) \equiv \frac{1}{K(p/\Lambda)}\phi(p) + \frac{1-K(p/\Lambda)}{p^2+m^2} \frac{\delta S_\Lambda}{\delta\phi(-p)}$ has the asymptotic behavior

$$[\phi]_\Lambda(p) \xrightarrow{\Lambda \rightarrow \infty} \phi(p). \quad (\text{F}\cdot 16)$$

(d) If $\mathcal{O}_\Lambda$ is a composite operator, so is

$$\begin{aligned} A(p) &\equiv K(p/\Lambda) \left(\mathcal{O}_\Lambda \frac{\delta S_\Lambda}{\delta\phi(-p)} + \frac{\delta \mathcal{O}_\Lambda}{\delta\phi(-p)} \right) \\ &= \exp[-S_\Lambda] K(p/\Lambda) \frac{\delta}{\delta\phi(-p)} (\mathcal{O}_\Lambda \exp[S_\Lambda]). \end{aligned} \quad (\text{F}\cdot 17)$$

$$\langle A(p)\phi(p_1)\cdots\phi(p_n) \rangle^\infty = - \sum_{i=1}^n (2\pi)^D \delta^{(D)}(p_i + p) \left\langle \cdots \widehat{\phi(p_i)} \mathcal{O}_\Lambda \cdots \right\rangle^\infty. \quad (\text{F}\cdot 18)$$

(e) If $\mathcal{O}_\Lambda$ is a composite operator, so is

$$B(p) \equiv \frac{\delta}{\delta J(-p)} \mathcal{O}_\Lambda + \frac{\delta}{\delta J(-p)} S_\Lambda \cdot \mathcal{O} = \exp[-S_\Lambda] \frac{\delta}{\delta J(-p)} (\mathcal{O} \exp[S_\Lambda]). \quad (\text{F}\cdot 19)$$

$$\langle B(p)\phi(p_1)\cdots\phi(p_n) \rangle^\infty = \frac{\delta}{\delta J(-p)} \langle \mathcal{O} \phi(p_1)\cdots\phi(p_n) \rangle^\infty. \quad (\text{F}\cdot 20)$$

10. For the ϕ^4 theory in $D = 4$, the composite operator $[\phi^2/2](p)$ is determined by the differential equation and the asymptotic behavior

$$[\phi^2/2](p) \xrightarrow{\Lambda \rightarrow \infty} c(\ln \Lambda/\mu) \frac{1}{2} \int_q \phi(q)\phi(p-q), \quad (\text{F}\cdot 21)$$

where $c(0)$ is an arbitrary normalization constant.

Ward-Takahashi (WT) identity

1. Under the symmetry transformation of the field $\phi(p)$

$$\delta\phi(p) = K(p/\Lambda) \mathcal{O}_\Lambda(p), \quad (\text{F}\cdot 22)$$

where $\mathcal{O}_\Lambda(p)$ is a composite operator, the Wilson action changes by

$$\begin{aligned} \Sigma_\Lambda &\equiv e^{-S_\Lambda} \int_p K(p/\Lambda) \frac{\delta}{\delta\phi(p)} (\mathcal{O}_\Lambda(p) e^{S_\Lambda}) \\ &= \int_p K(p/\Lambda) \left(\frac{\delta S_\Lambda}{\delta\phi(p)} \mathcal{O}_\Lambda(p) + \frac{\delta \mathcal{O}_\Lambda(p)}{\delta\phi(p)} \right). \end{aligned} \quad (\text{F}\cdot 23)$$

The WT composite operator Σ_Λ is a composite operator of type (d).

2. The WT identity is given by

$$\Sigma_\Lambda = 0. \quad (\text{F}\cdot 24)$$

Since Σ_Λ is a composite operator,

$$\lim_{\Lambda \rightarrow \infty} \Sigma_\Lambda = 0 \quad (\text{F}\cdot 25)$$

guarantees the WT identity for all Λ .

3. The WT identity is equivalent to

$$\sum_{i=1}^n \langle \phi(p_1) \cdots \mathcal{O}(p_i) \cdots \phi(p_n) \rangle^\infty = 0. \quad (\text{F}\cdot 26)$$

Quantum master equation

1. We generalize the Wilson action S_Λ to $\bar{S}_\Lambda$ by introducing an antifield ϕ^* , conjugate to ϕ . ϕ^* has the opposite statistics to ϕ . ϕ^* is a classical external source that generates the transformation of ϕ .
2. The quantum master operator

$$\begin{aligned} \bar{\Sigma}_\Lambda &\equiv e^{-\bar{S}_\Lambda} \int_p K(p/\Lambda) \frac{\delta}{\delta \phi(p)} \frac{\overrightarrow{\delta}}{\delta \phi^*(-p)} e^{\bar{S}_\Lambda} \\ &= \int_p K(p/\Lambda) \left[\frac{\delta \bar{S}_\Lambda}{\delta \phi(p)} \frac{\overrightarrow{\delta}}{\delta \phi^*(-p)} \bar{S}_\Lambda + \frac{\delta}{\delta \phi(p)} \frac{\overrightarrow{\delta}}{\delta \phi^*(-p)} \bar{S}_\Lambda \right] \end{aligned}$$

is a composite operator of type (d), since $\frac{\overrightarrow{\delta}}{\delta \phi^*(-p)} \bar{S}_\Lambda$ is a composite operator of type (e).

3. The quantum master equation

$$\bar{\Sigma}_\Lambda = 0 \quad (\text{F}\cdot 27)$$

is satisfied, if

$$\lim_{\Lambda \rightarrow \infty} \bar{\Sigma}_\Lambda = 0. \quad (\text{F}\cdot 28)$$

4. The quantum master equation is equivalent to the WT identity

$$\sum_{i=1}^n \frac{\overrightarrow{\delta}}{\delta \phi^*(-p_i)} \left\langle \phi(p_1) \cdots \widehat{\phi(p_i)} \cdots \phi(p_n) \right\rangle^\infty = 0. \quad (\text{F}\cdot 29)$$

5. The Wilson action $\bar{S}_\Lambda + \epsilon \mathcal{O}_\Lambda$, deformed by the composite operator $\mathcal{O}_\Lambda$, satisfies the quantum master equation, if and only if

$$\delta_Q \mathcal{O}_\Lambda = 0, \quad (\text{F}\cdot 30)$$

where the BRST transformation is defined by

$$\delta_Q \equiv \int_p K(p/\Lambda) \left\{ \frac{\overrightarrow{\delta}}{\delta \phi^*(-p)} \bar{S}_\Lambda \cdot \frac{\delta}{\delta \phi(p)} + \frac{\delta \bar{S}_\Lambda}{\delta \phi(p)} \frac{\overrightarrow{\delta}}{\delta \phi^*(-p)} + \frac{\overrightarrow{\delta}}{\delta \phi^*(-p)} \frac{\delta}{\delta \phi(p)} \right\}. \quad (\text{F}\cdot 31)$$

6. For any composite operator $\mathcal{O}$, $\delta_Q \mathcal{O}$ is a composite operator if $\bar{\Sigma} = 0$. Then,

$$\langle \phi(p_1) \cdots \phi(p_n) \delta_Q \mathcal{O} \rangle^\infty = - \sum_{i=1}^n \frac{\overrightarrow{\delta}}{\delta \phi^*(-p_i)} \left\langle \phi(p_1) \cdots \widehat{\phi(p_i)} \cdots \phi(p_n) \mathcal{O} \right\rangle^\infty. \quad (\text{F}\cdot 32)$$

For example,

$$\begin{aligned}\mathcal{N} &\equiv -\delta_Q \int_p \phi^*(-p)[\phi](p) \\ &= \int_p \left[\phi^*(-p) \frac{\vec{\delta}}{\delta\phi^*(-p)} \bar{S}_\Lambda - K(p/\Lambda) \left\{ [\phi](p) \frac{\delta \bar{S}_\Lambda}{\delta\phi(p)} + \frac{\delta[\phi](p)}{\delta\phi(p)} \right\} \right]\end{aligned}\quad (\text{F}\cdot 33)$$

satisfies

$$\langle \mathcal{N} \phi(p_1) \cdots \phi(p_n) \rangle^\infty = \left(\int_p \phi^*(-p) \frac{\vec{\delta}}{\delta\phi^*(-p)} + n \right) \langle \phi(p_1) \cdots \phi(p_n) \rangle^\infty. \quad (\text{F}\cdot 34)$$

7. The algebraic identity

$$\delta_Q \bar{S}_\Lambda = 0 \quad (\text{F}\cdot 35)$$

even if $\bar{S}_\Lambda \neq 0$.

8. The nilpotency $\delta_Q^2 = 0$ needs the QME $\bar{S}_\Lambda = 0$.

Classical nature of the quantum master equation

1. Loop expansions (bars omitted; $\bar{S}_\Lambda \rightarrow S_\Lambda$, etc.)

$$S_\Lambda = \sum_{l=0}^{\infty} S_l(\Lambda), \quad \Sigma_\Lambda = \sum_{l=0}^{\infty} \Sigma_l(\Lambda), \quad \dots \quad (\text{F}\cdot 36)$$

Loop expansions of the asymptotic parts:

$$S_\Lambda^{asympt} = \sum_{l=0}^{\infty} S_l^{asympt}(\Lambda), \quad \Sigma_\Lambda^{asympt} = \sum_{l=0}^{\infty} \Sigma_l^{asympt}(\Lambda), \quad \dots \quad (\text{F}\cdot 37)$$

2. **induction hypothesis:** Given $S_{0,\dots,l-1}$ that satisfy $\Sigma_{0,\dots,l-1} = 0$, we wish to construct S_l so that

$$\Sigma_l = 0. \quad (\text{F}\cdot 38)$$

This condition is equivalent to

$$\Sigma_l^{asympt} = 0. \quad (\text{F}\cdot 39)$$

3. Tree level: we start from the classical action S_{cl} with the classical BRST invariance

$$\int_x \frac{\vec{\delta}}{\delta\phi^*(x)} S_{cl} \cdot \frac{\delta S_{cl}}{\delta\phi(x)} = 0. \quad (\text{F}\cdot 40)$$

We then choose

$$S_0^{asympt} = S_{cl} \quad (\text{F}\cdot 41)$$

so that

$$\Sigma_0^{asympt} = \Sigma_0 = 0. \quad (\text{F}\cdot 42)$$

4. Under the induction hypothesis, the ERG differential equation Σ_l

$$-\Lambda \frac{\partial}{\partial \Lambda} \Sigma_l = \int_p \frac{\Delta(p/\Lambda)}{p^2 + m^2} \frac{\delta S_0}{\delta \phi(-p)} \frac{\delta \Sigma_l}{\delta \phi(p)} \quad (\text{F}\cdot 43)$$

implies that Σ_l^{asympt} is independent of Λ .

5. $S_l(\Lambda)$ satisfies the differential equation

$$-\Lambda \frac{\partial}{\partial \Lambda} S_l = \frac{1}{2} \int_p \frac{\Delta(p/\Lambda)}{p^2 + m^2} \left[\sum_{k=0}^l \frac{\delta S_k}{\delta \phi(-p)} \frac{\delta S_{l-k}}{\delta \phi(p)} + \frac{\delta^2 S_{l-1}}{\delta \phi(p) \delta \phi(-p)} \right]. \quad (\text{F}\cdot 44)$$

We split S_l into two parts:

$$S_l(\Lambda) = S_{1,l} + S_{2,l}(\Lambda) \quad (\text{F}\cdot 45)$$

(a) $S_{1,l}$ is independent of Λ , not determined by the differential equation.

(b) $S_{2,l}(\Lambda)$ is a particular solution of the differential equation for S_l .

We wish to fine tune $S_{1,l}$ so that $\Sigma_l = 0$.

6. We split Σ_l into two parts:

$$\Sigma_l = \Sigma_{1,l} + \Sigma_{2,l}, \quad (\text{F}\cdot 46)$$

where

$$\begin{aligned} \Sigma_{1,l} &\equiv \int_p K(p/\Lambda) \left[\frac{\vec{\delta}}{\delta \phi^*(-p)} S_0 \cdot \frac{\delta}{\delta \phi(p)} + \frac{\delta S_0}{\delta \phi(p)} \frac{\vec{\delta}}{\delta \phi^*(-p)} \right] S_{1,l}, \\ \Sigma_{2,l} &\equiv \int_p K(p/\Lambda) \left[\frac{\vec{\delta}}{\delta \phi^*(-p)} S_0 \cdot \frac{\delta}{\delta \phi(p)} + \frac{\delta S_0}{\delta \phi(p)} \frac{\vec{\delta}}{\delta \phi^*(-p)} \right] S_{2,l} \\ &\quad + \int_p K(p/\Lambda) \left[\sum_{k=1}^{l-1} \frac{\vec{\delta}}{\delta \phi^*(-p)} S_k \cdot \frac{\delta S_{l-k}}{\delta \phi(p)} + \frac{\vec{\delta}}{\delta \phi^*(-p)} \frac{\delta}{\delta \phi(p)} S_{l-1} \right]. \end{aligned}$$

(a) $S_0^{asympt} = S_{cl}$ implies $\Sigma_{1,l}^{asympt}$ is Λ independent.

(b) Σ_l^{asympt} is Λ independent, and so is $\Sigma_{2,l}^{asympt}$.

7. $\Sigma_l = 0$ is equivalent to

$$\Sigma_{1,l}^{asympt} + \Sigma_{2,l}^{asympt} = 0. \quad (\text{F}\cdot 47)$$

8. The classical BRST transformation is defined by

$$\delta_{cl} \equiv \int_x \left[\frac{\vec{\delta}}{\delta \phi^*(x)} S_{cl} \cdot \frac{\delta}{\delta \phi(x)} + \frac{\delta S_{cl}}{\delta \phi(x)} \frac{\vec{\delta}}{\delta \phi^*(x)} \right]. \quad (\text{F}\cdot 48)$$

This is nilpotent:

$$\delta_{cl}^2 = 0. \quad (\text{F}\cdot 49)$$

9. Since

$$\Sigma_{1,l}^{asympt} = \delta_{cl} S_{1,l}, \quad (\text{F}\cdot 50)$$

we obtain

$$\delta_{cl} \Sigma_{1,l}^{asympt} = 0. \quad (\text{F}\cdot 51)$$

10. $\delta_Q \Sigma_A = 0$ and the induction hypothesis imply

$$\int_p K(p/\Lambda) \left[\frac{\vec{\delta}}{\delta\phi^*(-p)} S_0 \cdot \frac{\delta}{\delta\phi(p)} + \frac{\delta S_0}{\delta\phi(p)} \frac{\vec{\delta}}{\delta\phi^*(-p)} \right] \Sigma_l = 0. \quad (\text{F}\cdot 52)$$

Taking the asymptotic form, we obtain

$$\delta_{cl} \Sigma_l^{asympt} = 0. \quad (\text{F}\cdot 53)$$

Hence, $\delta_{cl} \Sigma_{1,l}^{asympt} = 0$ implies

$$\delta_{cl} \Sigma_{2,l}^{asympt} = 0. \quad (\text{F}\cdot 54)$$

11. $\Sigma_l = 0$ is equivalent to

$$\delta_{cl} S_{1,l} = -\Sigma_{2,l}^{asympt}. \quad (\text{F}\cdot 55)$$

12. Thus, we must prove the triviality of the cohomology of δ_{cl} :

$$\boxed{\delta_{cl} \Sigma_{2,l}^{asympt} = 0 \stackrel{?}{\implies} \Sigma_{2,l}^{asympt} = \delta_{cl}(-S_{1,l}).} \quad (\text{F}\cdot 56)$$

References

- 1) M. G. Alford, Phys. Lett. B **336** (1994), 237.
- 2) K.-I. Aoki, Int. J. Mod. Phys. **B14** (2000), 1249.
- 3) K.-I. Aoki, *Non-perturbative renormalization group approach to dynamical chiral symmetry breaking in gauge theories* in *Nagoya 1996, Perspectives of strong coupling gauge theories*, (1996) 171, hep-ph/9706204.
- 4) K.-I. Aoki, K.-I. Morikawa, W. Souma, J.-I. Sumi, and H. Terao, Prog. Theor. Phys. **95** (1996), 409.
- 5) K.-I. Aoki, K.-I. Morikawa, W. Souma, J.-I. Sumi, and H. Terao, Prog. Theor. Phys. **99** (1998), 451.
- 6) S. Arnone, C. Fusi, and K. Yoshida, J. High Energy Phys. **02** (1999), 022.
- 7) S. Arnone, Y. A. Kubyshin, T. R. Morris, and J. F. Tighe, Int. J. Mod. Phys. **A17** (2002), 2283.
- 8) S. Arnone, T. R. Morris, and O. J. Rosten, Eur. Phys. J. **C50** (2007), 467.
- 9) C. Bagnuls and C. Bervillier, Phys. Rep. **348** (2001), 91.
- 10) R. D. Ball and R. S. Thorne, Ann. Phys. **236** (1994), 117.
- 11) R. D. Ball, P. E. Haagensen, J. I. Latorre, and E. Moreno, Phys. Lett. B **347** (1995), 80.
- 12) G. Barnich, F. Brandt, and M. Henneaux, Commun. Math. Phys. **174** (1995), 57.
- 13) G. Barnich, F. Brandt, and M. Henneaux, Commun. Math. Phys. **174** (1995), 93.
- 14) G. Barnich, F. Brandt, and M. Henneaux, Phys. Rep. **338** (2000), 439.
- 15) G. Barnich and M. Henneaux, Phys. Rev. Lett. **72** (1994), 1588.
- 16) I. A. Batalin and G. A. Vilkovisky, Phys. Lett. B **102** (1981), 27.
- 17) C. Becchi, *On the construction of renormalized gauge theories using renormalization group techniques*, hep-th/9607188.
- 18) F. Benitez, J.-P. Blaizot, H. Chaté, B. Delamotte, R. Méndez-Galain, and N. Wschebor, arXiv:0901.0128 [cond-mat.stat-mech].
- 19) F. Benitez, R. Méndez Galain, and N. Wschebor, Phys. Rev. B **77** (2008), 024431.
- 20) J. Berges, N. Tetradis, and C. Wetterich, Phys. Rep. **363** (2002), 223.
- 21) C. Bervillier, B. Boisseau, and H. Giacomini, Nucl. Phys. B **801** (2008), 296.
- 22) C. Bervillier, B. Boisseau, and H. Giacomini, Nucl. Phys. B **789** (2008), 525.
- 23) A. Bilal, Ann. of Phys. **323** (2008), 2311.
- 24) M. Bonini, M. D'Attanasio, and G. Marchesini, Nucl. Phys. B **409** (1993), 441.
- 25) M. Bonini, M. D'Attanasio, and G. Marchesini, Phys. Lett. B **329** (1994), 249.
- 26) M. Bonini, M. D'Attanasio, and G. Marchesini, Nucl. Phys. B **421** (1994), 429.
- 27) M. Bonini, M. D'Attanasio, and G. Marchesini, Nucl. Phys. B **418** (1994), 81.
- 28) M. Bonini, M. D'Attanasio, and G. Marchesini, Nucl. Phys. B **437** (1995), 163.
- 29) M. Bonini, M. D'Attanasio, and G. Marchesini, Phys. Lett. B **346** (1995), 87.
- 30) M. Bonini, M. D'Attanasio, and G. Marchesini, Nucl. Phys. B **444** (1995), 602.
- 31) M. Bonini, G. Marchesini, and M. Simionato, Nucl. Phys. B **483** (1997), 475.
- 32) M. Bonini and E. Tricarico, Nucl. Phys. B **585** (2000), 253.
- 33) M. Bonini and E. Tricarico, Nucl. Phys. B **606** (2001), 231.
- 34) M. Bonini and F. Vian, Nucl. Phys. B **511** (1998), 479.
- 35) M. Bonini and F. Vian, Nucl. Phys. B **532** (1998), 473.
- 36) F. Brandt, M. Henneaux, and A. Wilch, Phys. Lett. B **387** (1996), 320.
- 37) F. Brandt, M. Henneaux, and A. Wilch, Nucl. Phys. B **510** (1998), 640.
- 38) F. De Jonghe, hep-th/9403143.
- 39) B. Delamotte, cond-mat/0702365v1 [cond-mat.stat-mech].
- 40) U. Ellwanger, Phys. Lett. B **335** (1994), 364.
- 41) U. Ellwanger, M. Hirsch, and A. Weber, Z. Phys. C **69** **1996** (687).
- 42) M. E. Fisher, Rev. Mod. Phys. **70** (1998), 653.
- 43) F. Freire, D. F. Litim, and J. M. Pawłowski, Int. J. Mod. Phys. **A16** (2001), 2035.
- 44) F. Freire and C. Wetterich, Phys. Lett. B **380** (1996), 337.
- 45) F. Freire, D. F. Litim, and J. M. Pawłowski, Phys. Lett. B **495** (2000), 256.
- 46) H. Gies, hep-ph/0611146.
- 47) P. H. Ginsparg and K. G. Wilson, Phys. Rev. D **25** (1982), 2649.
- 48) G. R. Golner, hep-th/9801124.
- 49) J. Gomis, J. Paris, and S. Samuel, Phys. Rep. **259** (1995), 1.
- 50) A. Hasenfratz and P. Hasenfratz, Nucl. Phys. B **270** (1986), 687.

- 51) M. Henneaux and C. Teitelboim, *Quantization of Gauge Systems* (Princeton University Press, 1992).
- 52) T. Higashi, K. Higashijima, and E. Itou, Prog. Theor. Phys. Suppl. No. 164 (2007), 103.
- 53) T. Higashi, E. Itou, and T. Kugo, Prog. Theor. Phys. **118** (2007), 1115.
- 54) K. Higashijima and E. Itou, Prog. Theor. Phys. **108** (2002), 737.
- 55) K. Higashijima and E. Itou, Prog. Theor. Phys. **110** (2003), 563.
- 56) K. Higashijima and E. Itou, hep-th/0511300v1.
- 57) J. Hughes and J. Liu, Nucl. Phys. B **307** (1988), 183.
- 58) I. Ichinose and K. Nagao, Chin. J. Phys. **38** (2000), 671.
- 59) Y. Igarashi, M. Ishikake, K. Itoh, H. Sawanaka, and H. So, J. of Phys. **A39** (2006), 8023.
- 60) Y. Igarashi, K. Itoh, and H. So, Int. J. Mod. Phys. **A16** (2001), 2047.
- 61) Y. Igarashi, K. Itoh, and H. So, Prog. Theor. Phys. **104** (2000), 1053.
- 62) Y. Igarashi, K. Itoh, and H. So, Phys. Lett. B **479** (2000), 336.
- 63) Y. Igarashi, K. Itoh, and H. So, Prog. Theor. Phys. **106** (2001), 149.
- 64) Y. Igarashi, K. Itoh, and H. So, Phys. Lett. B **526** (2002), 164.
- 65) Y. Igarashi, K. Itoh, and H. Sonoda, Prog. Theor. Phys. **118** (2007), 121.
- 66) Y. Igarashi, K. Itoh, and H. Sonoda, Prog. Theor. Phys. **120** (2008), 1017.
- 67) Y. Igarashi, K. Itoh, and H. Sonoda, *Quantum master equation for yang-mills theory in the exact renormalization group in preparation.*
- 68) Y. Igarashi, H. So, and N. Ukita, Phys. Lett. B **535** (2002), 363.
- 69) Y. Igarashi, H. So, and N. Ukita, Nucl. Phys. B **640** (2002), 95.
- 70) G. Keller and C. Kopper, Commun. Math. Phys. **148** (1992), 445.
- 71) G. Keller and C. Kopper, Commun. Math. Phys. **153** (1993), 245.
- 72) G. Keller, C. Kopper, and M. Salmhofer, Helv. Phys. Acta **65** (1992), 32.
- 73) T. Kugo and S. Uehara, Nucl. Phys. B **197** (1982), 378.
- 74) O. Lauscher and M. Reuter, *Quantum Einstein gravity: Towards an asymptotically safe field theory of gravity*, volume 721 (Springer Berlin / Heidelberg, 2007).
- 75) D. F. Litim, Phys. Rev. D **64** (2001), 105007.
- 76) D. F. Litim, arXiv:0810.3675 [hep-th].
- 77) D. F. Litim and J. M. Pawłowski, hep-th/9901063.
- 78) D. F. Litim and J. M. Pawłowski, Phys. Lett. B **546** (2002), 279.
- 79) M. Lüscher, Phys. Lett. B **428** (1998), 342.
- 80) A. Margaritis, G. Odor, and A. Patkos, Z. Phys. **C39** (1988), 109.
- 81) T. R. Morris, Int. J. Mod. Phys. **A9** (1994), 2411.
- 82) T. R. Morris, Prog. Theor. Phys. Suppl. No. 131 (1998), 395.
- 83) T. R. Morris, Nucl. Phys. B **573** (2000), 97.
- 84) T. R. Morris, J. High Energy Phys. **12** (2000), 012.
- 85) T. R. Morris, J. High Energy Phys. **07** (2005), 027.
- 86) T. R. Morris and O. J. Rosten, Phys. Rev. D **73** (2006), 065003.
- 87) T. R. Morris and O. J. Rosten, J. of Phys. **A39** (2006), 11657.
- 88) J. F. Nicoll and T. S. Chang, Phys. Lett. A **62** (1977), 287.
- 89) J. F. Nicoll, T. S. Chang, and H. E. Stanley, Phys. Rev. Lett. **33** (1974), 540.
- 90) J. M. Pawłowski, Ann. of Phys. **322** (2007), 2831.
- 91) M. Pernici and M. Raciti, Nucl. Phys. B **531** (1998), 560.
- 92) M. Pernici, M. Raciti, and F. Riva, Nucl. Phys. B **520** (1998), 469.
- 93) M. Pernici, M. Raciti, and F. Riva, Phys. Lett. B **440** (1998), 305.
- 94) J. Polchinski, Nucl. Phys. B **231** (1984), 269.
- 95) J. Polonyi, Central Eur. J. Phys. **1** (2003), 1.
- 96) M. Reuter and C. Wetterich, Nucl. Phys. B **417** (1994), 181.
- 97) M. Reuter and C. Wetterich, Nucl. Phys. B **427** (1994), 291.
- 98) M. Reuter and F. Saueressig, arXiv:0708.1317 [hep-th].
- 99) O. J. Rosten, Int. J. Mod. Phys. **A21** (2006), 4627.
- 100) O. J. Rosten, J. of Phys. **A39** (2006), 8699.
- 101) O. J. Rosten, Phys. Rev. D **74** (2006), 125006.
- 102) O. J. Rosten, arXiv:0808.2150 [hep-th].
- 103) M. Salmhofer and C. Honerkamp, Prog. Theor. Phys. **105** (2001), 1.
- 104) N. Seiberg, Phys. Lett. B **318** (1993), 469.
- 105) R. Shankar, Rev. Mod. Phys. **66** (1994), 129.

- 106) H. Sonoda, Phys. Rev. D **67** (2003), 065011.
- 107) H. Sonoda, J. of Phys. **A40** (2007), 9675.
- 108) H. Sonoda, J. of Phys. **A40** (2007), 5733.
- 109) H. Sonoda, arXiv:0710.1662.
- 110) H. Sonoda and K. Ulker, Prog. Theor. Phys. **120** (2008), 197.
- 111) G. 't Hooft, Nucl. Phys. B **61** (1973), 455.
- 112) G. 't Hooft and M. J. G. Veltman, Nucl. Phys. B **44** (1972), 189.
- 113) N. Tetradis and C. Wetterich, Nucl. Phys. B **422** (1994), 541.
- 114) W. Troost, P. van Nieuwenhuizen, and A. Van Proeyen, Nucl. Phys. B **333** (1990), 727.
- 115) B. J. Warr, Ann. of Phys. **183** (1988), 1.
- 116) B. J. Warr, Ann. of Phys. **183** (1988), 59.
- 117) F. J. Wegner and A. Houghton, Phys. Rev. A **8** (1973), 401.
- 118) S. Weinberg, Phys. Rev. D **8** (1973), 3497.
- 119) S. Weinberg, Phys. Rev. Lett. **80** (1998), 3702.
- 120) J. Wess and B. Zumino, Phys. Lett. B **49** (1974), 52.
- 121) C. Wetterich, Nucl. Phys. B **352** (1991), 529.
- 122) C. Wetterich, Z. Phys. **C60** (1993), 461.
- 123) C. Wetterich, Phys. Lett. B **301** (1993), 90.
- 124) K. G. Wilson and J. B. Kogut, Phys. Rep. **12** (1974), 75.
- 125) J. Zinn-Justin, *Quantum field theory and critical phenomena, Fourth Edition* (Oxford Univ Press, 2002).